


RSC Advances



This article can be cited before page numbers have been issued, to do this please use: W. Zhao, P. Xie, Z. Bian, A. Zhou, H. Ge, B. Niu and Y. Ding, *RSC Adv.*, 2015, DOI: 10.1039/C5RA10763F.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Journal Name

COMMUNICATION

Generation of ArS-substituted Flavone Derivatives Using Aryl Thiols as Sulfenylating Agents

 Received 00th January 20xx,
Accepted 00th January 20xx

 Wannian Zhao,^a Ping Xie,^b Zhaogang Bian,^a Aihua Zhou,^{a*} Haibo Ge,^{c*} Ben Niu,^a and Yingcai Ding^a

DOI: 10.1039/x0xx00000x

www.rsc.org/

The regioselective and metal-free sulfonylation of flavones was achieved under aerobic conditions with ammonium iodide salt as an inducer instead of traditional iodine/oxidant combinations. This method enables the generation of various ArS-substituted flavone derivatives in good to excellent yields under environmentally friendly conditions, which significantly enriches current flavone chemistry.


The construction of C-S bond is a very important transformation in organic synthesis,¹ because C-S bond is widely present in natural products and drug molecules.² Therefore, many researchers have devoted their efforts on developing highly efficient, regioselective, and environmentally friendly C-S bond formation methods.³ Traditionally, C(sp³)-S bond can be constructed easily by the condensation of an alkyl halide with a metal thiolate.⁴ Recently, transition metal catalyzed C(sp²)-S bond formation were successfully developed via cross couplings between aryl halides and various sulfonylating reagents such as thiols (Scheme 1),⁵ sulfonyl chlorides,⁶ disulfides,⁷ sodium sulfonates,⁸ and sulfonyl hydrazides.⁹ However, these processes require the use of prefunctionalized halides and expensive catalysts.¹⁰

Recently, transition metal-catalyzed C-S bond formation via direct C-H bond functionalization has risen as an efficient alternative method, which allows direct conversion of C-H into C-S bonds. Although this method is highly efficient and atom-economic, these reactions still suffer from the high loading of transition metal catalysts, additives and harsh reaction conditions sometimes.¹¹

Very recently, metal-free iodine-induced sulfonylation method to construct C(sp²)-S bond was successfully developed (Scheme 1).¹² This method doesn't need the use of any toxic and transitional metal catalysts, and proceeded well under environmentally friendly reaction conditions, generating thioesters in good yields. Despite

this significant progress, the development of a new metal-free sulfonylation method is still highly desirable.

Herein, we developed a new and regioselective sulfonylation method in which a clean and colorless ammonium iodide salt was used as an reaction inducer instead of direct usage of purple iodine. In this study, air was used as the oxidant instead of pure oxygen and some oxidants (Scheme 1)¹²; and cheap aryl thiols were used as sulfonylating agents. To the best of our knowledge, there are no such reports to date. In this paper, this ammonium iodide-induced sulfonylation method was directly applied to electron-rich flavones to generate flavone derivatives which may be potentially valuable in drug discovery.



Scheme 1 Previously reported sulfonylation methods with arylthiols as sulfonylating agents.

To find the suitable reaction conditions for the ammonium iodide-induced sulfonylation between flavones and aryl thiol, flavone **1a** and thiol **2a** were used as the representative reactants. Different catalysts/inducers, oxidants, and solvents were screened

^aPharmacy School, Jiangsu University, Xuefu Road 301, Zhenjiang, Jiangsu 212013, People's Republic of China, Email: ahz@ujs.edu.cn.

^bScientific Information Research Institute, Jiangsu University (Library), Xuefu Road 301, Zhenjiang, Jiangsu 212013, People's Republic of China.

^cDepartment of Chemistry and Chemical Biology, Indiana University Purdue University Indianapolis, Indianapolis IN 46202 (USA), Email: geh@iupui.edu


† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/c000000x/

under different reaction temperature (Table 1). First, CuCl_2 and FeCl_3 were used as catalysts, and TBHP (*tert*-butyl hydroperoxide, 70 wt % in water) was employed as an oxidant. Both reactions gave a less than 5% yield or trace amount of expected product **3a** (entries 1 and 2) at 80 °C in CH_3CN . Using CuI as a catalyst with TBHP also provided a less than 5% yield of **3a** (entry 3) in CH_3CN . The combination of KI and TBHP only produced a 20% yield of **3a** (entry 4). When I_2 was employed with TBHP in CH_3CN , 45% of **3a** was observed (entry 5). The TBAI/TBHP combination in CH_3CN didn't afford **3a** at all (entry 6), while the combination of NH_4I /TBHP in CH_3CN afforded a 53% yield of **3a** (entry 7). The combination of I_2 /TBHP or I_2 /DTBP afforded **3a** in 70% and 65% yields, respectively (entries 8 and 9). When the reaction temperature was raised to 135 °C, the combination of NH_4I /TBHP in CH_3CN gave a 78% yield (entry 10). By converting NH_4I into I_2 , the reaction afforded 75% yield of **3a** (entry 11). Using DMF instead of CH_3CN afforded 83% yield of **3a** (entry 12). Surprisingly, when NH_4I was used without TBHP, the reaction also gave an 86% yield of **3a** (entry 13). Decreasing the amount of NH_4I from 2.0 to 1.2 equivalents led to 64% yield (entry 14). Using CH_3CN or THF as a solvent generated 55% yield or trace amount of product (entries 15 and 16). When toluene was used as a solvent, 73% yield of **3a** (entry 17) was obtained. No presence of

NH_4I in DMF didn't produce any product (entry 18). Using H_2O_2 instead of TBHP as an oxidant in the presence of NH_4I afforded **3a** a 65% yield (entry 19). While the combination of I_2 / H_2O_2 gave a 70% yield of **3a** (entry 20). After the above screening, the suitable conditions selected for the coupling of flavone and aryl thiol are: flavone (1.0 equiv.), thiols (1.2 equiv.), NH_4I (2.0 equiv.), DMF is used as a solvent and temperature: 135°C.


After suitable reaction conditions have been obtained, different aryl thiols were reacted with different flavones which were synthesized based on existing literature¹³. From Table 2, it can be found that all flavones with electron-donating functions gave better yields, while flavones with electron-deficient functions gave a little bit lower yields. Based on ^1H and ^{13}C -NMR spectra of all products, it was found that -SAr was regioselectively added to the α -position of


Table 1 Screening of reaction conditions^a

					
Entry	Catalyst /Inducer	Reagent	Temp. (°C)	Solvent	Yield ^b (%)
1	CuCl_2	TBHP	80	CH_3CN	5
2	FeCl_3	TBHP	80	CH_3CN	trace
3	CuI	TBHP	80	CH_3CN	<5
4	KI	TBHP	80	CH_3CN	20
5	I_2	TBHP	80	CH_3CN	45
6	TBAI	TBHP	80	CH_3CN	0
7	NH_4I	TBHP	80	CH_3CN	53
8	I_2	DCP	80	CH_3CN	70
9	I_2	DTBP	80	CH_3CN	65
10	NH_4I	TBHP	135	CH_3CN	78
11	I_2	TBHP	135	CH_3CN	75
12	NH_4I	TBHP	135	DMF	83
13	NH_4I	--	135	DMF	86
14	NH_4I^c	--	135	DMF	64
15	NH_4I	--	135	CH_3CN	55
16	NH_4I	--	135	THF	trace
17	NH_4I	--	135	Toluene	73
18	--	--	135	DMF	0
19	NH_4I	H_2O_2	135	DMF	65
20	I_2	H_2O_2	135	DMF	70

^a Reaction conditions: flavone (1 equiv.), aryl thiol (1.2 equiv.), NH_4I (2.0 equiv.), aqueous TBHP (70 wt % in water, 2.0 equiv.), metal catalyst (10 mol%), DMF (0.5 mL). ^b Isolated yields are based on reactant **1a**, the reaction was run for 15 hours with air. ^c NH_4I (1.2 equiv.)

Table 2 Ammonium iodide-mediated sulfonylation reactions using thiols as sulfonylating agents^a

			
Entry	Reactant 1	Product 3	Yield ^b (%)
1			85
2			86
3			80
4			82
5			84
6			81
7			83
8			81




^a Reaction conditions: flavone (1.0 equiv.), aryl thiol (1.2 equiv.), NH_4I (2.0 equiv.). ^b Isolated yields are based on reactant **1a**, the reaction was run for 15 hours.

the ketone function of flavones, and no β -substituted products were isolated.


To further explore how the substituents on α and β -position of flavones influence the regioselective sulfonylation, two methyl-substituted flavones were synthesized with methyl function on α and β -position of the flavone (see Scheme 2, entries 1 and 2). When the α -position reaction site was blocked, no any expected product was isolated, indicating that the regioselectivity of this sulfonylation is very good, and the reaction only happened on the α -position of flavone. When the methyl group on the β -position of flavone, a decreased sulfonylation yield was observed, possibly due to the steric effects caused by neighbouring methyl function.

To determine if a radical process is involved in this sulfonylation protocol, TEMPO (2,2,6,6-tetramethylpiperidine) and BHT (butylated hydroxytoluene) were used as radical scavengers in the reaction of producing **3a**. In the presence of TEMPO or BHT, **3a** was still produced in good yields (Scheme 2). This fact indicated that radical intermediate is not involved in the sulfonylation process. Based on the above results and existing literature,^{12a,14} a plausible nucleophilic substitution reaction mechanism is proposed below. At



Scheme 2 Control reactions and radical trapping experiments.

135 °C, NH_4I was split into NH_3 and HI , and the resulting HI was further oxidized by air to generate iodine. Thiophenol was then reacted with iodine to form electrophilic species Ar-S-I , which reacted further with electron-rich flavone **1** to give reaction intermediate A. After the loss of a proton from intermediate A, final product **3** was obtained.



Scheme 3 Proposed reaction mechanism

Conclusion

In summary, an efficient and regioselective ammonium iodide-induced sulfonylation of flavones is herein reported with different thiophenols as sulfonylating agents, generating various Ar-S- substituted flavone derivatives in good to excellent yields under metal-free conditions. This method has improved the current reported protocol represented by using iodine as a catalyst. This method enriches current flavone chemistry, making it a highly valuable and practical approach in pharmaceutical industry. Investigation on biological activities of flavone derivatives currently underway. The method is also quite suitable for compound library production.


Acknowledgment

Aihua Zhou acknowledge Jiangsu University for financial support (1281290006 and PAPD).

Notes and references

- 1 (a) M. Iwasaki, M. Iyanaga, Y. Tsuchiya, Y. Nishimura, W. Li, Z. Li and Y. Nishihara, *Chem. Eur. J.*, 2014, **20**, 2459; (b) W. Ali, S. Guin, S. K. Rout, A. Gogoi and B. K. Patel, *Adv. Synth. Catal.*, 2014, **356**, 3099; (c) H. Xi, B. Deng, Z. Zong, S. Lu and Z. Li, *Org. Lett.*, 2015, **17**, 1180; (d) C. Shen, P. Zhang, Q. Sun, S. Bai, T. S. A. Hor and X. Liu, *Chem. Soc. Rev.*, 2015, **44**, 291; (e) H. Wang, L. Wang, J. Shang, X. Li, H. Wang, J. Gui and A. Lei, *Chem. Commun.*, 2012, **48**, 76; (f) C.-P. Zhang and D. A. Vicic, *J. Am. Chem. Soc.*, 2012, **134**, 183.
- 2 (a) M. Halim, D. J. Yee and D. Sames, *J. Am. Chem. Soc.*, 2008, **130**, 14123; (b) M. S. C. Pedras, Q.-A. Zheng and S. Strelkov, *J. Agric. Food Chem.*, 2008, **56**, 9949; (c) X. Wang, L. Cui, N. Zhou, W. Zhu, R. Wang, X. Qian and Y. Xu, *Chem. Sci.*, 2013, **4**, 2936; (d) A. J. Wilson, J. K. Kerns, J. F. Callahan and C. J. Moody, *J. Med. Chem.*, 2013, **56**, 7463.
- 3 (a) J.-K. Qiu, W.-J. Hao, D.-C. Wang, P. Wei, J. Sun, B. Jiang and S.-J. Tu, *Chem. Commun.*, 2014, **50**, 14782; (b) L. D. Tran, I. Popov and O. Daugulis, *J. Am. Chem. Soc.*, 2012, **134**, 18237; (c) E. L. Tyson, M. S. Ament and T. P. Yoon, *J. Org. Chem.*, 2012, **78**, 2046; (d) F.-L. Yang, F.-X. Wang, T.-T. Wang, Y.-J. Wang and S.-K. Tian, *Chem. Commun.*, 2014, **50**, 2111; (e) Y. Yang, L. Tang, S. Zhang, X. Guo, Z. Zha and Z. Wang, *Green Chem.*, 2014, **16**, 4106; (f) M. Yoshimatsu, K. Ohta and N. Takahashi, *Chem. Eur. J.*, 2012, **18**, 15602.
- 4 (a) J. R. Campbell, *J. Org. Chem.*, 1964, **29**, 1830; (b) T. Migita, T. Shimizu, Y. Asami, J.-i. Shiobara, Y. Kato and M. Kosugi, *Bull. Chem. Soc. Jpn.*, 1980, **53**, 1385; (c) B. C. Ranu and T. Mandal, *J. Org. Chem.*, 2004, **69**, 5793.
- 5 (a) C.-K. Chen, Y.-W. Chen, C.-H. Lin, H.-P. Lin and C.-F. Lee, *Chem. Commun.*, 2010, **46**, 282; (b) A. Correa, O. G. Mancheño and C. Bolm, *Chem. Soc. Rev.*, 2008, **37**, 1108; (c) M. A. Fernández-Rodríguez, Q. Shen and J. F. Hartwig, *J. Am. Chem. Soc.*, 2006, **128**, 2180; (d) P. F. Larsson, A. Correa, M. Carril, P. O. Norrby and C. Bolm, *Angew. Chem.*, 2009, **121**, 5801; (e) M. Sayah and M. G. Organ, *Chem. Eur. J.*, 2011, **17**, 11719; (f) C. Yu, C. Zhang and X. Shi, *Eur. J. Org. Chem.*, 2012, **10**, 1953.
- 6 (a) M. Chen, Z.-T. Huang and Q.-Y. Zheng, *Chem. Commun.*, 2012, **48**, 11686; (b) O. Saidi, J. Marafie, A. E. Ledger, P. M. Liu, M. F. Mahon, G. Kociok-Köhn, M. K. Whittlesey and C. G. Frost, *J. Am. Chem. Soc.*, 2011, **133**, 19298; (c) D. Zhang, X. Cui, Q. Zhang and Y. Wu, *J. Org. Chem.*, 2015, **80**, 1517.
- 7 (a) W. Ge and Y. Wei, *Green Chem.*, 2012, **14**, 2066; (b) D. Huang, J. Chen, W. Dan, J. Ding, M. Liu and H. Wu, *Adv. Synth. Catal.*, 2012, **354**, 2123; (c) P. Sang, Z. Chen, J. Zou and Y. Zhang, *Green Chem.*, 2013, **15**, 2096; (d) S. Zhang, P. Qian, M. Zhang, M. Hu and J. Cheng, *J. Org. Chem.*, 2010, **75**, 6732.
- 8 (a) M. Yang, H. Shen, Y. Li, C. Shen and P. Zhang, *RSC Adv.*, 2014, **4**, 26295; (b) F. Xiao, H. Chen, H. Xie, S. Chen, L. Yang and G.-J. Deng, *Org. Lett.*, 2014, **16**, 50; (c) Y. Xu, X. Tang, W. Hu, W. Wu and H. Jiang, *Green Chem.*, 2014, **16**, 3720.
- 9 (a) X. Li, X. Xu, P. Hu, X. Xiao and C. Zhou, *J. Org. Chem.*, 2013, **78**, 7343; (b) X. Li, Y. Xu, W. Wu, C. Jiang, C. Qi and H. Jiang, *Chem. Eur. J.*, 2014, **20**, 7911; (c) M. Zhang, P. Xie, W. Zhao, B. Niu, W. Wu, Z. Bian, C. U. Pittman Jr and A. Zhou, *J. Org. Chem.*, 2015, **80**, 4176; (d) J. Zhang, Y. Shao, H. Wang, Q. Luo, J. Chen, D. Xu and X. Wan, *Org. Lett.*, 2014, **16**, 3312.
- 10 (a) C. Dai, Z. Xu, F. Huang, Z. Yu and Y.-F. Gao, *J. Org. Chem.*, 2012, **77**, 4414; (b) S. Mao, Y.-R. Gao, X.-Q. Zhu, D.-D. Guo and Y.-Q. Wang, *Org. Lett.*, 2015, **17**, 1692; (c) S. Tang, Y. Wu, W. Liao, R. Bai, C. Liu and A. Lei, *Chem. Commun.*, 2014, **50**, 4496; (d) Z. Wu, H. Song, X. Cui, C. Pi, W. Du and Y. Wu, *Org. Lett.*, 2013, **15**, 1270; (e) Y. Xu, P. Liu, S.-L. Li and P. Sun, *J. Org. Chem.*, 2015, **80**, 1269; (f) X. Zhao, E. Dimitrijevic and V. M. Dong, *J. Am. Chem. Soc.*, 2009, **131**, 3466.
- 11 (a) K. Inamoto, Y. Arai, K. Hiroya and T. Doi, *Chem. Commun.*, 2008, **43**, 5529; (b) K. Inamoto, C. Hasegawa, K. Hiroya and T. Doi, *Org. Lett.*, 2008, **10**, 5147; (c) L. L. Joyce and R. A. Batey, *Org. Lett.*, 2009, **11**, 2792; (d) R. Samanta and A. P. Antonchik, *Angew. Chem. Int. Ed.*, 2011, **50**, 5217; (e) C. Shen, H. Xia, H. Yan, X. Chen, S. Ranjit, X. Xie, D. Tan, R. Lee, Y. Yang and B. Xing, *Chem. Sci.*, 2012, **3**, 2388; (f) R. Xu, J.-P. Wan, H. Mao and Y. Pan, *J. Am. Chem. Soc.*, 2010, **132**, 15531; (g) Y. Yang, V. Hou, L. Qin, J. Du, H. Feng, B. Zhou and Y. Li, *Chem. Eur. J.*, 2014, **20**, 416; (h) J. Zhu, Z. Chen, H. Xie, S. Li and Y. Wu, *Org. Lett.*, 2010, **12**, 2434.
- 12 (a) M.-A. Hiebel and S. Berteina-Raboin, *Green Chem.*, 2015, **17**, 937; (b) Y. Liao, P. Jiang, S. Chen, H. Qi and G.-J. Deng, *Green Chem.*, 2013, **15**, 3302.
- 13 (a) D. Ameen and T. J. Snape, *Synthesis*, 2015, **47**, 141; (b) S.-H. Wang, Y. Wang, Y.-Y. Zhu, S.-J. Liu, J. Han, Y.-F. Zhou, D.-W. Li, D. Koirala and C. Hu, *Chem. Res. Chinese Univ.*, 2011, **27**, 54; (c) Y. Zheng, M. Zhao, Q. Qiao, H. Liu, H. Lang and Z. Xu, *Dyes and Pigments*, 2013, **98**, 367.
- 14 (a) F. L. Yang and S. K. Tian, *Angew. Chem. Int. Ed.*, 2013, **52**, 4929.

Graphic abstract:



The regioselective and metal-free sulfenylation of flavones was achieved under aerobic conditions with ammonium iodide as an inducer