


Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: E. Yoo, B. M. Crall, R. Balakrishna, S. S. Malladi, L. Fox, A. R. Hermanson and S. David, *Org. Biomol. Chem.*, 2013, DOI: 10.1039/C3OB40816G.



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

Structure-Activity Relationships in Toll-like Receptor 7 Agonistic 1*H*-Imidazo[4,5-*c*]pyridines**Euna Yoo, Breanna M. Crall, Rajalakshmi Balakrishna, Subbalakshmi S. Malladi, Lauren M. Fox, Alec R. Hermanson and Sunil A. David***

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

Engagement of TLR7 in plasmacytoid dendritic cells leads to the induction of IFN- α/β which plays essential functions in the control of adaptive immunity. We had previously examined structure-activity relationships (SAR) in TLR7/8-agonistic imidazoquinolines with a focus on substituents at the N^1 , C^2 , N^3 and N^4 positions, and we now report SAR on 1*H*-imidazo[4,5-*c*]pyridines. 1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine was found to be a pure TLR7-agonist with negligible activity on TLR8. Increase in potency was observed in N^6 -substituted analogues, especially in those compounds with electron-rich substituents. Direct aryl-aryl connections at C6 abrogated activity, but TLR7 agonism was reinstated in 6-benzyl and 6-phenethyl analogues. Consistent with the pure TLR7-agonistic behavior, prominent IFN- α induction in human PBMCs was observed with minimal proinflammatory cytokine induction. A benzologue of imidazoquinoline was also synthesized which showed substantial improvements in potency over the parent imidazopyridine. Distinct differences in N^6 -substituted analogues were observed with respect to IFN- α induction in human PBMCs on the one hand, and CD69 upregulation in lymphocytic subsets, on the other.

Introduction

Host responses to pathogens are mediated via highly coordinated mechanisms involving both the innate and adaptive limbs of the immune system. The innate immune system utilizes germline-encoded pattern recognition receptors (PRRs) to detect pathogen-associated molecular patterns (PAMPs) that are distinct and unique to the pathogen.¹⁻³ PRRs encompass a broad range of molecules⁴ that are secreted into the extracellular environment (such as the collectins,⁵ ficolins,⁶ pentraxins,⁷ alarmins⁸), exist in the cytosol (examples of which include the retinoic acid-inducible gene I-like receptors,⁹ and the nucleotide-binding domain and leucine-rich repeat-containing receptors¹⁰), or are present on membranes.

Important among the transmembrane PRRs include the Toll-like receptors¹¹ (TLRs) which are either expressed on the plasma membrane or in the endolysosomal compartments.^{1,12} At least 10 functional TLRs are encoded in the human genome, each with an extracellular domain having leucine-rich repeats (LRR) and a cytosolic domain called the Toll/IL-1 receptor (TIR) domain.¹³ The ligands for these receptors are highly conserved microbial molecules such as lipopolysaccharides (LPS) (recognized by TLR4), lipopeptides (TLR2 in combination with TLR1 or TLR6), flagellin (TLR5), single stranded RNA (TLR7 and TLR8), double stranded RNA (TLR3), CpG motif-containing DNA (recognized by TLR9), and profilin present on uropathogenic bacteria (TLR11).^{14,15} TLR1, -2, -4, -5, and -6 recognize extracellular

stimuli, while TLR3, -7, -8 and -9 function within the endolysosomal compartment.¹³ The activation of TLRs by their cognate ligands leads to production of inflammatory cytokines, and up-regulation of major histocompatibility complex (MHC) molecules and co-stimulatory signals in antigen-presenting cells as well as activating natural killer (NK) cells (innate immune response), which leads to the priming and amplification of T-, and B-cell effector functions (adaptive immune responses).¹⁶⁻¹⁹

The Type I interferon (IFN) family in humans include approximately 20 IFN- α subtype genes in addition to individual genes encoding IFN- β , - κ , - ϵ and - ω ; these monomeric secreted proteins bind to a single IFN- α/β receptor, which is constitutively expressed in virtually all cell types.²⁰ Occupancy of TLR7²¹⁻²³ or TLR9^{24,25} in professional antigen-presenting cells (APCs), particularly plasmacytoid dendritic cells (pDCs), leads to the induction of IFN- α/β . Although the Type I IFNs are best known historically for their antiviral activities,²⁶ recent studies show that they have many essential functions in the control of adaptive immunity.² First, Type I IFNs promote cross-priming through direct stimulation of DCs, leading to specific CD8⁺ lymphocytic responses to soluble antigens.²⁷ Second, Type I IFNs potently enhance the primary antibody responses to soluble antigens, inducing sustained and durable humoral responses with appropriate isotype switching, as well as the induction of immunological memory.²⁸ B lymphocytes can differentiate into two distinct types of functionally polarized effectors: B-effector-

1-cells (Be-1), producing a Th-1-like cytokine pattern, or Be-2, characterized by a Th-2-like profile.^{29,30} It is of particular interest that recent reports suggest that IFN- α may serve as an initial trigger for Be-1-biased differentiation pattern.³¹ Third, Type I interferons secondarily induce Type II IFN (IFN- γ) secretion, also driving Th-1-biased adaptive immune responses.³² Type I IFN-inducing TLR ligands may therefore hold promise as vaccine adjuvants.

In an effort to identify optimal immunostimulatory chemotypes, we have screened representative members of virtually the entire compendium of known TLR agonists in a series of hierarchical assays including primary TLR-reporter assays, secondary indices of immune activation such as IFN- $\alpha/\beta/\gamma$ and cytokine induction, activation of lymphocytic subsets in whole human blood, and tertiary screens characterizing transcriptomal activation patterns.³³ In these assays, small-molecule agonists of TLR7 were uniquely immunostimulatory; they were potent inducers of Type I IFN and, unlike TLR-4, -5, or -8 agonists,³³ did not evoke dominant proinflammatory cytokine responses, suggesting that they may be effective, yet safe vaccine adjuvants, a premise that we have been actively exploring.³⁴ Small molecule TLR7 agonists are also being investigated as orally bioavailable, endogenous Type I IFN inducers for the management of chronic viral diseases,³⁵ especially hepatitis C and hepatitis B. Current therapeutic regimens for the therapy of hepatitis C and hepatitis B include parenteral IFN- α .³⁶ Clinical trials of TLR7 agonists for hematological malignancies are also currently underway.³⁷

The currently known small molecule agonists of TLR7 occupy a very small chemical space, and are represented by the 1*H*-imidazo[4,5-*c*]quinolines,³⁸ 8-hydroxy-³⁹⁻⁴¹ or 8-oxoadenines,⁴² and guanine nucleoside analogues.^{43,44} We had previously reported structure-activity relationships (SAR) in the imidazoquinolines with a focus on substituents at the *N*¹, *C*², *N*³ and *N*⁴ positions,^{45,46} and we had observed that relatively minor structural modifications at these positions yielded compounds with widely differing immunomodulatory properties.^{34,47-49} It was of interest, therefore, to extend our SAR studies to the quinoline ring system. We asked if a part-structure (imidazopyridine) or a benzologue (benzoimidazoquinoline) would alter the biological properties of the parent imidazoquinoline compound. Examination of the structures of 3M-003⁵⁰ and the 8-hydroxy- and 8-oxoadenines (Fig. 1) suggested that the quinoline system may be dispensable, and activity would be retained in imidazopyridines. Indeed, imidazopyridine derivatives with alkyl groups at *C*⁶ and *C*⁷ positions,⁵¹ hydroxyalkyl,⁵² oxime and hydroxylamine-bearing substituents⁵³ at *C*², and alkylsulfonamide substituents at the *N*¹ position⁵⁴ have been reported in the patent literature. Detailed activity profiles of these compounds, however, are not available, perhaps owing to the fact that the investigations of such compounds precede the discovery of the TLRs.

Incorporating substituents that we had previously determined to be optimal in the imidazoquinolines (*N*¹-benzyl and *C*²-butyl; IMDQ, Fig. 1), we embarked on the syntheses and biological evaluation of novel 1*H*-imidazo[4,5-*c*]pyridine analogues with modifications at the *N*¹- and *C*⁶ positions. The parent imidazopyridine compound, 1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]





Fig. 1 Structures of small molecule agonists of TLR7 represented by the 8-oxoadenine (SM360320, Ref. 42), 8-hydroxyadenines (CL264, Ref. 39), 1*H*-imidazo[4,5-*c*]quinolines 3M-003 (Ref. 49), 1-isobutyl-1*H*-imidazo[4,5-*c*]quinolin-4-amine (Imiquimod) and 1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]quinolin-4-amine (IMDQ, Ref. 45).


pyridin-4-amine, exhibited moderate TLR7-agonistic activity. *N*¹-acyl or -alkyl substitutions abrogated activity. The majority of *C*⁶ derivatives bearing aryl groups were also inactive, but analogues with *N*⁶-benzyl substituents gained TLR7-specific activity. Particular *N*⁶ substituents were found to augment TLR7-specific agonistic potency without compromising specificity at TLR7; consistent with their pure TLR7 activity (and undetectable TLR8 agonism), these compounds potently induced IFN- α in human peripheral blood mononuclear cells (PBMCs), upregulated CD69 in lymphocytic subsets, and yet showed very weak proinflammatory cytokine-inducing activities. Strong Type I IFN inducers, especially in conjunction with attenuated proinflammatory profiles are expected to be potently adjuvant without inducing prominent local or systemic inflammation.

Results and discussion

Our interest in exploring TLR7 agonists as vaccine adjuvants has been greatly reinforced by our observations that pure TLR7 agonists, unlike other TLR ligands, are potently immunostimulatory without prominently activating inflammatory programs in human whole blood model systems.⁵⁵ As mentioned earlier, the structures of 3M-003⁵⁰ and the 8-hydroxy- and 8-oxoadenines (Fig. 1), as well as patent literature suggested that the quinoline system may be dispensable, and activity would be retained in imidazopyridines. Our previous SAR studies on the imidazoquinolines had established that *N*¹-benzyl and *C*²-butyl substituents were optimal;⁴⁶ our point of departure in examining structure-activity relationships in the imidazopyridines consequently began with the evaluation of 1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine (**5**), following the synthetic strategy described earlier (Scheme 1).^{38,46} Compound **5**, itself a novel and unprecedented structure, was found to possess TLR7-specific agonistic activity (EC₅₀: 1.57 μ M, Fig. 2, Table 1), with negligible TLR8 activity. The potency of the lead TLR7-specific imidazoquinoline (1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]quinolin-4-amine, structure in Fig. 1) was 0.06 μ M (Fig. 2).⁴⁶




Scheme 1. Reagents: (i) BrNH_2 , NEt_3 , CH_2Cl_2 ; (ii) Zn , HCOONH_4 , MeOH ; (iii) a. $\text{C}_4\text{H}_9\text{COCl}$, NEt_3 , THF ; b. NaOH , EtOH ; (iv) *m*CPBA, CHCl_3 ; (v) a. Benzoyl isocyanate, CH_2Cl_2 ; b. NaOMe , MeOH .



Scheme 2. Reagents: (i) RCOCl , NEt_3 , CH_2Cl_2

5 Acylation (**6a**, **6b**, Scheme 2) of the $\text{C}^4\text{-NH}_2$ resulted in complete abrogation of activity (Table 1). C^6 -modified analogues were synthesized via an alternate route. Nitration of 4-amino-2-chloropyridine resulted, as expected, in a mixture of the 3- and 5-nitro intermediates **7a** and **7b**, which were taken forward to obtain the 4- and 6-chloroimidazopyridines **10a** and **10b** (Scheme 3). Excellent chromatographic separation of these advanced intermediates was possible. Pd-catalyzed C-N cross-coupling reactions using *n*-butylamine and benzylamine furnished the C^4 -*N*-alkylated analogues **11a** and **11b**, respectively (Scheme 3). A 4-butoxy analogue **11c** was also obtained by *ipso*-chloro displacement with 1-butanol. Compounds **11a-c** were, however, inactive (Table 1). We had envisaged utilizing the 6-chloroimidazopyridine intermediate **10b** for synthesizing C^6 -functionalized analogues. However, this intermediate exhibited unexpectedly low reactivity to displacement with nucleophiles or to Buchwald-Hartwig coupling reactions. As outlined in Scheme 4, we utilized 4-amino-2,6-dichloropyridine as the starting material and obtained **13** as a key intermediate which, upon reaction with *tert*-octylamine⁵⁶ provided exclusively the N^2 -



Scheme 3. Reagents: (i) a. H_2SO_4 , HNO_3 ; b. H_2SO_4 ; (ii) BnBr , NaH , THF ; (iii) Zn , HCOONH_4 , MeOH ; (iv) a. $\text{C}_4\text{H}_9\text{COCl}$, NEt_3 , THF ; b. NaOH , EtOH ; (v) For **11a** and **11b**, amines (*n*- BuNH_2 and BnNH_2 , respectively), $\text{Pd}_2(\text{dba})_3$, DavePhos, KOfBu , dioxane; For **11c**, BuOH , NaH , THF .






Fig. 2 TLR7 agonistic activity of imidazopyridine compounds. Data points represent means and standard deviations on quadruplicates.

alkylated intermediate **14**. The 6-chloro-*N*-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazopyridin-4-amine intermediate **16** was obtained without difficulty, and we were able to synthesize the N^6 -substituted analogues **19a-o** under conventional Buchwald-Hartwig conditions (Scheme 4). TLRs signal via ligand-induced dimerization, but since that the crystal structure of human TLR7 and of its ligand binding modes are as yet unknown, we utilized intermediate **16** in constructing ‘dimeric’ imidazopyridines (using *p*- and *m*-xylylenediamine, Scheme 5) to ascertain if such pre-organized dimeric ligands could yield high-potency agonists. For the C^6 -substituted compounds **23a-j** (Scheme 6), we observed mediocre yields in pilot Suzuki coupling reactions with the advanced intermediate **16**. We therefore exploited the electron-withdrawing resonance effect of the 3-nitro group in **14**. As expected, the classical Suzuki reaction on intermediate **14** using various aliphatic and aromatic boronic acids/boronate esters resulted in the intermediates **20a-j** (Scheme 6), which were further derivatized to obtain the desired C^6 alkyl/aryl substituted imidazopyridines **23a-j**.

The 6-chloro imidazopyridine, **17** (Scheme 4) was inactive (Table 1). Buchwald-Hartwig-derived N^6 -substituted analogues **19a-q**, however, showed a distinctive SAR. Compound **19a** with a free NH_2 at C^6 , obtained by coupling the *tert*-octylamine and subsequent *N*-dealkylation with HCl (Scheme 4, Table 1) displayed TLR7-specific agonism with a potency comparable to that of the parent C^6 -unsubstituted compound **5**.


Table 1. EC₅₀ values of compounds in human TLR7-specific reporter gene assay

Compound Number	R ¹	R ²	hTLR7-agonistic Activity (μM)
5	H	H	1.57
6a	COCH ₃	H	Inactive ^a
6b	COC ₃ H ₇	H	Inactive
11a	C ₄ H ₉	H	Inactive
11b	CH ₂ C ₆ H ₅	H	Inactive
17	H	Cl	Inactive
19a	H	NH ₂	1.25
19b	H	NHC ₄ H ₉	0.34
19c	H	NHC ₇ H ₁₅	0.76
19d	H		0.32
19e	H	NHC ₆ H ₅	1.72
19f	H	NHCH ₂ C ₆ H ₅	0.21
19g	H		0.075
19h	H		0.61
19i	H		1.04
19j	H		0.64
19k	H		0.075
19l	H		0.25
19m	H		0.25

Table 1. (contd.)

Compound Number	R ¹	R ²	hTLR7-agonistic Activity (μM)
19n	H		Inactive
19o	H		1.09
19p	H		0.26
19q	H		0.37
19r	H		1.08
19s	H		Inactive
23a	H	C ₄ H ₉	0.28
23b	H	C ₆ H ₅	Inactive
23c	H		Inactive
23d	H		Inactive
23e	H		Inactive
23f	H		Inactive
23g	H		0.28
23h	H		1.80
23i	H		Inactive
23j	H		0.57
30	Benzologue (Scheme 7)		0.22

^aInactive: no activity was detected up to a concentration of 500 μg/mL.



Scheme 4. Reagents: (i) a. H_2SO_4 , HNO_3 ; b. H_2SO_4 ; (ii) BnBr , NaH , THF ; (iii) *t*-Octylamine, NEt_3 , CH_2Cl_2 ; (iv) Zn , HCOONH_4 , MeOH ; (v) a. $\text{C}_4\text{H}_9\text{COCl}$, NEt_3 , THF ; b. NaOH , EtOH ; (vi) RNH_2 , $\text{Pd}_2(\text{dba})_3$, DavePhos , KOtBu , dioxane; (vii) HCl .

50

Modest gains in potency were obtained in analogues with short aliphatic substituents with N^6 -butyl (**19b**) and N^6 -cyclohexylmethyl (**19d**), but potency diminished in the N^6 -heptyl analogue (**19c**). The N^6 -phenyl-substituted compound **19e** was marginally weaker than **5**; however, the potency of the N^6 -benzyl analogue **19f** was ~ 7.6 times that of **5** (Table 1, Fig. 2), triggering a detailed SAR investigation on various aryl substituents at N^6 . Both steric and electronic effects appear to play a role in governing TLR7-agonistic potency, since the biphenylmethyl-substituted compound **19o** was active, whereas the naphthylmethyl analogue **19n** was quiescent; to a first approximation, electron-rich N^6 substituents appear to be preferred, with the methoxybenzyl derivatives (**19g** and **19h**) and the pyridinylmethyl compounds (**19l** and **19m**) being marginally more active than the trifluoromethyl- (**19i**) or chloro- (**19j**) substituted analogues. Compounds **19p** and **19q** were also active in primary screens, with EC_{50} values of 0.26 and 0.37 μM , respectively (Table 1). In the C^6 -alkyl or -aryl analogues (Scheme 6), the SAR appeared more stringent. Whereas the C^6 -butyl compound **23a** was more active than **5**, direct aryl-aryl connections at C^6 (**23b-f**) abrogated activity, but TLR7 agonistic properties were restored in the 6-benzyl (**23g**) and 6-phenethyl analogues (**23j**). Taken together with activity data of compounds of the **19** series, we surmised that rotational constraints about the






Fig. 3 Dose-response profiles of TLR7 agonistic activity of compounds **5** and **30**. Data points represent means and standard deviations on quadruplicates.


C^6 -aryl groups may hinder TLR7 occupancy. Unlike TLR2, TLR3, TLR4,⁵⁷ and TLR5⁵⁸ for which crystal structures are available as complexes with their cognate ligands, a detailed structural characterization of the mode of binding of TLR7 ligands is not yet available to guide structure-based design of small molecule agonists of TLR7, necessitating classical SAR approaches to refine successive iterations of ligand design.



Scheme 5. Reagents: (i) $\text{Pd}_2(\text{dba})_3$, DavePhos , KOtBu , dioxane; (ii) HCl .



Scheme 6. Reagents: (i) For **20a-f** and **20j**, R-Boronic acid, Pd(dppf)Cl₂, Cs₂CO₃, dioxane; For **20c**, 4-Cyanophenyl boronic acid, Pd(dppf)Cl₂, Cs₂CO₃, dioxane; For **20g-i**, R-Boronic acid pinacol ester, Pd(dppf)Cl₂, Cs₂CO₃, dioxane; (ii) Zn, HCOONH₄, MeOH; (iii) a. C₄H₉COCl, NEt₃, THF; b. NaOH, EtOH; (iv) HCl.



Scheme 7. Reagents: (i) a. HCl, HON=CHCH₂NO₂, H₂O; b. (CH₃CO)₂O, CH₃COOK; (ii) POCl₃; (iii) BnNH₂, NEt₃, CH₂Cl₂; (iv) Zn, HCOONH₄, MeOH; (v) a. C₄H₉COCl, NEt₃, THF; b. NaOH, EtOH; (vi) mCPBA, CH₂Cl₂, CHCl₃, MeOH; (vii) a. Benzoyl isocyanate, CH₂Cl₂; b. NaOMe, MeOH.

5

The benzologue **30** was synthesized as shown in Scheme 7. It showed substantial improvements in potency over the parent imidazopyridine **5** (Fig. 3, Table 1), but the two most potent compounds in the entire series as adjudged by primary screens were the N⁶-(4-methoxybenzyl) and N⁶-(furan-2-ylmethyl) analogues (**19g** and **19k**, respectively), both of which were approximately twenty-fold more potent than **5** (Fig. 2, Table 1).

We chose the nine most active compounds (**19b**, **19d**, **19f-g**, **19k-m**, **19p** and **23a**) for evaluation in secondary screens using IFN- α and cytokine release in human PBMCs. We used, as reference compounds, imiquimod, a known TLR7 agonist,^{59,60} as




Fig. 4 Dose-response profiles of Type I interferon (IFN- α) and proinflammatory cytokine (IL-8, IL-1 β , and TNF- α) induction by selected imidazopyridine (and reference) compounds. Representative data from three independent experiments are presented.

well as CL075^{61,62} (2-propylthiazolo[4,5-*c*]quinolin-4-amine), a predominantly TLR8-active agonist with an EC₅₀ of 1.32 μ M in hTLR8 assays.⁶³ Given that the imidazopyridine compounds are pure TLR7 agonists, we expected to find prominent IFN- α induction,³³ and this was indeed the case, with **19p**, **19m** and **19k** being the most potent (EC₅₀: 0.3 μ M, 0.4 μ M and 0.7 μ M, respectively; Fig. 4). CL075 was among the least potent in IFN- α induction (EC₅₀: 2.6 μ M; Fig. 4), and as expected for a TLR8 agonist, CL075 was dramatically more active in inducing




Fig. 5 CD69 upregulation in human natural killer (NK), cytokine-induced killer (CIK) and nominal B lymphocytes by select imidazopyridine (and reference) compounds.

proinflammatory cytokines such as TNF- α , IL-1 β , and IL-8 (Fig. 4). We do not yet understand the basis for the slight discrepancy between rank-order potency in primary screens ($19k \approx 19g > 19f > 19p$; Fig. 2, Table 1) vis-à-vis IFN- α -inducing potency in human PBMCs ($19p \approx 19m > 19k$; Fig. 4), and we surmise that analogues with more basic C6 substituents may allow for higher endolysosomal partitioning. The dose-response profiles show characteristic biphasic responses (dose-dependent activation, followed by apparent suppression) as we had previously observed in several chemotypes. We verified that the apparent suppression was not due to cytotoxicity using LDH release and mitochondrial redox-based assays.

We had previously shown that of all the various classes of innate immune stimuli, TLR7 agonists were extraordinarily immunostimulatory, stimulating virtually all subsets of lymphocytes (assessed by quantifying CD69 expression), and yet without inducing dominant proinflammatory cytokine responses,³³ and we wished to confirm the rank-order potency observed in IFN- α induction assays described above. We observed considerable dissociation between Type I IFN induction on the one hand (Fig. 4), and CD69 upregulation in lymphocytic subsets on the other (Fig. 5). Whereas the subset of active compounds induced IFN- α with similar potencies (EC_{50} values between 0.3-2 μ M; Fig. 4), pronounced differences were observed in CD69 expression in natural killer, cytokine-induced killer and B lymphocytic subsets with **19p** being as active as the reference TLR7 agonist IMDQ, and **19d** showing virtually no activity (Fig. 5). Possible mechanisms underlying the differential activity in these two compounds are being investigated.

Conclusions

These findings raise the possibility of utilizing these compounds in selectively targeting Type I IFN induction versus lymphocytic activation, and are being explored in greater detail. The potential advantages of strong Type I IFN inducers as candidate vaccine adjuvants have been discussed earlier. Such compounds, especially in conjunction with attenuated proinflammatory cytokines, are expected to be potently adjuvantic without inducing prominent local or systemic inflammation. As mentioned earlier, the prominent Type I IFN inducing abilities of the imidazopyridines may also find utility as an alternative therapeutic strategy to address disease states wherein systemic IFN- α is of proven benefit. A clear delineation of structural features that confer TLR specificity not only charts a rational course for the development of effective, yet safe vaccine adjuvants, but also provides tools to understand innate immune function in greater detail.

Experimental

Materials and equipment. All of the solvents and reagents used were obtained commercially and used as such unless noted otherwise. Moisture- or air-sensitive reactions were conducted under nitrogen atmosphere in oven-dried (120 $^{\circ}$ C) glass apparatus. The solvents were removed under reduced pressure using standard rotary evaporators. Flash column chromatography was carried out using RediSep R_f 'Gold' high performance silica columns on CombiFlash R_f instrument unless otherwise mentioned, while thin-layer chromatography was carried out on silica gel (200 μ m) CCM pre-coated aluminum sheets. Purity for all final compounds was confirmed to be greater than 97% by LC-MS using a Zorbax Eclipse Plus 4.6 mm x 150 mm, 5 μ m analytical reverse phase C_{18} column with H₂O-isopropanol or H₂O-CH₃CN gradients and an Agilent 6520 ESI-QTOF Accurate Mass spectrometer (mass accuracy of 5 ppm) operating in the positive ion (or negative ion, as appropriate) acquisition mode.

Synthesis of compound 1: N-Benzyl-3-nitropyridin-4-amine.

To a solution of 4-chloro-3-nitropyridine (1.0 g, 6.31 mmol) in 25 mL of CH₂Cl₂ were added triethylamine (1.32 mL, 9.47 mmol) and benzyl amine (0.83 mL, 7.57 mmol). The reaction mixture was refluxed for 18 h. The solvent was then evaporated under vacuum and H₂O was added to the residue. The solution was extracted with CH₂Cl₂ (3 × 20 mL), washed with water and dried over sodium sulfate. The solvent was evaporated and the residue was purified using silica gel column chromatography (5% MeOH/CH₂Cl₂) to obtain compound **1** as a yellow solid (1.4 g, 94%). ¹H NMR (500 MHz, CDCl₃) δ 9.22 (s, 1H), 8.53 (s, 1H), 8.25 (dd, *J* = 6.1, 0.6 Hz, 1H), 7.41 – 7.35 (m, 2H), 7.32 (dd, *J* = 7.2, 5.3 Hz, 3H), 6.69 (d, *J* = 6.2 Hz, 1H), 4.56 (d, *J* = 5.7 Hz, 2H). ¹³C NMR (126 MHz, CDCl₃) δ 153.38, 149.06, 148.68, 135.96, 130.04, 129.21, 128.22, 127.10, 108.25, 46.85. MS (ESI) calculated for C₁₂H₁₂N₃O₂, *m/z* 230.0924, found 230.0949 (M + H)⁺.

Synthesis of compound 2: N⁴-Benzylpyridine-3,4-diamine. To a solution of compound **1** (1.0 g, 4.36 mmol) in 40 mL of MeOH were added zinc dust (1.4 g, 21.8 mmol) and ammonium formate (1.4 g, 21.8 mmol). The reaction mixture was stirred at room temperature for 10 min and filtered through celite. Then the solvent was evaporated and the residue was dissolved in water. This was extracted with EtOAc (3 × 20 mL), washed with water and dried over sodium sulfate. The solvent was evaporated under vacuum to obtain the compound **2** (0.8 g, 92%). ¹H NMR (500 MHz, CDCl₃) δ 7.81 (d, *J* = 5.4 Hz, 1H), 7.77 (s, 1H), 7.29 – 7.19 (m, 5H), 6.36 (d, *J* = 5.4 Hz, 1H), 4.87 (s, 1H), 4.28 (d, *J* = 5.4 Hz, 2H), 3.28 (s, 2H). ¹³C NMR (126 MHz, CDCl₃) δ 144.52, 143.51, 138.22, 137.53, 128.96, 128.79, 127.55, 127.48, 105.29, 47.22. MS (ESI) calculated for C₁₂H₁₄N₃, *m/z* 200.1182, found 200.1242 (M + H)⁺.

Synthesis of compound 3: 1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine. To a solution of compound **2** (400 mg, 2.00 mmol) in 20 mL of anhydrous THF were added triethylamine (0.29 mL, 2.10 mmol) and valeryl chloride (0.27 mL, 2.20 mmol). The reaction mixture was refluxed for 2 h. The solvent was then removed under vacuum, and the residue was dissolved in EtOAc and washed with water. The EtOAc fraction was dried using sodium sulfate and evaporated under vacuum to obtain the intermediate amide compound. This was dissolved in 20 mL of EtOH and NaOH (160 mg, 4.00 mmol) in 2 mL of H₂O was added. The reaction mixture was refluxed for 4 h. The solvent was then removed under vacuum, and the residue was dissolved in EtOAc and washed with water. The organic layer was dried using sodium sulfate and evaporated and purified using column chromatography (5% MeOH/CH₂Cl₂) to obtain the compound **3** (210 mg, 40%). ¹H NMR (500 MHz, CDCl₃) δ 9.04 (d, *J* = 0.6 Hz, 1H), 8.34 (d, *J* = 5.6 Hz, 1H), 7.34 – 7.28 (m, 3H), 7.14 (dd, *J* = 5.6, 1.0 Hz, 1H), 7.01 (dd, *J* = 7.7, 1.8 Hz, 2H), 5.33 (s, 2H), 2.86 – 2.81 (m, 2H), 1.82 (dt, *J* = 15.5, 7.7 Hz, 2H), 1.46 – 1.36 (m, 2H), 0.91 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 157.30, 142.14, 142.00, 140.48, 140.01, 135.26, 129.28, 128.37, 126.27, 105.12, 47.25, 29.44, 27.38, 22.61, 13.85. MS (ESI) calculated for C₁₇H₂₀N₃, *m/z* 266.1652, found 266.1715 (M + H)⁺.

Synthesis of compound 4: 1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine 5-oxide. To a solution of compound **3** (210 mg, 0.79 mmol) in 15 mL of was added *m*-chloroperoxybenzoic acid (443 mg, 1.98 mmol), and the solution was refluxed at 45–50 °C for 1 h. The solvent was then removed and the residue was purified using column chromatography (10% MeOH/CH₂Cl₂) to obtain the *N*-oxide derivative (188 mg, 85%). ¹H NMR (500 MHz, CDCl₃) δ 8.71 (d, *J* = 1.3 Hz, 1H), 8.05 (dd, *J* = 7.0, 1.6 Hz, 1H), 7.36 – 7.30 (m, 3H), 7.01 (dd, *J* = 10.4, 4.3 Hz, 3H), 5.31 (s, 2H), 2.85 – 2.80 (m, 2H), 1.79 (dt, *J* = 15.4, 7.6 Hz, 2H), 1.44 – 1.34 (m, 2H), 0.90 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 160.72, 140.89, 134.42, 134.32, 133.94, 131.52, 129.48, 128.75, 126.24, 106.63, 47.75, 29.25, 27.49, 22.52, 13.78. MS (ESI) calculated for C₁₇H₂₀N₃O, *m/z* 282.1601, found 282.1612 (M + H)⁺.

Synthesis of compound 5: 1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine. To a solution of compound **4** (188 mg, 0.67 mol) in 15 mL of CH₂Cl₂ was added benzoyl isocyanate (197 mg, 1.34 mmol) and heated at 45 °C for 2 h. The solvent was then removed under vacuum, and the residue was dissolved in 15 mL of anhydrous MeOH, followed by the addition of excess sodium methoxide. The reaction mixture was then heated at 80 °C for 1 h. The solvent was removed under vacuum and the residue was purified using column chromatography (7% MeOH/CH₂Cl₂) to obtain the compound **5** (56 mg, 30%). ¹H NMR (500 MHz, CDCl₃) δ 7.78 (d, *J* = 5.8 Hz, 1H), 7.34 – 7.28 (m, 3H), 7.03 (d, *J* = 6.4 Hz, 2H), 6.59 (d, *J* = 5.8 Hz, 1H), 5.27 (s, 2H), 5.15 (s, 2H), 2.83 – 2.75 (m, 2H), 1.72 (ddd, *J* = 13.0, 9.0, 7.7 Hz, 2H), 1.44 – 1.34 (m, 2H), 0.90 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 154.06, 151.00, 140.61, 140.41, 135.76, 129.21, 128.23, 126.33, 97.74, 47.52, 30.10, 27.42, 22.68, 13.89. HRMS (ESI) calculated for C₁₇H₂₁N₄, *m/z* 281.1761, found 281.1795 (M + H)⁺.

Synthesis of compound 6a: N-(1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl)acetamide. To a solution of compound **5** (30 mg, 0.11 mmol) in 2 mL of CH₂Cl₂ were added triethylamine (17 μL, 0.12 mmol) and acetyl chloride (8 μL, 0.11 mmol). The reaction mixture was stirred at room temperature for 3 h and purified using column chromatography (5% MeOH/CH₂Cl₂) to obtain the compound **6a** as white solid (6 mg, 16%). ¹H NMR (500 MHz, CDCl₃) δ 11.60 (s, 1H), 8.20 (s, 1H), 7.37 (dd, *J* = 7.7, 5.5 Hz, 3H), 7.20 (d, *J* = 5.6 Hz, 1H), 7.05 (dd, *J* = 6.3, 2.5 Hz, 2H), 5.46 (s, 2H), 3.03 – 2.94 (m, 2H), 2.55 (s, 3H), 1.85 (dt, *J* = 15.0, 7.6 Hz, 2H), 1.43 (dq, *J* = 14.6, 7.3 Hz, 2H), 0.92 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 141.48, 129.64, 129.40, 129.17, 126.30, 126.25, 48.47, 29.06, 24.87, 22.44, 13.60. HRMS (ESI) calculated for C₁₉H₂₃N₄O, *m/z* 323.1866, found 323.1911 (M + H)⁺.

110 Compounds **6b** was synthesized similarly as compound **6a**.

N-(1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-4-yl)butyramide (6b**).** Butyryl chloride was used as a reagent. (4 mg, 14 %). ¹H NMR (500 MHz, CDCl₃) δ 11.31 (s, 1H), 8.30 (d, *J* = 4.8 Hz, 1H), 7.38 – 7.34 (m, 3H), 7.27 (s, 1H), 7.04 (dd, *J* = 6.4, 2.6 Hz, 2H), 5.50 (s, 2H), 2.94 (t, *J* = 7.5 Hz, 2H), 2.78 (t, *J* = 7.4

Hz, 2H), 1.80 (dp, $J = 22.1, 7.5$ Hz, 4H), 1.41 (dt, $J = 14.7, 7.4$ Hz, 2H), 1.01 (t, $J = 7.4$ Hz, 3H), 0.91 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 174.32, 141.35, 133.50, 129.69, 129.16, 126.46, 103.38, 48.58, 39.27, 29.16, 27.45, 22.55, 18.38, 13.75. HRMS (ESI) calculated for $\text{C}_{21}\text{H}_{27}\text{N}_4\text{O}$, m/z 351.2179, found 351.2240 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 10a: 1-Benzyl-2-butyl-4-chloro-1H-imidazo[4,5-c]pyridine. 4-Amino-2-chloropyridine (2.0 g, 15.6 mmol) was taken in 20 mL of conc. H_2SO_4 in an ice-bath to which was added 10 mL of conc. HNO_3 slowly. The reaction mixture was gradually brought to room temperature and stirred for 1 h. The reaction was quenched by pouring the reaction mixture on ice. Ammonium hydroxide solution was slowly added until a pH of 3.0 was reached. A white solid was obtained which was filtered, washed with water, and dried. This (*N*-nitro)aminopyridine intermediate was taken up in 10 mL of conc. H_2SO_4 and the reaction solution was heated at 90 °C for 30 min. It was cooled to room temperature and poured into ice. It was slowly neutralized with ammonium hydroxide solution until a pH of 7 and the formed yellow solid was filtered, washed with water and dried to obtain compound 7 as a mixture of 2-chloro-3-nitropyridin-4-amine and 2-chloro-5-nitropyridin-4-amine intermediates. Sodium hydride (275 mg, 6.90 mmol) was carefully added to 20 mL of THF under N_2 and compound 7 (1.0 g, 5.76 mmol) was slowly added to the solution at 0 °C. The reaction mixture was stirred for 1 h, followed by the addition of benzyl bromide (0.75 mL, 6.34 mmol). The reaction mixture was stirred at room temperature for 2 h and poured into ice water. Then it was extracted with EtOAc (3×20 mL), washed with water, dried over sodium sulfate. The solvent was removed and the crude residue was purified using column chromatography (20% EtOAc/hexane) to obtain the compound 8 as a mixture of regioisomeric *N*-benzyl-2-chloro-3-nitropyridin-4-amine and *N*-benzyl-2-chloro-5-nitropyridin-4-amine intermediates. To this regioisomeric mixture (1.0 g, 4.2 mmol) in 20 mL of MeOH were added zinc dust (1.4 g, 21.0 mmol) and ammonium formate (1.4 g, 21.0 mmol). The reaction mixture was stirred at room temperature for 10 min and filtered through celite. Then the solvent was evaporated and the residue was dissolved in water. This was extracted with EtOAc (3×20 mL), washed with water and dried over sodium sulfate. The filtrate evaporated under vacuum, and chromatographed (20% EtOAc/hexane) to obtain the required *N*⁴-benzyl-2-chloropyridine-3,4-diamine compound 9a. Also obtained was *N*⁴-benzyl-6-chloropyridine-3,4-diamine as a side-product. To a solution of compound 9a (495 mg, 2.12 mmol) in 20 mL of anhydrous THF were added triethylamine (0.31 mL, 2.23 mmol) and valeryl chloride (0.28 mL, 2.33 mmol). The reaction mixture was refluxed for 1 h. The solvent was then removed under vacuum, and the residue was dissolved in 20 mL of EtOH and NaOH (170 mg, 4.24 mmol) in 2 mL of H_2O was added. The reaction mixture was refluxed for 2 h. The solvent was then removed under vacuum, and the residue was dissolved in EtOAc and washed with water. The EtOAc fraction was dried using sodium sulfate and evaporated and purified using column chromatography (5% MeOH/ CH_2Cl_2) to obtain the compound 10a (203 mg, 32%). ^1H NMR (500 MHz, CDCl_3) δ 8.10 (d, $J = 5.6$ Hz, 1H), 7.35 – 7.30 (m, 3H), 7.08 (d, $J = 5.6$ Hz,

1H), 7.01 (dd, $J = 7.4, 2.0$ Hz, 2H), 5.34 (s, 2H), 2.91 – 2.86 (m, 2H), 1.78 (dt, $J = 15.7, 7.7$ Hz, 2H), 1.45 – 1.36 (m, 2H), 0.90 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 158.03, 141.88, 141.75, 141.12, 137.03, 134.84, 129.39, 128.58, 126.25, 105.18, 47.90, 30.04, 27.63, 22.73, 13.82. MS (ESI) calculated for $\text{C}_{17}\text{H}_{19}\text{ClN}_3$, m/z 300.1262, found 300.1159 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 11a: 1-Benzyl-*N*,2-dibutyl-1H-imidazo[4,5-c]pyridin-4-amine. To a solution of compound 10 (50 mg, 0.17 mmol) in 1 mL of dioxane were added potassium *tert*-butoxide (57 mg, 0.51 mmol), catalytic amount of 2-dicyclohexylphosphino-2'-(*N,N*-dimethylamino)biphenyl (DavePhos) and *tris*(dibenzylideneacetone)dipalladium(0) ($\text{Pd}_2(\text{dba})_3$) and butyl amine (83 μL , 0.83 mmol). The reaction mixture was then heated under microwave conditions (500 W, 100 °C) in a sealed vial for 1 h. It was cooled to room temperature and filtered through celite and washed with MeOH. The solvent was removed and the crude residue was purified using column chromatography (5% MeOH/ CH_2Cl_2) to obtain the compound 11a (21 mg, 36%). ^1H NMR (500 MHz, CDCl_3) δ 7.84 (d, $J = 5.9$ Hz, 1H), 7.33 – 7.27 (m, 3H), 7.03 (dd, $J = 4.5, 3.6$ Hz, 2H), 6.48 (d, $J = 5.9$ Hz, 1H), 5.41 (s, 1H), 5.25 (s, 2H), 3.60 (dt, $J = 12.9, 6.5$ Hz, 2H), 2.80 – 2.74 (m, 2H), 1.73 – 1.67 (m, 4H), 1.49 (dt, $J = 15.0, 7.4$ Hz, 3H), 1.38 (dd, $J = 15.0, 7.5$ Hz, 2H), 0.96 (t, $J = 7.4$ Hz, 3H), 0.89 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 153.15, 151.39, 135.95, 129.24, 129.15, 128.15, 127.25, 126.34, 126.18, 96.16, 47.43, 41.09, 32.22, 30.27, 27.41, 22.69, 20.47, 14.11, 13.89. HRMS (ESI) calculated for $\text{C}_{21}\text{H}_{29}\text{N}_4$, m/z 337.2387, found 337.2451 ($\text{M} + \text{H}$) $^+$.

Compounds 11b was synthesized similarly as compound 11a.

***N*,1-Dibenzyl-2-butyl-1H-imidazo[4,5-c]pyridin-4-amine (11b).** Benzyl amine was used as a reagent. (33 mg, 52%). ^1H NMR (500 MHz, CDCl_3) δ 7.86 (d, $J = 5.9$ Hz, 1H), 7.45 (dd, $J = 7.9, 0.9$ Hz, 2H), 7.35 – 7.27 (m, 6H), 7.03 (d, $J = 6.4$ Hz, 2H), 6.54 (d, $J = 5.9$ Hz, 1H), 5.76 (s, 1H), 5.26 (s, 2H), 4.83 (d, $J = 5.6$ Hz, 2H), 2.79 – 2.74 (m, 2H), 1.72 – 1.67 (m, 2H), 1.37 (dq, $J = 14.8, 7.4$ Hz, 2H), 0.88 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 153.39, 150.92, 140.51, 139.81, 139.78, 135.88, 129.18, 128.62, 128.23, 128.19, 127.24, 126.36, 126.24, 96.73, 47.46, 45.40, 30.14, 27.38, 22.67, 13.88. HRMS (ESI) calculated for $\text{C}_{24}\text{H}_{27}\text{N}_4$, m/z 371.2230, found 371.2303 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 11c: 1-Benzyl-4-butoxy-2-butyl-1H-imidazo[4,5-c]pyridine. To a suspension of sodium hydride (48 mg, 2.00 mmol) in 2 mL of anhydrous THF was added 1-butanol (0.18 mL, 2.00 mmol). It was stirred at room temperature for 1 h, followed by the addition of compound 10 (100 mg, 0.33 mmol). The reaction mixture was heated at 60 °C for 18 h and then solvent was evaporated under vacuum. The residue was extracted with EtOAc (3×10 mL), washed with water and dried over sodium sulfate. The solvent was removed and the crude residue was purified using column chromatography (5% MeOH/ CH_2Cl_2) to obtain the compound 11c (61 mg, 55%). ^1H NMR (500 MHz, CDCl_3) δ 7.85 (d, $J = 5.8$ Hz, 1H), 7.33 – 7.28 (m, 3H), 7.01 (dd, $J = 7.7, 1.7$ Hz, 2H), 6.77 (d, $J = 5.8$ Hz, 1H), 5.30 (s, 2H), 4.52 (t, $J = 7.0$ Hz, 2H), 2.87 – 2.82 (m, 2H), 1.90 (dd, $J = 15.0, 7.1$

Hz, 2H), 1.75 (dt, $J = 15.7, 7.7$ Hz, 2H), 1.58 – 1.48 (m, 2H), 1.38 (dq, $J = 14.7, 7.4$ Hz, 2H), 0.97 (t, $J = 7.4$ Hz, 3H), 0.88 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 156.04, 155.18, 142.02, 138.95, 135.59, 129.21, 128.27, 127.41, 126.27, 100.47, 66.25, 47.61, 31.34, 30.21, 27.44, 22.75, 19.45, 14.09, 13.85. HRMS (ESI) calculated for $\text{C}_{21}\text{H}_{28}\text{N}_3\text{O}$, m/z 338.2227, found 338.2307 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 12: 2,6-Dichloro-3-nitropyridin-4-amine. 4-Amino-2,6-dichloropyridine (2.0 g, 12.27 mmol) was added to 20 mL of conc. H_2SO_4 . The mixture was cooled to 0 °C and 10 mL of conc. HNO_3 was dropwise at 0 °C. The reaction mixture was stirred at room temperature for 1 h and then poured into crushed ice. The white solid was filtered, washed with water and dried. This intermediate was dissolved in 10 mL of conc. H_2SO_4 and the reaction solution was heated at 90 °C for 30 min. It was cooled to room temperature and poured into ice. It was slowly neutralized with ammonium hydroxide solution until a pH of 9 and the formed yellow solid was filtered, washed with water and dried to obtain compound **12** as light yellow solid. ^1H NMR (500 MHz, MeOD) δ 6.84 (s, 1H). ^{13}C NMR (126 MHz, MeOD) δ 152.53, 151.18, 144.57, 111.12.

Synthesis of compound 13: *N*-Benzyl-2,6-dichloro-3-nitropyridin-4-amine. Sodium hydride (138 mg, 5.77 mmol) was carefully suspended in 10 mL of dry THF under N_2 . The grey suspension was cooled to 0 °C and compound **12** (1.0 g, 4.81 mmol) was slowly added to the suspension at 0 °C. The reaction mixture was stirred for 1 h, followed by the addition of benzyl bromide (0.5 mL, 5.29 mmol). The reaction mixture was stirred at room temperature for 2 h and poured into ice water. Then it was extracted with EtOAc (3 \times 20 mL), washed with water, dried over sodium sulfate. The solvent was removed and the crude residue was purified using column chromatography (20% EtOAc/hexane) to obtain the compound **13** as yellow solid. ^1H NMR (500 MHz, CDCl_3) δ 7.44 – 7.34 (m, 3H), 7.32 – 7.27 (m, 2H), 6.99 (s, 1H), 6.67 (s, 1H), 4.47 (d, $J = 5.4$ Hz, 2H). ^{13}C NMR (126 MHz, CDCl_3) δ 152.07, 149.97, 145.68, 135.01, 129.48, 128.71, 127.32, 106.62, 47.74.

Synthesis of compound 14: *N*⁴-Benzyl-6-chloro-3-nitro-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine. To a solution of compound **13** (300 mg, 1.02 mmol) in 20 mL of CH_2Cl_2 were added triethylamine (0.21 mL, 1.53 mmol) and *tert*-octylamine (0.5 mL, 3.06 mmol). The reaction mixture was refluxed for 48 h and the solvent was removed under vacuum. The crude residue was purified using column chromatography (20% EtOAc/hexane) to obtain the compound **14** as yellow solid (360 mg, 91%). ^1H NMR (500 MHz, CDCl_3) δ 9.57 (s, 1H), 9.51 (s, 1H), 7.39 (ddd, $J = 7.5, 4.4, 1.3$ Hz, 2H), 7.36 – 7.30 (m, 3H), 5.92 (s, 1H), 4.44 (d, $J = 5.4$ Hz, 2H), 1.95 (s, 2H), 1.56 (s, 6H), 0.98 (s, 9H). ^{13}C NMR (126 MHz, CDCl_3) δ 155.05, 154.25, 153.44, 136.10, 129.24, 128.24, 127.43, 115.80, 94.21, 57.32, 51.54, 47.64, 31.94, 31.66, 29.84. MS (ESI) calculated for $\text{C}_{20}\text{H}_{28}\text{ClN}_4\text{O}_2$, m/z 391.1895, found 391.1901 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 16: 1-Benzyl-2-butyl-6-chloro-*N*-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-

amine. To a solution of compound **14** (260 mg, 0.66 mmol) in 20 mL of MeOH were added zinc dust (434 mg, 6.60 mmol) and ammonium formate (416 mg, 6.60 mmol). The reaction mixture was stirred at room temperature for 10 min and filtered through celite. Then the solvent was evaporated and the residue was dissolved in water. This was extracted with EtOAc (3 \times 20 mL), washed with water and dried over sodium sulfate. The solvent was removed under vacuum to obtain compound **15**, brown oil (184 mg, 77%). To a solution of compound **15** (184 mg, 0.50 mmol) in 10 mL of anhydrous THF were added triethylamine (74 μL , 0.52 mmol) and valeryl chloride (62 μL , 0.50 mmol). The reaction mixture was refluxed for 1 h. The solvent was then removed under vacuum, and the residue was dissolved in 10 mL of EtOH and NaOH (40 mg, 1.00 mmol) in 1 mL of H_2O was added. The reaction mixture was refluxed for 5 h. The solvent was then removed under vacuum, and the residue was dissolved in EtOAc and washed with water. The EtOAc fraction was dried using sodium sulfate and evaporated and purified using column chromatography (20% EtOAc/hexane) to obtain the compound **16** as yellow solid (120 mg, 56%). ^1H NMR (500 MHz, CDCl_3) δ 7.34 – 7.28 (m, 3H), 7.02 (d, $J = 6.5$ Hz, 2H), 6.42 (s, 1H), 5.38 (s, 1H), 5.16 (s, 2H), 2.76 – 2.69 (m, 2H), 2.02 (s, 2H), 1.71 – 1.65 (m, 4H), 1.60 (s, 6H), 1.37 (dd, $J = 15.0, 7.5$ Hz, 2H), 1.01 (s, 9H), 0.89 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 153.37, 149.45, 141.67, 140.77, 135.64, 129.21, 128.22, 126.31, 125.50, 93.95, 55.74, 51.42, 47.42, 31.91, 31.73, 30.12, 29.96, 27.37, 22.66, 13.87. MS (ESI) calculated for $\text{C}_{25}\text{H}_{36}\text{ClN}_4$, m/z 427.2623, found 427.2635 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 17: 1-Benzyl-2-butyl-6-chloro-1*H*-imidazo[4,5-*c*]pyridin-4-amine. Compound **16** (34 mg, 0.082 mmol) was dissolved in 1 mL of HCl (4M in dioxane) and stirred at room temperature for 30 min. Then the solvent was removed under vacuum to obtain compound **17** (11 mg, 42%). ^1H NMR (500 MHz, CDCl_3) δ 7.36 – 7.30 (m, 3H), 7.01 (d, $J = 6.4$ Hz, 2H), 6.58 (s, 1H), 5.23 (s, 2H), 5.21 (s, 2H), 2.80 – 2.73 (m, 2H), 1.72 (dt, $J = 15.5, 7.7$ Hz, 2H), 1.39 (dq, $J = 14.8, 7.4$ Hz, 2H), 0.90 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 154.83, 149.94, 142.31, 141.73, 135.26, 129.32, 128.57, 128.41, 126.26, 96.29, 47.59, 29.92, 27.36, 22.63, 13.87. HRMS (ESI) calculated for $\text{C}_{17}\text{H}_{20}\text{ClN}_4$, m/z 315.1371, found 315.1422 ($\text{M} + \text{H}$) $^+$.

Synthesis of compound 19a: 1-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine. To a solution of compound **16** (70 mg, 0.16 mmol) in 1 mL of dioxane were added potassium *tert*-butoxide (92 mg, 0.82 mmol), catalytic amount of DavePhos and *tert*-octylamine (83 μL , 0.83 mmol). The reaction mixture was then heated under microwave conditions (500 W, 100 °C) in a sealed vial for 1 h. It was cooled to room temperature and filtered through celite and washed with MeOH. The solvent was removed and the crude residue was purified using column chromatography (20% EtOAc/hexane) to obtain the compound **18a**, brown solid (41 mg, 49%). Compound **18a** (33 mg, 0.063 mmol) was dissolved in 1 mL of HCl (4M in dioxane) and stirred at room temperature for 30 min. Then the solvent was removed under vacuum to obtain compound **19a** as brown solid (11 mg, 58%). ^1H NMR (500 MHz, DMSO) δ 7.37 – 7.31 (m, 2H), 7.30 – 7.25 (m, 1H), 7.09 – 7.04 (m, 2H), 6.41 (s, 2H), 5.65 (s, 1H), 5.24

(s, 2H), 2.73 – 2.67 (m, 2H), 1.59 (dt, $J = 15.3, 7.5$ Hz, 2H), 1.31 (dq, $J = 14.7, 7.4$ Hz, 2H), 0.83 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, DMSO) δ 151.81, 147.80, 144.08, 136.91, 128.81, 128.77, 127.47, 126.29, 117.96, 76.19, 46.11, 29.27, 26.17, 21.80, 13.68. HRMS (ESI) calculated for $\text{C}_{17}\text{H}_{22}\text{N}_5$, m/z 296.1870, found 296.1906 (M + H) $^+$.

Compounds **19b-19o** were synthesized similarly as compound **19a**.

1-Benzyl-2-butyl- N^6 -dibutyl-1H-imidazo[4,5-c]pyridine-4,6-diamine (19b). Butyl amine was used as a reagent. Brown solid (23 mg, 79%). ^1H NMR (500 MHz, CDCl_3) δ 7.31 (ddd, $J = 8.6, 6.4, 3.4$ Hz, 3H), 7.05 (d, $J = 6.7$ Hz, 2H), 5.46 (s, 1H), 5.26 (s, 1H), 5.16 (s, 2H), 3.03 (t, $J = 7.1$ Hz, 2H), 2.72 – 2.66 (m, 2H), 1.68 (dd, $J = 15.5, 7.6$ Hz, 2H), 1.55 (dd, $J = 14.8, 7.2$ Hz, 2H), 1.43 – 1.33 (m, 4H), 0.90 (dt, $J = 16.4, 7.4$ Hz, 6H). ^{13}C NMR (126 MHz, CDCl_3) δ 152.42, 148.50, 144.42, 135.99, 129.14, 128.06, 126.33, 74.80, 47.20, 43.09, 31.41, 30.03, 27.37, 22.66, 20.40, 14.02, 13.89. HRMS (ESI) calculated for $\text{C}_{21}\text{H}_{30}\text{N}_5$, m/z 352.2496, found 352.2515 (M + H) $^+$.

1-Benzyl-2-butyl- N^6 -heptyl- N^4 -(2,4,4-trimethylpentan-2-yl)-1H-imidazo[4,5-c]pyridine-4,6-diamine (18c). Heptylamine was used as a reagent. Brown oil (43 mg, 65%). ^1H NMR (500 MHz, CDCl_3) δ 7.32 – 7.26 (m, 3H), 7.05 (d, $J = 6.7$ Hz, 2H), 5.39 (s, 1H), 5.13 (s, 1H), 5.11 (s, 2H), 3.13 (t, $J = 7.2$ Hz, 2H), 2.68 – 2.64 (m, 2H), 2.04 (s, 2H), 1.65 – 1.55 (m, 11H), 1.41 – 1.22 (m, 12H), 1.01 (s, 9H), 0.89 – 0.84 (m, 6H). ^{13}C NMR (126 MHz, CDCl_3) δ 154.28, 150.38, 149.22, 142.39, 136.63, 128.97, 128.05, 127.76, 126.40, 120.44, 95.52, 73.57, 55.27, 51.72, 47.00, 43.63, 31.97, 31.94, 31.90, 31.79, 30.47, 30.29, 29.97, 29.32, 27.44, 27.39, 22.77, 22.71, 14.25, 13.90. MS (ESI) calculated for $\text{C}_{32}\text{H}_{52}\text{N}_5$, m/z 506.4217, found 506.4209 (M + H) $^+$.

1-Benzyl-2-butyl- N^6 -heptyl-1H-imidazo[4,5-c]pyridine-4,6-diamine (19c). Light brown solid (24 mg, 80%). ^1H NMR (500 MHz, DMSO) δ 8.03 (s, 2H), 7.38 – 7.33 (m, 2H), 7.32 – 7.28 (m, 1H), 7.12 (d, $J = 7.2$ Hz, 2H), 6.84 (s, 1H), 5.98 (s, 1H), 5.39 (s, 2H), 3.08 (t, $J = 7.0$ Hz, 2H), 2.73 – 2.68 (m, 2H), 1.59 (dt, $J = 15.3, 8.2$ Hz, 2H), 1.50 (dd, $J = 14.3, 7.2$ Hz, 2H), 1.33 – 1.21 (m, 12H), 0.85 (t, $J = 7.0$ Hz, 3H), 0.81 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, DMSO) δ 154.62, 145.66, 145.15, 136.32, 128.96, 128.86, 127.73, 126.64, 126.55, 74.21, 46.34, 42.24, 31.25, 28.78, 28.42, 28.09, 26.34, 26.18, 22.07, 21.71, 13.98, 13.62. HRMS (ESI) calculated for $\text{C}_{24}\text{H}_{36}\text{N}_5$, m/z 394.2965, found 394.3001 (M + H) $^+$.

1-Benzyl-2-butyl- N^6 -(cyclohexylmethyl)- N^4 -(2,4,4-trimethylpentan-2-yl)-1H-imidazo[4,5-c]pyridine-4,6-diamine (18d). *N*-Cyclohexylmethylamine was used as a reagent. Yellow solid (50 mg, 62%). ^1H NMR (500 MHz, CDCl_3) δ 7.32 – 7.27 (m, 3H), 7.06 (d, $J = 6.7$ Hz, 2H), 5.37 (s, 1H), 5.10 (s, 2H), 2.99 (d, $J = 6.7$ Hz, 2H), 2.69 – 2.64 (m, 2H), 2.04 (s, 2H), 1.79 (d, $J = 13.2$ Hz, 2H), 1.73 – 1.60 (m, 9H), 1.58 (s, 6H), 1.35 (dd, $J = 15.0, 7.5$ Hz, 2H), 1.27 – 1.13 (m, 4H), 1.01 (s, 9H), 0.98 – 0.91 (m, 2H), 0.86 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ

154.37, 150.36, 149.19, 142.41, 136.65, 128.97, 127.76, 126.44, 120.35, 73.62, 55.26, 51.67, 50.18, 47.02, 37.88, 31.91, 31.80, 31.48, 30.51, 30.31, 27.46, 26.80, 26.17, 22.72, 13.91. MS (ESI) calculated for $\text{C}_{32}\text{H}_{50}\text{N}_5$, m/z 504.4061, found 504.41 (M + H) $^+$.

1-Benzyl-2-butyl- N^6 -(cyclohexylmethyl)-1H-imidazo[4,5-c]pyridine-4,6-diamine (19d). Light brown (28 mg, 80%). ^1H NMR (500 MHz, DMSO) δ 7.35 (t, $J = 7.3$ Hz, 2H), 7.31 – 7.27 (m, 1H), 7.10 (d, $J = 7.2$ Hz, 2H), 6.34 (d, $J = 3.1$ Hz, 1H), 5.85 (s, 1H), 5.33 (s, 2H), 2.92 (t, $J = 6.1$ Hz, 2H), 2.71 – 2.66 (m, 2H), 1.75 – 1.64 (m, 4H), 1.57 (dt, $J = 15.3, 7.6$ Hz, 3H), 1.49 (dtd, $J = 11.2, 7.4, 3.7$ Hz, 1H), 1.30 (dt, $J = 14.8, 7.4$ Hz, 2H), 1.21 – 1.09 (m, 4H), 0.91 (dd, $J = 22.3, 10.5$ Hz, 2H), 0.81 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, DMSO) δ 153.32, 146.28, 145.05, 136.69, 128.79, 127.61, 126.53, 116.94, 74.21, 48.49, 46.22, 36.76, 30.47, 29.02, 26.21, 26.05, 25.43, 21.75, 13.65. HRMS (ESI) calculated for $\text{C}_{24}\text{H}_{34}\text{N}_5$, m/z 392.2809, found 392.2806 (M + H) $^+$.

1-Benzyl-2-butyl- N^6 -phenyl- N^4 -(2,4,4-trimethylpentan-2-yl)-1H-imidazo[4,5-c]pyridine-4,6-diamine (18e). Aniline was used as a reagent. Yellow solid (51 mg, 55%). ^1H NMR (500 MHz, CDCl_3) δ 7.31 (dt, $J = 16.5, 5.5$ Hz, 3H), 7.22 (t, $J = 7.8$ Hz, 2H), 7.16 (d, $J = 7.6$ Hz, 2H), 7.05 (d, $J = 6.7$ Hz, 2H), 6.90 (t, $J = 6.9$ Hz, 1H), 6.16 (s, 1H), 6.02 (s, 1H), 5.23 (s, 1H), 5.12 (s, 2H), 2.74 – 2.67 (m, 2H), 2.03 (s, 2H), 1.66 (dd, $J = 15.8, 8.2$ Hz, 4H), 1.60 (s, 6H), 1.38 (dd, $J = 15.0, 7.5$ Hz, 2H), 1.02 (s, 9H), 0.89 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 151.49, 149.42, 149.11, 142.46, 141.51, 136.32, 129.16, 129.06, 127.93, 126.53, 121.98, 120.91, 118.83, 55.42, 51.80, 47.26, 31.93, 31.80, 30.44, 30.19, 27.46, 22.71, 13.91. MS (ESI) calculated for $\text{C}_{31}\text{H}_{42}\text{N}_5$, m/z 484.3435, found 484.3459 (M + H) $^+$.

1-Benzyl-2-butyl- N^6 -phenyl-1H-imidazo[4,5-c]pyridine-4,6-diamine (19e). Light yellow solid (28 mg, 76%). ^1H NMR (500 MHz, DMSO) δ 9.10 (s, 1H), 8.08 (s, 1H), 7.39 – 7.35 (m, 2H), 7.33 – 7.28 (m, 3H), 7.11 (t, $J = 7.3$ Hz, 4H), 7.01 (t, $J = 7.3$ Hz, 1H), 6.41 (s, 1H), 5.44 (s, 2H), 2.82 (t, $J = 7.5$ Hz, 2H), 1.63 (dt, $J = 15.3, 7.6$ Hz, 2H), 1.33 (dd, $J = 14.9, 7.4$ Hz, 2H), 0.84 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, DMSO) δ 146.34, 140.41, 136.06, 129.52, 128.92, 127.84, 126.69, 122.44, 118.87, 46.68, 28.89, 26.15, 21.72, 13.63. HRMS (ESI) calculated for $\text{C}_{23}\text{H}_{26}\text{N}_5$, m/z 372.2183, found 372.2219 (M + H) $^+$.

N^6 -Dibenzyl-2-butyl- N^4 -(2,4,4-trimethylpentan-2-yl)-1H-imidazo[4,5-c]pyridine-4,6-diamine (18f). Benzyl amine was used as a reagent. Light brown solid (30 mg, 46%). ^1H NMR (500 MHz, CDCl_3) δ 7.35 (d, $J = 7.4$ Hz, 2H), 7.26 (dddd, $J = 9.4, 7.2, 6.1, 1.6$ Hz, 6H), 7.03 (d, $J = 6.8$ Hz, 2H), 5.44 (s, 1H), 5.17 (s, 1H), 5.07 (s, 2H), 4.42 (s, 2H), 2.70 – 2.64 (m, 2H), 1.67 – 1.60 (m, 2H), 1.55 (s, 6H), 1.35 (dq, $J = 14.7, 7.4$ Hz, 2H), 1.02 – 0.94 (m, 9H), 0.87 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3) δ 153.83, 150.58, 149.20, 142.27, 140.52, 136.49, 128.98, 128.56, 127.80, 127.68, 126.97, 126.46, 126.39, 74.17, 55.29, 51.62, 47.59, 47.07, 31.87, 31.76, 30.45, 30.28, 30.20, 27.42, 22.70, 13.90. MS (ESI) calculated for $\text{C}_{32}\text{H}_{44}\text{N}_5$, m/z 498.3591, found 498.3599 (M + H) $^+$.

***N*⁶-1-Dibenzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19f).** White solid (22 mg, 85%). ¹H NMR (500 MHz, DMSO) δ 8.07 (s, 2H), 7.38 – 7.35 (m, 2H), 7.34 – 7.29 (m, 5H), 7.29 – 7.25 (m, 1H), 7.07 (d, *J* = 6.3 Hz, 2H), 6.06 (s, 1H), 5.33 (s, 2H), 4.38 (s, 2H), 2.75 – 2.69 (m, 2H), 1.58 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.30 (dt, *J* = 14.8, 7.4 Hz, 2H), 0.81 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 154.68, 145.29, 137.99, 136.12, 128.85, 128.45, 127.78, 127.51, 127.23, 126.74, 75.08, 46.43, 45.67, 28.76, 26.16, 21.70, 13.62. HRMS (ESI) calculated for C₂₄H₂₈N₅, m/z 386.2339, found 386.2389 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(4-methoxybenzyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19g). 4-Methoxybenzyl amine was used as a reagent. Light yellow solid (22 mg, 69%). ¹H NMR (500 MHz, DMSO) δ 8.07 (s, 2H), 7.34 – 7.28 (m, 5H), 7.08 (d, *J* = 6.4 Hz, 2H), 6.87 (d, *J* = 8.7 Hz, 2H), 6.04 (s, 1H), 5.34 (s, 2H), 4.29 (s, 2H), 3.72 (s, 3H), 2.75 – 2.70 (m, 2H), 1.58 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.29 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.81 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 158.48, 154.67, 145.28, 145.27, 136.13, 129.73, 128.92, 128.84, 127.78, 126.74, 113.82, 75.11, 55.08, 46.43, 45.18, 28.76, 26.16, 21.71, 13.62. HRMS (ESI) calculated for C₂₅H₃₀N₅O, m/z 416.2445, found 416.2490 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(3-methoxybenzyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19h). 3-Methoxybenzyl amine was used as a reagent. Brown solid (27 mg, 69%). ¹H NMR (500 MHz, DMSO) δ 7.32 – 7.27 (m, 3H), 7.23 (t, *J* = 7.9 Hz, 1H), 7.06 (d, *J* = 6.1 Hz, 2H), 6.94 (dd, *J* = 12.3, 4.8 Hz, 2H), 6.83 (dd, *J* = 8.1, 2.2 Hz, 1H), 6.03 (s, 1H), 5.31 (s, 2H), 4.33 (d, *J* = 5.8 Hz, 2H), 3.71 (s, 3H), 2.70 (t, *J* = 7.6 Hz, 2H), 1.57 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.29 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.81 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 159.37, 136.32, 129.52, 128.80, 127.72, 126.70, 119.55, 113.13, 112.50, 75.05, 55.00, 46.33, 45.63, 40.11, 40.02, 39.94, 28.89, 26.21, 21.72, 13.63. HRMS (ESI) calculated for C₂₅H₃₀N₅O, m/z 416.2445, found 416.2506 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(4-(trifluoromethyl)benzyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19i). 4-(Trifluoromethyl)benzyl amine was used as a reagent. Brown solid (17 mg, 77%). ¹H NMR (500 MHz, DMSO) δ 7.60 (d, *J* = 8.2 Hz, 2H), 7.51 (d, *J* = 8.1 Hz, 2H), 7.29 – 7.21 (m, 3H), 7.02 (d, *J* = 6.7 Hz, 2H), 6.23 (s, 1H), 5.70 (s, 2H), 5.57 (s, 1H), 5.15 (s, 2H), 4.41 (d, *J* = 6.3 Hz, 2H), 2.69 – 2.63 (m, 2H), 1.56 (dt, *J* = 15.3, 7.5 Hz, 2H), 1.29 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.82 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 153.45, 150.32, 149.41, 146.48, 143.12, 137.24, 128.58, 127.82, 127.31, 126.51, 124.88, 124.85, 118.91, 75.32, 46.04, 44.94, 29.43, 26.19, 21.82, 13.69. HRMS (ESI) calculated for C₂₅H₂₇F₃N₅, m/z 454.2213, found 454.2284 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(4-chlorobenzyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19j). 4-Chlorobenzylamine was used as a reagent. Yellow solid (15 mg, 63%). ¹H NMR (500 MHz, DMSO) δ 7.33 – 7.25 (m, 8H), 7.02 (d, *J* = 6.6 Hz, 2H), 6.18 (s, 1H), 5.83 (s, 2H), 5.59 (s, 1H), 5.16 (s, 2H), 4.30 (d, *J* = 6.0 Hz, 2H), 2.68 – 2.64 (m, 2H), 1.56 (dt, *J* = 15.3, 7.5 Hz, 2H), 1.29

(dd, *J* = 14.9, 7.4 Hz, 2H), 0.82 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 150.57, 149.14, 143.25, 140.18, 137.18, 130.77, 129.08, 128.63, 127.96, 127.37, 126.56, 118.72, 75.34, 46.06, 44.71, 40.11, 40.02, 39.94, 29.40, 26.20, 21.82, 13.69. HRMS (ESI) calculated for C₂₄H₂₇ClN₅, m/z 420.1950, found 420.1976 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(furan-2-ylmethyl)-*N*⁴-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (18k). Furfuryl amine was used as a reagent. Yellow oil (53 mg, 54%). ¹H NMR (500 MHz, CDCl₃) δ 7.30 (ddd, *J* = 10.9, 4.6, 1.1 Hz, 4H), 7.04 (d, *J* = 6.6 Hz, 2H), 6.28 (dd, *J* = 3.2, 1.9 Hz, 1H), 6.15 (dd, *J* = 3.2, 0.6 Hz, 1H), 5.50 (s, 1H), 5.10 (s, 2H), 4.43 (s, 2H), 2.72 – 2.64 (m, 2H), 2.03 (s, 2H), 1.68 – 1.59 (m, 4H), 1.58 (s, 6H), 1.35 (dt, *J* = 14.7, 7.4 Hz, 2H), 1.00 (s, 9H), 0.87 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 153.92, 153.28, 150.77, 149.15, 142.19, 141.69, 136.47, 129.00, 127.83, 126.42, 110.36, 106.53, 74.85, 55.32, 51.68, 47.09, 40.78, 31.89, 31.77, 30.47, 30.31, 27.43, 22.71, 13.90. MS (ESI) calculated for C₃₀H₄₂N₅O, m/z 488.3384, found 488.3446 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(furan-2-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19k). Brown solid (21 mg, 55%). ¹H NMR (500 MHz, DMSO) δ 7.55 (s, 1H), 7.34 (t, *J* = 7.3 Hz, 2H), 7.31 – 7.27 (m, 1H), 7.10 (d, *J* = 7.3 Hz, 2H), 6.36 (dd, *J* = 3.0, 1.9 Hz, 1H), 6.29 (d, *J* = 2.6 Hz, 1H), 5.99 (s, 1H), 5.31 (s, 2H), 4.35 (d, *J* = 5.8 Hz, 2H), 2.73 – 2.67 (m, 2H), 1.58 (dt, *J* = 15.2, 7.6 Hz, 2H), 1.30 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.82 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 142.17, 136.66, 128.79, 127.62, 126.61, 110.39, 107.43, 75.34, 60.19, 46.26, 38.71, 29.07, 26.21, 21.76, 13.65. HRMS (ESI) calculated for C₂₂H₂₆N₅O, m/z 376.2132, found 376.2184 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(pyridin-4-ylmethyl)-*N*⁴-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (18l). 4-Picolylamine was used as a reagent. Yellow solid (25 mg, 31%). ¹H NMR (500 MHz, CDCl₃) δ 8.48 (dd, *J* = 4.5, 1.6 Hz, 2H), 7.29 – 7.25 (m, 3H), 7.24 (dd, *J* = 4.5, 1.5 Hz, 2H), 7.01 (d, *J* = 6.1 Hz, 2H), 5.40 (s, 1H), 5.06 (s, 2H), 4.49 (s, 2H), 2.70 – 2.64 (m, 2H), 1.92 (s, 2H), 1.63 (dt, *J* = 15.5, 7.7 Hz, 2H), 1.48 (s, 6H), 1.36 (dt, *J* = 14.9, 7.4 Hz, 2H), 0.95 (s, 9H), 0.87 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 149.83, 136.31, 129.01, 127.89, 126.40, 122.20, 74.67, 55.25, 51.56, 47.13, 46.09, 31.81, 31.75, 31.71, 30.44, 30.23, 27.41, 22.69, 13.89. MS (ESI) calculated for C₃₁H₄₃N₆, m/z 499.3544, found 499.3475 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(pyridin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19l). Brown solid (7 mg, 47%). ¹H NMR (500 MHz, DMSO) δ 8.43 (dd, *J* = 4.5, 1.5 Hz, 2H), 7.27 (ddd, *J* = 9.7, 6.8, 4.4 Hz, 5H), 7.03 (d, *J* = 6.7 Hz, 2H), 6.50 – 6.28 (m, 2H), 5.65 (s, 1H), 5.18 (s, 2H), 4.37 (d, *J* = 6.3 Hz, 2H), 2.69 – 2.65 (m, 2H), 1.57 (dt, *J* = 15.3, 7.5 Hz, 2H), 1.30 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.81 (d, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 149.28, 137.09, 130.51, 128.65, 127.43, 126.53, 122.35, 113.97, 75.37, 46.09, 44.38, 29.34, 26.19, 21.80, 13.68. HRMS (ESI) calculated for C₂₃H₂₇N₆, m/z 387.2292, found 387.2299 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(pyridin-3-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19m). 3-Picolylamine was used as a reagent. Light brown solid (33 mg, 63%). ¹H NMR (500 MHz, DMSO) δ 8.92 (s, 1H), 8.80 (d, *J* = 5.3 Hz, 1H), 8.46 (d, *J* = 8.0 Hz, 1H), 8.12 (s, 2H), 7.94 (dd, *J* = 7.9, 5.7 Hz, 1H), 7.69 (s, 1H), 7.33 – 7.25 (m, 3H), 7.04 (d, *J* = 6.9 Hz, 2H), 6.03 (s, 1H), 5.32 (s, 2H), 4.64 (s, 2H), 2.72 (t, *J* = 7.6 Hz, 2H), 1.56 (dt, *J* = 15.2, 7.6 Hz, 2H), 1.28 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.80 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 154.65, 145.80, 145.01, 143.55, 141.29, 141.12, 138.17, 136.10, 128.83, 127.76, 126.68, 126.52, 75.49, 46.45, 42.51, 28.77, 26.12, 21.68, 13.60. HRMS (ESI) calculated for C₂₃H₂₇N₆, *m/z* 387.2292, found 387.2296 (M + H)⁺.

1-Benzyl-2-butyl-*N*⁶-(naphthalen-1-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19n). 1-Naphthylmethylamine was used as a reagent. Light brown solid (28 mg, 70%). ¹H NMR (500 MHz, DMSO) δ 12.12 (s, 1H), 8.10 (d, *J* = 8.0 Hz, 1H), 8.04 (s, 2H), 7.99 – 7.96 (m, 1H), 7.88 (d, *J* = 8.2 Hz, 1H), 7.61 – 7.53 (m, 3H), 7.45 (dd, *J* = 8.1, 7.1 Hz, 1H), 7.33 – 7.28 (m, 3H), 7.08 (d, *J* = 6.4 Hz, 2H), 6.22 (s, 1H), 5.35 (s, 2H), 4.83 (s, 2H), 2.74 – 2.68 (m, 2H), 1.58 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.29 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.81 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 154.69, 145.51, 145.31, 136.23, 133.44, 132.90, 130.92, 128.83, 128.63, 127.99, 127.76, 126.67, 126.39, 125.97, 125.49, 125.43, 123.63, 74.91, 46.39, 43.99, 28.80, 26.20, 21.71, 13.62. HRMS (ESI) calculated for C₂₈H₃₀N₅, *m/z* 436.2496, found 436.2549 (M + H)⁺.

***N*⁶-([1,1'-Biphenyl]-4-ylmethyl)-1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19o).** [1,1'-biphenyl]-4-yl methanamine was used as a reagent. Light brown solid (21 mg, 51%). ¹H NMR (500 MHz, DMSO) δ 12.36 (s, 1H), 8.04 (s, 2H), 7.63 (dd, *J* = 12.4, 7.8 Hz, 4H), 7.50 – 7.42 (m, 4H), 7.40 – 7.33 (m, 1H), 7.28 (t, *J* = 7.3 Hz, 2H), 7.23 (t, *J* = 7.2 Hz, 1H), 7.07 (d, *J* = 7.3 Hz, 2H), 6.07 (s, 1H), 5.33 (s, 2H), 4.42 (s, 2H), 2.71 (t, *J* = 7.6 Hz, 2H), 1.57 (dt, *J* = 15.2, 7.7 Hz, 2H), 1.29 (dq, *J* = 14.6, 7.3 Hz, 2H), 0.81 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 154.65, 145.40, 145.35, 139.80, 139.07, 137.24, 136.17, 128.95, 128.80, 128.06, 127.67, 127.43, 126.71, 126.69, 126.57, 75.11, 46.38, 45.31, 28.81, 26.18, 21.69, 13.61. HRMS (ESI) calculated for C₃₀H₃₂N₅, *m/z* 462.2652, found 462.2698 (M + H)⁺.

Synthesis of compound 19p and 19r: *N*⁶-(4-(aminomethyl)benzyl)-1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine. To a solution of compound 16 (70 mg, 0.16 mmol) in 1 mL of dioxane were added potassium *tert*-butoxide (90 mg, 0.80 mmol), catalytic amount of DavePhos and Pd₂(dba)₃ and *p*-xylylenediamine (109 mg, 0.80 mmol). The reaction mixture was then heated under microwave conditions (500 W, 100 °C) in a sealed vial for 1 h. It was cooled to room temperature and filtered through celite and washed with MeOH. The solvent was removed and the crude residue was purified using column chromatography (20% EtOAc/hexane) to obtain the compound 18p and 18r. Compound 18p (33 mg, 0.063 mmol) was dissolved in 1 mL of HCl (4M in dioxane) and stirred at room temperature for 30 min. Then the solvent was removed under vacuum to obtain compound

19p as light yellow solid (5 mg, 63%). ¹H NMR (500 MHz, DMSO) δ 8.37 (s, 2H), 8.06 (s, 2H), 7.42 (q, *J* = 8.5 Hz, 4H), 7.36 – 7.30 (m, 3H), 7.08 (d, *J* = 6.8 Hz, 2H), 6.08 (s, 1H), 5.34 (s, 2H), 4.40 (s, 2H), 3.98 (q, *J* = 5.8 Hz, 2H), 2.75 – 2.67 (m, 2H), 1.57 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.28 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.80 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 154.71, 145.41, 145.34, 138.43, 136.19, 132.98, 129.02, 128.89, 127.79, 127.75, 126.65, 46.40, 45.31, 41.89, 28.78, 26.18, 21.70, 13.62. HRMS (ESI) calculated for C₂₅H₃₁N₆, *m/z* 415.2605, found 415.2606 (M + H)⁺.

***N*⁶,*N*⁶-(1,4-phenylenebis(methylene))bis(1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19r).** Compound 18r (35 mg, 0.038 mmol) was dissolved in 1 mL of HCl (4M in dioxane) and stirred at room temperature for 30 min. Then the solvent was removed under vacuum to obtain compound 19r as light yellow solid (12 mg, 45%). ¹H NMR (500 MHz, MeOD) δ 7.27 (s, 4H), 7.21 – 7.17 (m, 6H), 7.02 – 6.98 (m, 4H), 5.20 (s, 4H), 4.33 (s, 4H), 2.78 – 2.73 (m, 4H), 1.67 (dt, *J* = 15.3, 7.6 Hz, 4H), 1.37 (dd, *J* = 15.0, 7.5 Hz, 4H), 0.89 (t, *J* = 7.4 Hz, 6H). ¹³C NMR (126 MHz, MeOD) δ 155.94, 150.96, 148.16, 146.47, 138.77, 137.12, 130.00, 129.03, 128.70, 127.70, 118.63, 48.01, 47.26, 30.28, 27.76, 23.29, 14.05. HRMS (ESI) calculated for C₄₂H₄₉N₁₀, *m/z* 693.4136, found 693.4331 (M + H)⁺.

Compounds 19q and 19s were synthesized similarly as compounds 19p and 19r.

***N*⁶-(3-(aminomethyl)benzyl)-1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19q).** *m*-xylylenediamine was used as a reagent. Light yellow solid. ¹H NMR (500 MHz, MeOD) δ 7.49 (s, 1H), 7.41 (dd, *J* = 3.9, 1.5 Hz, 2H), 7.39 – 7.35 (m, 1H), 7.32 – 7.28 (m, 3H), 7.05 – 7.02 (m, 2H), 5.31 (s, 2H), 4.44 (s, 2H), 4.07 (s, 2H), 2.79 – 2.75 (m, 2H), 1.70 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.38 (dq, *J* = 14.8, 7.4 Hz, 2H), 0.89 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 157.36, 147.23, 146.79, 140.00, 136.86, 134.96, 130.63, 130.12, 129.26, 129.19, 129.14, 129.00, 127.62, 48.10, 47.15, 44.18, 30.00, 27.75, 23.24, 14.03. HRMS (ESI) calculated for C₂₅H₃₁N₆, *m/z* 415.2605, found 415.2610 (M + H)⁺.

***N*⁶,*N*⁶-(1,3-phenylenebis(methylene))bis(1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridine-4,6-diamine (19s).** Light yellow solid. ¹H NMR (500 MHz, DMSO) δ 12.38 (s, 2H), 8.04 (s, 4H), 7.38 (s, 1H), 7.30 – 7.22 (m, 9H), 7.04 (d, *J* = 6.8 Hz, 4H), 6.01 (s, 2H), 5.28 (s, 4H), 4.34 (s, 4H), 2.73 – 2.68 (m, 4H), 1.56 (dd, *J* = 15.3, 7.7 Hz, 4H), 1.29 (dd, *J* = 14.9, 7.4 Hz, 4H), 0.81 (t, *J* = 7.4 Hz, 6H). ¹³C NMR (126 MHz, DMSO) δ 154.62, 145.36, 145.31, 138.22, 136.15, 128.80, 128.61, 127.72, 126.66, 126.56, 126.38, 46.36, 45.73, 28.80, 26.19, 21.71, 13.62. HRMS (ESI) calculated for C₄₂H₄₉N₁₀, *m/z* 693.4136, found 693.4131 (M + H)⁺.

Synthesis of compound 23a: To a solution of compound 14 (120 mg, 0.31 mmol) in 1 mL of dioxane were added cesium carbonate (303 mg, 0.93 mmol) in H₂O (0.5 mL), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (Pd(dppf)Cl₂) (15 mg, 0.019 mmol) and *n*-butylboronic acid (98 μL, 0.46 mmol) under N₂. The reaction mixture was then heated

at 90 °C in a sealed vial for 18 h. It was cooled to room temperature and filtered through celite and washed with MeOH. The solvent was removed and the crude residue was purified using column chromatography (15% EtOAc/hexane) to obtain the compound **20a** (97 mg, 76%). To a solution of compound **20a** (94 mg, 0.23 mmol) in 10 mL of MeOH were added zinc dust (149 mg, 2.30 mmol) and ammonium formate (145 mg, 2.30 mmol). The reaction mixture was stirred at room temperature for 10 min and filtered through celite. Then the solvent was evaporated and the residue was dissolved in water. This was extracted with EtOAc (3 × 20 mL), washed with water and dried over sodium sulfate. The solvent was removed under vacuum to obtain compound **21a** (45 mg, 51%). To a solution of compound **21a** (42 mg, 0.11 mmol) in 7 mL of anhydrous THF were added triethylamine (16 µL, 0.12 mmol) and valeryl chloride (13 µL, 0.11 mmol). The reaction mixture was refluxed for 1 h. The solvent was then removed under vacuum, and the residue was dissolved in 5 mL of EtOH and NaOH (10 mg, 0.22 mmol) in 1 mL of H₂O was added. The reaction mixture was refluxed for 18 h. The solvent was then removed under vacuum, and the residue was dissolved in EtOAc and washed with water. The EtOAc fraction was dried using sodium sulfate and evaporated and purified using column chromatography (20% EtOAc/hexane) to obtain the compound **22a** (25 mg, 51%). Compound **22a** (22 mg, 0.049 mmol) was dissolved in 1 mL of HCl (4M in dioxane) and stirred at room temperature for 30 min. Then the solvent was removed under vacuum to obtain compound **23a** (11 mg, 69%).

N⁴-Benzyl-6-butyl-3-nitro-N²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20a). Yellow oil (97 mg, 76%). ¹H NMR (500 MHz, CDCl₃) δ 9.51 (t, *J* = 4.4 Hz, 1H), 9.40 (s, 1H), 7.37 (ddd, *J* = 7.1, 4.4, 1.6 Hz, 2H), 7.34 – 7.30 (m, 3H), 5.74 (s, 1H), 4.45 (d, *J* = 5.4 Hz, 2H), 2.44 (t, *J* = 7.5 Hz, 2H), 2.01 (s, 2H), 1.67 – 1.61 (m, 2H), 1.30 (dd, *J* = 15.0, 7.4 Hz, 2H), 0.96 (s, 9H), 0.89 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 166.49, 154.69, 152.58, 136.94, 129.07, 127.92, 127.42, 115.70, 94.41, 56.59, 51.17, 47.44, 38.87, 31.95, 31.64, 30.61, 30.19, 22.43, 14.11. MS (ESI) calculated for C₂₄H₃₇N₄O₂, *m/z* 413.2911, found 413.3267 (M + H)⁺.

1-Benzyl-2,6-dibutyl-1H-imidazo[4,5-c]pyridin-4-amine (23a). White solid (11 mg, 69%). ¹H NMR (500 MHz, CDCl₃) δ 7.37 – 7.31 (m, 3H), 7.10 (d, *J* = 6.9 Hz, 2H), 6.81 (s, 1H), 5.47 (s, 2H), 2.85 (t, 2H), 2.72 (t, 2H), 1.74 (dd, *J* = 15.3, 7.6 Hz, 2H), 1.67 (ddd, *J* = 15.3, 10.4, 7.0 Hz, 2H), 1.38 (ddq, *J* = 14.8, 10.2, 7.4 Hz, 4H), 0.94 (t, *J* = 7.4 Hz, 3H), 0.90 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 157.75, 150.13, 143.74, 137.10, 130.16, 129.22, 127.59, 124.62, 97.63, 35.61, 32.84, 30.27, 27.85, 23.31, 23.22, 14.12, 14.04. HRMS (ESI) calculated for C₂₁H₂₉N₄, *m/z* 337.2387, found 337.2476 (M + H)⁺.

Compounds **23b–23j** were synthesized similarly as compound **23a**.

N⁴-Benzyl-3-nitro-6-phenyl-N²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20b). Phenylboronic acid was used as a reagent. Yellow solid (28 mg, 57%). ¹H NMR (500 MHz, CDCl₃) δ 9.68 (s, 1H), 9.54 (s, 1H), 7.92 (ddd, *J* = 4.4, 2.5, 1.4

Hz, 2H), 7.45 – 7.42 (m, 3H), 7.40 – 7.38 (m, 3H), 7.34 – 7.31 (m, 1H), 7.26 (s, 1H), 6.39 (s, 1H), 4.58 (d, *J* = 5.4 Hz, 2H), 2.09 (s, 2H), 1.64 (s, 6H), 0.99 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 159.01, 154.74, 153.13, 138.88, 136.87, 130.20, 129.19, 128.66, 128.06, 127.50, 127.44, 92.16, 56.72, 51.69, 47.62, 32.02, 31.74, 30.15. MS (ESI) calculated for C₂₆H₃₃N₄O₂, *m/z* 433.2598, found 433.2615 (M + H)⁺.

1-Benzyl-2-butyl-6-phenyl-N-(2,4,4-trimethylpentan-2-yl)-1H-imidazo[4,5-c]pyridin-4-amine (22b). Yellow solid (31 mg, 47%). ¹H NMR (500 MHz, CDCl₃) δ 8.02 (dd, *J* = 8.3, 1.2 Hz, 2H), 7.39 (t, *J* = 7.7 Hz, 2H), 7.34 – 7.27 (m, 4H), 7.07 (d, *J* = 6.7 Hz, 2H), 6.94 (s, 1H), 5.34 (s, 1H), 5.28 (s, 2H), 2.79 – 2.73 (m, 2H), 2.18 (s, 2H), 1.73 – 1.66 (m, 8H), 1.39 (dd, *J* = 15.0, 7.5 Hz, 2H), 1.03 (s, 9H), 0.89 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 153.22, 150.01, 148.43, 141.28, 140.32, 136.16, 129.61, 129.42, 129.15, 128.41, 128.07, 127.67, 127.54, 126.72, 126.36, 126.28, 126.22, 91.77, 55.45, 51.63, 47.28, 31.98, 31.81, 30.33, 30.25, 27.50, 22.72, 13.90. MS (ESI) calculated for C₃₁H₄₁N₄, *m/z* 469.3326, found 469.3350 (M + H)⁺.

1-Benzyl-2-butyl-6-phenyl-1H-imidazo[4,5-c]pyridin-4-amine (23b). White solid (13 mg, 73%). ¹H NMR (500 MHz, CDCl₃) δ 7.91 (d, *J* = 7.2 Hz, 2H), 7.54 – 7.44 (m, 3H), 7.42 – 7.34 (m, 3H), 7.05 (dd, *J* = 6.4, 1.1 Hz, 2H), 6.83 (s, 1H), 5.37 (s, 2H), 2.88 – 2.78 (m, 2H), 1.78 (dd, *J* = 15.2, 7.8 Hz, 2H), 1.42 (dq, *J* = 14.7, 7.4 Hz, 2H), 0.93 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 157.54, 149.18, 142.11, 141.93, 134.08, 132.07, 130.82, 129.68, 129.66, 129.01, 127.38, 126.25, 95.25, 48.13, 29.31, 27.40, 22.52, 13.84. HRMS (ESI) calculated for C₂₃H₂₅N₄, *m/z* 357.2074, found 357.2155 (M + H)⁺.

4-(4-(Benzylamino)-5-nitro-6-((2,4,4-trimethylpentan-2-yl)amino)pyridin-2-yl)benzonitrile (20c). 4-Cyanophenylboronic acid was used as a reagent. Yellow solid (30 mg, 55%). ¹H NMR (500 MHz, CDCl₃) δ 9.72 (s, 1H), 9.51 (s, 1H), 7.97 (d, *J* = 8.6 Hz, 2H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.44 – 7.31 (m, 5H), 6.38 (s, 1H), 4.59 (d, *J* = 5.4 Hz, 2H), 2.06 (s, 2H), 1.63 (s, 6H), 0.99 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 156.75, 154.69, 153.26, 143.14, 136.53, 132.50, 129.28, 128.23, 127.93, 127.35, 118.79, 113.38, 93.05, 56.87, 51.72, 47.68, 32.01, 31.72, 30.10. MS (ESI) calculated for C₂₇H₃₂N₅O₂, *m/z* 458.2551, found 458.2571 (M + H)⁺.

4-(1-Benzyl-2-butyl-4-((2,4,4-trimethylpentan-2-yl)amino)-1H-imidazo[4,5-c]pyridin-6-yl)benzamide (22c). Cyano group was converted into amide in the basic condition. Yellow solid (22 mg, 41%). ¹H NMR (500 MHz, CDCl₃) δ 8.10 (d, *J* = 8.5 Hz, 2H), 7.84 (d, *J* = 8.5 Hz, 2H), 7.34 – 7.28 (m, 3H), 7.08 – 7.04 (m, 2H), 6.99 (s, 1H), 6.14 (s, 1H), 5.79 (s, 1H), 5.41 (s, 1H), 5.29 (s, 2H), 2.81 – 2.74 (m, 2H), 2.17 (s, 2H), 1.72 – 1.66 (m, 8H), 1.43 – 1.33 (m, 2H), 1.03 (s, 9H), 0.89 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 169.46, 153.74, 150.08, 147.00, 144.87, 140.11, 135.97, 131.83, 129.20, 128.15, 127.62, 126.76, 126.74, 126.31, 92.63, 55.50, 51.55, 47.31, 31.95, 31.79, 30.25, 30.17, 27.46, 22.69, 13.89. MS (ESI) calculated for C₃₂H₄₂N₅O, *m/z* 512.3384, found 512.3408 (M + H)⁺.

4-(4-Amino-1-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-6-yl)benzamide (23c). White solid (11 mg, 79%). ¹H NMR (500 MHz, MeOD) δ 8.09 – 8.05 (m, 2H), 7.89 – 7.85 (m, 2H), 7.51 (s, 1H), 7.41 – 7.32 (m, 3H), 7.17 (d, *J* = 7.0 Hz, 2H), 5.62 (s, 2H), 3.76 – 3.72 (m, 1H), 3.68 – 3.64 (m, 1H), 3.58 (dd, *J* = 7.0, 2.7 Hz, 1H), 2.94 – 2.89 (m, 2H), 1.78 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.42 (dd, *J* = 15.0, 7.5 Hz, 2H), 0.92 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 171.03, 159.91, 149.79, 143.90, 141.45, 137.23, 136.75, 136.73, 130.29, 129.74, 129.43, 128.43, 127.76, 125.80, 98.94, 73.57, 72.45, 62.18, 43.73, 30.01, 27.99, 23.28, 14.04. HRMS (ESI) calculated for C₂₄H₂₆N₅O, *m/z* 400.2132, found 400.2177 (M + H)⁺.

***N*⁴-Benzyl-5-nitro-*N*⁶-(2,4,4-trimethylpentan-2-yl)-[2,3'-bipyridine]-4,6-diamine (20d).** 4-Pyridinylboronic acid was used as a reagent. Orange solid (135 mg, 82%). ¹H NMR (500 MHz, CDCl₃) δ 9.74 (t, *J* = 5.1 Hz, 1H), 9.54 (s, 1H), 9.10 (dd, *J* = 2.2, 0.6 Hz, 1H), 8.65 (dd, *J* = 4.8, 1.6 Hz, 1H), 8.21 – 8.16 (m, 1H), 7.42 – 7.36 (m, 5H), 7.35 – 7.31 (m, 1H), 6.38 (s, 1H), 4.59 (d, *J* = 5.5 Hz, 2H), 2.07 (s, 2H), 1.63 (s, 6H), 0.99 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 156.52, 154.78, 153.22, 150.86, 148.93, 136.60, 134.75, 134.36, 129.26, 128.17, 127.35, 123.54, 116.37, 92.38, 56.86, 51.65, 47.65, 32.01, 31.72, 30.11. MS (ESI) calculated for C₂₅H₃₂N₅O₂, *m/z* 434.2551, found 434.2563 (M + H)⁺.

1-Benzyl-2-butyl-6-(pyridin-3-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23d). Light yellow solid (10 mg, 72%). ¹H NMR (500 MHz, MeOD) δ 9.28 (d, *J* = 1.9 Hz, 1H), 8.94 (dd, *J* = 5.5, 1.2 Hz, 1H), 8.83 (d, *J* = 8.2 Hz, 1H), 8.11 (ddd, *J* = 8.2, 5.6, 0.5 Hz, 1H), 7.72 (s, 1H), 7.40 – 7.32 (m, 3H), 7.18 (d, *J* = 6.9 Hz, 2H), 5.65 (s, 2H), 2.98 – 2.91 (m, 2H), 1.78 (dd, *J* = 15.3, 7.7 Hz, 2H), 1.42 (dd, *J* = 15.0, 7.5 Hz, 2H), 0.92 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 150.19, 143.97, 143.57, 143.11, 136.48, 130.41, 130.32, 129.52, 128.03, 127.74, 127.73, 100.47, 48.55, 29.93, 27.90, 23.26, 14.02. HRMS (ESI) calculated for C₂₂H₂₄N₅, *m/z* 358.2026, found 358.2067 (M + H)⁺.

***N*⁴-Benzyl-6-(furan-3-yl)-3-nitro-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20e).** 3-Furylboronic acid was used as a reagent. Yellow solid (95 mg, 70%). ¹H NMR (500 MHz, CDCl₃) δ 9.68 (t, *J* = 4.8 Hz, 1H), 9.49 (s, 1H), 7.95 (dd, *J* = 1.5, 0.8 Hz, 1H), 7.45 (t, *J* = 1.7 Hz, 1H), 7.41 – 7.34 (m, 4H), 7.34 – 7.30 (m, 1H), 6.69 (dd, *J* = 1.9, 0.8 Hz, 1H), 6.07 (s, 1H), 4.53 (d, *J* = 5.4 Hz, 2H), 2.05 (s, 2H), 1.60 (s, 6H), 0.98 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 154.88, 153.66, 153.09, 143.99, 143.68, 136.85, 129.19, 128.06, 127.39, 127.36, 108.83, 91.66, 56.68, 51.51, 47.61, 31.99, 31.71, 30.13. MS (ESI) calculated for C₂₄H₃₁N₄O₃, *m/z* 423.2391, found 423.2401 (M + H)⁺.

1-Benzyl-2-butyl-6-(furan-3-yl)-*N*-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (22e). Brown oil (30 mg, 41%). ¹H NMR (500 MHz, CDCl₃) δ 7.94 (dd, *J* = 1.6, 0.7 Hz, 1H), 7.41 (t, *J* = 1.7 Hz, 1H), 7.34 – 7.27 (m, 3H), 7.05 (d, *J* = 6.6 Hz, 2H), 6.76 (dd, *J* = 1.8, 0.8 Hz, 1H), 6.61 (s, 1H), 5.30 (s, 1H), 5.24 (s, 2H), 2.76 – 2.71 (m, 2H), 2.13 (s, 2H), 1.70 – 1.65 (m, 2H), 1.64 (s, 6H), 1.36 (dt, *J* = 14.7, 7.4 Hz, 2H), 1.02 (s, 9H),

0.88 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 152.80, 150.15, 143.24, 142.70, 140.71, 140.01, 136.16, 129.15, 128.83, 128.06, 126.33, 125.88, 108.87, 91.20, 55.44, 51.45, 47.25, 31.93, 31.79, 30.29, 30.16, 27.45, 22.71, 13.90. MS (ESI) calculated for C₂₉H₃₉N₄O, *m/z* 459.3118, found 459.3168 (M + H)⁺.

1-Benzyl-2-butyl-6-(furan-3-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23e). Light yellow solid (12 mg, 64%). ¹H NMR (500 MHz, MeOD) δ 8.23 – 8.19 (m, 1H), 7.73 – 7.70 (m, 1H), 7.42 (s, 1H), 7.40 – 7.32 (m, 3H), 7.15 (d, *J* = 6.9 Hz, 2H), 6.97 (dd, *J* = 2.0, 0.9 Hz, 1H), 5.58 (s, 2H), 2.90 – 2.85 (m, 2H), 1.75 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.40 (dq, *J* = 14.8, 7.4 Hz, 2H), 0.91 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 159.50, 149.42, 146.46, 144.24, 142.92, 136.72, 135.38, 130.27, 129.41, 127.69, 124.89, 121.45, 109.21, 96.82, 48.27, 29.95, 27.91, 23.26, 14.03. HRMS (ESI) calculated for C₂₁H₂₃N₄O, *m/z* 347.1866, found 347.1921 (M + H)⁺.

***N*⁴-Benzyl-3-nitro-6-(thiophen-3-yl)-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20f).** 3-Thienylboronic acid was used as a reagent. Yellow solid (107 mg, 76%). ¹H NMR (500 MHz, CDCl₃) δ 9.69 (s, 1H), 9.52 (s, 1H), 7.89 (dd, *J* = 3.0, 1.2 Hz, 1H), 7.49 (dd, *J* = 5.1, 1.2 Hz, 1H), 7.42 – 7.30 (m, 6H), 6.26 (s, 1H), 4.56 (d, *J* = 5.4 Hz, 2H), 2.08 (s, 2H), 1.63 (s, 6H), 0.99 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 154.88, 154.61, 153.25, 142.21, 136.90, 129.19, 128.05, 127.38, 126.59, 126.56, 126.28, 91.99, 56.68, 51.59, 47.61, 32.02, 31.72, 30.16. MS (ESI) calculated for C₂₄H₃₁N₄O₂S, *m/z* 439.2162, found 439.2183 (M + H)⁺.

1-Benzyl-2-butyl-6-(thiophen-3-yl)-*N*-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (22f). Brown solid (25 mg, 35%). ¹H NMR (500 MHz, CDCl₃) δ 7.79 (dd, *J* = 3.1, 1.2 Hz, 1H), 7.58 (dd, *J* = 5.0, 1.2 Hz, 1H), 7.33 – 7.27 (m, 4H), 7.06 (d, *J* = 6.6 Hz, 2H), 6.79 (s, 1H), 5.35 (s, 1H), 5.25 (s, 2H), 2.78 – 2.70 (m, 2H), 2.17 (s, 2H), 1.70 (dd, *J* = 8.7, 7.0 Hz, 3H), 1.67 (s, 6H), 1.38 (dq, *J* = 14.7, 7.4 Hz, 2H), 1.26 (dd, *J* = 8.7, 5.6 Hz, 1H), 1.02 (s, 9H), 0.89 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 153.04, 150.02, 144.89, 144.32, 140.11, 136.13, 129.14, 128.07, 126.45, 126.34, 125.94, 125.41, 121.55, 91.54, 55.43, 51.49, 50.96, 47.26, 31.96, 31.80, 30.26, 30.21, 27.46, 22.71, 13.89. MS (ESI) calculated for C₂₉H₃₉N₄S, *m/z* 475.2890, found 475.2922 (M + H)⁺.

1-Benzyl-2-butyl-6-(thiophen-3-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23f). White solid (9 mg, 53%). ¹H NMR (500 MHz, MeOD) δ 8.04 (dd, *J* = 2.9, 1.4 Hz, 1H), 7.66 (dd, *J* = 5.1, 2.9 Hz, 1H), 7.58 (dd, *J* = 5.1, 1.4 Hz, 1H), 7.48 (s, 1H), 7.41 – 7.32 (m, 3H), 7.17 (d, *J* = 6.9 Hz, 2H), 5.59 (s, 2H), 2.94 – 2.86 (m, 2H), 1.76 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.41 (dq, *J* = 14.8, 7.4 Hz, 2H), 0.91 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 159.63, 149.38, 144.24, 137.90, 136.73, 135.24, 130.28, 129.42, 129.31, 127.74, 126.73, 126.03, 124.88, 97.21, 49.07, 29.96, 27.94, 23.26, 14.03. HRMS (ESI) calculated for C₂₁H₂₃N₄S, *m/z* 363.1638, found 363.1686 (M + H)⁺.

***N*^{4,6}-Dibenzyl-3-nitro-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20g).**

Benzylboronic acid pinacol ester was used as a reagent. Orange oil (135 mg, 91%). ¹H NMR (500 MHz, CDCl₃) δ 9.54 (t, *J* = 5.0 Hz, 1H), 9.37 (s, 1H), 7.36 (ddd, *J* = 7.4, 4.5, 1.4 Hz, 2H), 7.31 (dt, *J* = 9.8, 4.4 Hz, 1H), 7.27 (ddd, *J* = 6.7, 3.3, 1.4 Hz, 5H), 7.22 – 7.18 (m, 3H), 5.75 (s, 1H), 4.40 (d, *J* = 5.4 Hz, 2H), 3.75 (s, 2H), 1.89 (s, 2H), 1.49 (s, 6H), 0.90 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 164.70, 154.68, 152.84, 138.77, 136.80, 129.45, 129.07, 128.54, 127.91, 127.46, 126.50, 124.96, 115.72, 94.56, 56.63, 51.05, 47.44, 45.61, 31.87, 31.60, 30.09, 24.86. MS (ESI) calculated for C₂₇H₃₅N₄O₂, *m/z* 447.2755, found 447.2737 (M + H)⁺.

1,6-Dibenzyl-2-butyl-*N*-(2,4,4-trimethylpentan-2-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (22g).

Light brown solid (55 mg, 60%). ¹H NMR (500 MHz, CDCl₃) δ 7.33 – 7.27 (m, 5H), 7.24 (t, *J* = 7.6 Hz, 2H), 7.14 (t, *J* = 7.3 Hz, 1H), 7.01 (d, *J* = 6.4 Hz, 2H), 6.31 (s, 1H), 5.22 (s, 1H), 5.16 (s, 2H), 3.96 (s, 2H), 2.74 – 2.67 (m, 2H), 2.02 (s, 2H), 1.66 – 1.61 (m, 2H), 1.56 (s, 6H), 1.35 (dd, *J* = 15.0, 7.5 Hz, 2H), 0.93 (s, 9H), 0.86 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, CDCl₃) δ 152.37, 151.90, 150.06, 141.57, 140.10, 136.28, 129.31, 129.06, 128.21, 127.95, 126.38, 125.78, 124.94, 93.59, 55.35, 50.98, 47.19, 45.39, 31.85, 31.68, 30.41, 30.27, 27.42, 22.67, 13.87. MS (ESI) calculated for C₃₂H₄₃N₄, *m/z* 483.3482, found 483.3526 (M + H)⁺.

1,6-Dibenzyl-2-butyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23g).

White solid (23 mg, 58%). ¹H NMR (500 MHz, DMSO) δ 13.63 (s, 1H), 8.27 (s, 2H), 7.37 – 7.29 (m, 7H), 7.27 – 7.23 (m, 1H), 7.15 (s, 1H), 7.11 (d, *J* = 6.9 Hz, 2H), 5.52 (s, 2H), 4.08 (s, 2H), 2.86 – 2.79 (m, 2H), 1.61 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.31 (dt, *J* = 14.9, 7.4 Hz, 2H), 0.81 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 156.72, 148.03, 142.12, 141.72, 137.39, 135.93, 128.96, 128.72, 128.70, 127.92, 126.98, 126.63, 123.29, 97.60, 46.84, 40.11, 38.20, 28.89, 26.28, 21.69, 13.61. HRMS (ESI) calculated for C₂₄H₂₇N₄, *m/z* 371.2230, found 371.2285 (M + H)⁺.

***N*⁴-Benzyl-6-(4-methylbenzyl)-3-nitro-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20h).**

4-Methylbenzylboronic acid pinacol ester was used as a reagent. Yellow oil (130 mg, 91%). ¹H NMR (500 MHz, CDCl₃) δ 9.52 (t, *J* = 5.0 Hz, 1H), 9.37 (s, 1H), 7.37 – 7.30 (m, 3H), 7.28 – 7.24 (m, 3H), 7.10 – 7.03 (m, 7H), 5.74 (s, 1H), 4.39 (d, *J* = 5.4 Hz, 2H), 3.70 (s, 2H), 2.31 (s, 3H), 1.90 (s, 2H), 1.51 (s, 6H), 0.90 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 165.00, 154.68, 152.83, 136.82, 135.98, 135.70, 135.50, 134.26, 129.28, 129.23, 129.12, 129.05, 128.98, 127.89, 127.47, 115.70, 94.47, 83.50, 56.63, 51.03, 47.44, 45.22, 31.87, 31.57, 30.12, 24.87, 21.19, 21.11. MS (ESI) calculated for C₂₈H₃₇N₄O₂, *m/z* 461.2911, found 461.3003 (M + H)⁺.

1-Benzyl-2-butyl-6-(4-methylbenzyl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23h).

White solid (19 mg, 66%). ¹H NMR (500 MHz, MeOD) δ 7.38 – 7.32 (m, 3H), 7.16 (d, *J* = 7.9 Hz, 2H), 7.13 – 7.08 (m, 4H), 6.89 (s, 1H), 5.49 (s, 2H), 4.08 (s, 2H), 2.92 – 2.87 (m, 2H), 2.32 (s, 3H), 1.76 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.40 (dt, *J* = 14.8, 7.4 Hz, 2H), 0.91 (t, *J* = 7.4 Hz, 3H). ¹³C NMR

(126 MHz, MeOD) δ 159.11, 149.35, 144.27, 143.93, 138.37, 136.53, 134.52, 130.68, 130.25, 129.85, 129.42, 127.74, 124.41, 99.41, 39.40, 30.01, 27.84, 23.26, 21.08, 14.03. HRMS (ESI) calculated for C₂₅H₂₉N₄, *m/z* 385.2387, found 385.2451 (M + H)⁺.

***N*⁴-Benzyl-3-nitro-6-(4-(trifluoromethoxy)benzyl)-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20i).**

4-(Trifluoromethoxy)benzylboronic acid pinacol ester was used as a reagent. Yellow solid (115 mg, 70%). ¹H NMR (500 MHz, CDCl₃) δ 9.57 (t, *J* = 5.0 Hz, 1H), 9.37 (s, 1H), 7.38 – 7.29 (m, 3H), 7.28 (t, *J* = 1.8 Hz, 2H), 7.21 – 7.17 (m, 2H), 7.10 (d, *J* = 7.9 Hz, 2H), 5.74 (s, 1H), 4.43 (d, *J* = 5.5 Hz, 2H), 1.83 (s, 2H), 1.45 (s, 6H), 0.88 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 163.87, 154.70, 152.91, 137.54, 136.71, 130.77, 129.10, 127.95, 127.33, 121.09, 115.75, 94.57, 56.63, 50.96, 47.44, 44.69, 31.82, 31.55, 30.02. MS (ESI) calculated for C₂₈H₃₄F₃N₄O₃, *m/z* 531.2578, found 531.2738 (M + H)⁺.

1-Benzyl-2-butyl-6-(4-(trifluoromethoxy)benzyl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23i).

White solid (21 mg, 47%). ¹H NMR (500 MHz, CDCl₃) δ 7.37 – 7.29 (m, 5H), 7.21 (d, *J* = 8.0 Hz, 2H), 7.08 (d, *J* = 6.5 Hz, 2H), 6.79 (s, 1H), 5.45 (s, 2H), 4.09 (s, 2H), 2.87 (dd, *J* = 9.3, 6.1 Hz, 2H), 1.74 (dt, *J* = 15.3, 7.6 Hz, 2H), 1.41 (dt, *J* = 15.0, 7.4 Hz, 2H), 0.91 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 157.96, 150.50, 149.43, 149.42, 143.48, 138.54, 136.96, 131.59, 130.16, 129.25, 127.66, 124.97, 122.37, 99.02, 40.76, 30.29, 27.86, 23.31, 14.05. HRMS (ESI) calculated for C₂₅H₂₆F₃N₄O, *m/z* 455.2053, found 455.2131 (M + H)⁺.

***N*⁴-Benzyl-3-nitro-6-phenethyl-*N*²-(2,4,4-trimethylpentan-2-yl)pyridine-2,4-diamine (20j).**

2-Phenylethylboronic acid was used as a reagent. Yellow oil (85 mg, 59%). ¹H NMR (500 MHz, CDCl₃) δ 9.52 (t, *J* = 4.9 Hz, 1H), 9.43 (s, 1H), 7.36 (t, *J* = 7.2 Hz, 2H), 7.33 – 7.28 (m, 4H), 7.27 – 7.25 (m, 3H), 7.19 (d, *J* = 7.3 Hz, 1H), 7.17 (d, *J* = 7.0 Hz, 2H), 5.69 (s, 1H), 4.39 (d, *J* = 5.4 Hz, 2H), 2.99 (dd, *J* = 9.1, 6.8 Hz, 2H), 2.75 (dd, *J* = 9.1, 6.8 Hz, 2H), 2.04 (s, 2H), 1.58 (s, 6H), 0.98 (s, 9H). ¹³C NMR (126 MHz, CDCl₃) δ 165.06, 154.74, 152.65, 141.77, 136.85, 129.09, 128.52, 128.49, 127.95, 127.39, 126.10, 115.78, 94.52, 56.68, 51.35, 47.41, 40.87, 34.61, 31.97, 31.69, 30.20. MS (ESI) calculated for C₂₈H₃₇N₄O₂, *m/z* 461.2911, found 461.3096 (M + H)⁺.

1-Benzyl-2-butyl-6-phenethyl-1*H*-imidazo[4,5-*c*]pyridin-4-amine (23j).

Yellow solid (11 mg, 42%). ¹H NMR (500 MHz, CDCl₃) δ 7.36 – 7.30 (m, 3H), 7.23 – 7.18 (m, 2H), 7.13 (ddd, *J* = 5.7, 3.7, 1.5 Hz, 3H), 7.02 (d, *J* = 6.5 Hz, 2H), 6.68 (s, 1H), 5.38 (s, 2H), 2.99 (s, 4H), 2.85 – 2.79 (m, 2H), 1.69 (ddd, *J* = 13.2, 8.5, 6.7 Hz, 2H), 1.37 (dq, *J* = 14.8, 7.4 Hz, 2H), 0.89 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, MeOD) δ 157.35, 150.48, 148.64, 143.39, 141.89, 137.14, 130.14, 129.53, 129.43, 129.16, 127.59, 127.22, 124.68, 97.93, 48.17, 38.42, 36.84, 30.28, 27.83, 23.31, 14.03. HRMS (ESI) calculated for C₂₅H₂₉N₄, *m/z* 385.2387, found 385.2446 (M + H)⁺.

Synthesis of compound 24: 3-Nitrobenzo[*g*]quinolin-4-ol.

Nitromethane (0.96 mL, 18 mmol) was added dropwise to a solution of NaOH (2.2 g, 54 mmol) in water (5 mL), at 0 °C. The mixture was then warmed to 40 °C and nitromethane (0.96 mL, 18 mmol) was again added slowly at 40-45 °C. The temperature was maintained until a clear solution was obtained. The reaction mixture was then heated to 55 °C for 2-5 minutes, cooled to 30 °C, poured onto crushed ice and acidified with conc. HCl (5 mL). The resultant solution of methazoic acid was added immediately to a filtered solution of 3-amino-2-naphtholic acid (3 g, 16 mmol) and conc. HCl (1 mL) in water (20 mL). The reaction mixture was allowed to stand at room temperature for 12 h. After filtration, the residue obtained was washed with water, and dried (1.1 g, 90%). A solution of intermediate (2 g, 7.75 mmol) in acetic anhydride (10 mL) was placed in a 2-neck flask fitted with a reflux condenser. It was stirred and heated to 105 °C until a clear solution was obtained. Heating was then discontinued and potassium acetate (0.77 g, 7.90 mmol) was added. The mixture was then refluxed for 15 min with vigorous stirring, until a solid started to precipitate. The reaction mixture was then slowly cooled to room temperature. The residue was filtered, washed with glacial acetic acid until the washings were colorless, then suspended in water, filtered, washed with water and dried at 110 °C to get compound **24** (0.93 g, 50%). ¹H NMR (500 MHz, DMSO) δ 13.12 (s, 1H), 9.28 (s, 1H), 8.95 (s, 1H), 8.27 – 8.21 (m, 2H), 8.11 (d, *J* = 8.4 Hz, 1H), 7.69 (t, *J* = 7.1 Hz, 1H), 7.61 (t, *J* = 7.1 Hz, 1H). ¹³C NMR (126 MHz, DMSO) δ 168.61, 144.40, 135.05, 134.63, 130.14, 129.54, 128.94, 128.71, 127.46, 127.41, 126.65, 126.48, 116.78. MS (ESI) calculated for C₁₃H₈N₂O₃, *m/z* 240.05, found 263.04 (M + Na)⁺.

Synthesis of compound 25: 4-Chloro-3-nitrobenzo[g]quinoline. A suspension of compound **24** (2.0 g, 8.30 mmol) in phosphorus(V) oxychloride was placed in a pressure vessel and it was heated at 150 °C. After a clear solution was obtained, the reaction mixture was kept at 150 °C for 1 h. Then it was slowly cooled to room temperature and the solvent was evaporated under vacuum. The residue was poured over crushed ice while stirring and the formed solid was filtered, washed with water and dried to obtain compound **25** (1.95 g, 91%). ¹H NMR (500 MHz, DMSO) δ 13.34 (d, *J* = 6.9 Hz, 1H), 9.26 (d, *J* = 7.3 Hz, 1H), 8.95 (s, 1H), 8.25 (d, *J* = 9.4 Hz, 2H), 8.11 (d, *J* = 8.4 Hz, 1H), 7.72 – 7.67 (m, 1H), 7.61 (ddd, *J* = 8.1, 6.8, 1.1 Hz, 1H). ¹³C NMR (126 MHz, DMSO) δ 168.64, 144.23, 134.96, 134.64, 130.16, 129.56, 128.98, 128.70, 127.48, 127.43, 126.69, 126.48, 116.74.

Synthesis of compound 26: *N*-Benzyl-3-nitrobenzo[g]quinolin-4-amine. To a solution of compound **25** (1.0 g, 3.90 mol) in 20 mL of CH₂Cl₂ was added triethylamine (0.81 mL, 5.80 mmol) and benzylamine (0.50 mL, 4.60 mmol). The reaction mixture was refluxed for 2 h. The solvent was then evaporated under vacuum and H₂O was added to the residue. The solution was extracted with CH₂Cl₂ (3 × 20 mL), washed with water and dried over sodium sulfate. The solvent was evaporated and the residue was purified using silica gel column chromatography (5% MeOH/CH₂Cl₂) to obtain compound **26** as a yellow solid (1.1 g, 88%). ¹H NMR (500 MHz, CDCl₃) δ 10.57 (s, 1H), 9.41 (s, 1H), 8.87 (s, 1H), 8.50 (s, 1H), 8.02 (d, *J* = 8.4 Hz, 1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.65 – 7.60 (m, 1H), 7.55 – 7.51 (m, 1H), 7.50 –

7.47 (m, 4H), 7.42 (ddd, *J* = 11.0, 5.4, 3.0 Hz, 1H), 5.28 (d, *J* = 5.9 Hz, 2H). ¹³C NMR (126 MHz, CDCl₃) δ 152.50, 147.73, 145.88, 136.85, 135.48, 130.53, 129.64, 129.12, 129.03, 128.98, 128.83, 128.11, 128.04, 127.29, 126.86, 124.21, 118.09, 53.19. MS (ESI) calculated for C₂₀H₁₆N₃O₂, *m/z* 330.1237, found 330.1304 (M + H)⁺.

Synthesis of compound 30: 1-Benzyl-2-butyl-1*H*-benzo[g]imidazo[4,5-*c*]quinolin-4-amine. To a solution of compound **26** (300 mg, 0.91 mmol) in 20 mL of MeOH were added zinc dust (594 mg, 9.10 mmol) and ammonium formate (574 mg, 9.10 mmol). The reaction mixture was stirred at room temperature for 30 min and filtered through celite. Then the solvent was evaporated and the residue was dissolved in water. This was extracted with EtOAc (3 × 20 mL), washed with water and dried over sodium sulfate. The solvent was removed under vacuum to obtain compound **27** (100 mg, 37%). To a solution of compound **27** (98 mg, 0.33 mmol) in 10 mL of anhydrous THF were added triethylamine (48 μL, 0.35 mmol) and valeryl chloride (40 μL, 0.33 mmol). The reaction mixture was refluxed for 2 h. The solvent was then removed under vacuum, and the residue was dissolved in 10 mL of EtOH and NaOH (26 mg, 0.66 mmol) in 1 mL of H₂O was added. The reaction mixture was refluxed for 2 h. The solvent was then removed under vacuum, and the residue was dissolved in EtOAc and washed with water. The EtOAc fraction was dried using sodium sulfate and evaporated and purified using column chromatography (10% MeOH/CH₂Cl₂) to obtain the compound **28** (76 mg, 63%). To a solution of compound **28** (76 mg, 0.21 mmol) in a solvent mixture of MeOH:CH₂Cl₂:CHCl₃ (1:10:10) was added 3-chloroperoxy benzoic acid (443 mg, 1.98 mmol), and the solution was refluxed at 45-50 °C for 1 h. The solvent was then removed and the residue was purified using column chromatography (10% MeOH/CH₂Cl₂) to obtain the *N*-oxide derivative **29** (64 mg, 80%). To a solution of compound **29** (64 mg, 0.17 mol) in 10 mL of CH₂Cl₂ was added benzoyl isocyanate (37 mg, 0.25 mmol) and heated at 45 °C for 18 h. The solvent was then removed under vacuum, and the residue was dissolved in 15 mL of anhydrous MeOH, followed by the addition of excess sodium methoxide. The reaction mixture was then heated at 80 °C for 2 h. The solvent was removed under vacuum and the residue was purified using column chromatography (10% MeOH/CH₂Cl₂) to obtain the compound **30** (20 mg, 30%). ¹H NMR (500 MHz, DMSO) δ 8.40 (s, 1H), 8.05 (s, 1H), 7.88 (d, *J* = 8.3 Hz, 1H), 7.77 (d, *J* = 8.3 Hz, 1H), 7.38 (d, *J* = 7.2 Hz, 1H), 7.33 (dd, *J* = 12.9, 5.2 Hz, 3H), 7.24 (d, *J* = 7.4 Hz, 1H), 7.16 (d, *J* = 7.4 Hz, 2H), 6.91 (s, 2H), 6.02 (s, 2H), 3.00 – 2.92 (m, 2H), 1.74 (dt, *J* = 15.4, 7.6 Hz, 2H), 1.40 (dd, *J* = 14.9, 7.4 Hz, 2H), 0.88 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (126 MHz, DMSO) δ 136.73, 131.80, 128.91, 127.99, 127.81, 127.46, 126.82, 125.81, 123.63, 118.72, 48.20, 29.71, 26.27, 21.85, 13.69. HRMS (ESI) calculated for C₂₅H₂₅N₄, *m/z* 381.2074, found 381.2089 (M + H)⁺.

Human TLR-7/-8 Reporter Gene assays (NF-κB induction). The induction of NF-κB was quantified using HEK-Blue-7 (hTLR7-specific) and HEK-Blue-8 (hTLR8-specific) cells as previously described by us.^{33,45,46} HEK293 cells stably co-transfected with human TLR7 or human TLR8, MD2, and

secreted alkaline phosphatase (sAP), were maintained in HEK-Blue™ Selection medium containing zeocin and normocin. Stable expression of secreted alkaline phosphatase (sAP) under control of NF-κB/AP-1 promoters is inducible by appropriate TLR agonists, and extracellular sAP in the supernatant is proportional to NF-κB induction. HEK-Blue cells were incubated at a density of ~10⁵ cells/ml in a volume of 80 μl/well, in 384-well, flat-bottomed, cell culture-treated microtiter plates until confluency was achieved, and subsequently stimulated with graded concentrations of stimuli. sAP was assayed spectrophotometrically using an alkaline phosphatase-specific chromogen (present in HEK-detection medium as supplied by the vendor) at 620 nm.

Immunoassays for Interferon (IFN)-α, and cytokines. Fresh human peripheral blood mononuclear cells (hPBMC) were isolated from human blood obtained by venipuncture with informed consent and as per institutional guidelines on Ficoll-Hypaque gradients as described elsewhere.⁶⁴ Aliquots of PBMCs (10⁵ cells in 100 μL/well) were stimulated for 12 h with graded concentrations of test compounds. Supernatants were isolated by centrifugation, and were assayed in triplicates using either high-sensitivity multi-subtype IFN-α ELISA kits (PBL Interferon Source, Piscataway, NJ and R&D Systems, Inc., Minneapolis, MN), or analyte-specific multiplexed cytokine/chemokine bead array assays as reported by us previously.⁶⁵

Flow-cytometric immunostimulation experiments. CD69 upregulation was determined by flow cytometry using protocols published by us previously,³³ and modified for rapid-throughput. Briefly, heparin-anticoagulated whole blood samples were obtained by venipuncture from healthy human volunteers with informed consent and as per guidelines approved by the University of Kansas Human Subjects Experimentation Committee. Serial dilutions of selected imidazopyridine compounds (and imiquimod, used as a reference compound) were performed using a Bio-Tek Precision 2000 XS liquid handler in sterile 96-well polypropylene plates, to which were added 100 μL aliquots of anticoagulated whole human blood. The plates were incubated at 37°C for 16.5 h. Negative (endotoxin free water) controls were included in each experiment. Following incubation, fluorochrome-conjugated antibodies (CD3-PE, CD56-APC, CD69-PE-Cy7, 10 μL of each antibody, Becton-Dickinson Biosciences, San Jose, CA) were added to each well with a liquid handler, and incubated at 37 °C in the dark for 30 min. Following staining, erythrocytes were lysed and leukocytes fixed by mixing 200 L of the samples in 2 mL pre-warmed Whole Blood Lyse/Fix Buffer (Becton-Dickinson Biosciences, San Jose, CA) in 96 deep-well plates. After washing the cells twice at 200 g for 8 minutes in saline, the cells were transferred to a 96-well plate. Flow cytometry was performed using a BD FACSAria instrument in the tri-color mode (tri-color flow experiment) and two-color mode (two-color flow experiment) for acquisition on 100,000 gated events. Compensation for spillover was computed for each experiment on singly-stained samples. CD69 activation in the major lymphocytic populations, viz., natural killer lymphocytes (NK cells: CD3⁺CD56⁺), cytokine-induced killer phenotype (CIK cells: CD3⁺CD56⁺), nominal B lymphocytes (CD3⁻CD56⁻), and

nominal T lymphocytes (CD3⁺CD56⁻) were quantified using FlowJo v 7.0 software (Treestar, Ashland, OR).

Acknowledgments

This work was supported by NIH/NIAID contract HSN272200900033C.

Notes and references

Department of Medicinal Chemistry, University of Kansas, Multidisciplinary Research Building, Room 320D, 2030 Becker Drive, Lawrence KS 66047. Tel: 785-864-1610; Fax: 785-864-1961
E-mail: sdavid@ku.edu

† Electronic Supplementary Information (ESI) available: Characterization of intermediates and final compounds (¹H, ¹³C, mass spectra).

References

1. A. Iwasaki and R. Medzhitov, *Science*, 2010, **327**, 291.
2. A. Iwasaki and R. Medzhitov, *Nat. Immunol.*, 2004, **5**, 987.
3. C. A. Janeway, Jr. and R. Medzhitov, *Annu. Rev. Immunol.*, 2002, **20**, 197.
4. M. S. Lee and Y. J. Kim, *Mol. Cells*, 2007, **23**, 1.
5. B. A. Seaton, E. C. Crouch, F. X. McCormack, J. F. Head, K. L. Hartshorn, and R. Mendelsohn, *Innate. Immun.*, 2010, **16**, 143.
6. V. L. Runza, W. Schwaeble, and D. N. Mannel, *Immunobiol.*, 2008, **213**, 297.
7. A. Agrawal, P. P. Singh, B. Bottazzi, C. Garlanda, and A. Mantovani, *Adv. Exp. Med. Biol.*, 2009, **653**, 98.
8. M. E. Bianchi, *J. Leukoc. Biol.*, 2007, **81**, 1.
9. M. Yoneyama and T. Fujita, *J. Biol. Chem.*, 2007, **282**, 15315.
10. M. H. Shaw, T. Reimer, Y. G. Kim, and G. Nunez, *Curr. Opin. Immunol.*, 2008, **20**, 377.
11. T. Kawai and S. Akira, *Nat. Immunol.*, 2010, **11**, 373.
12. A. Iwasaki and R. Medzhitov, *Nat. Immunol.*, 2004, **5**, 987.
13. Y. Kumagai, O. Takeuchi, and S. Akira, *J. Infect. Chemother.*, 2008, **14**, 86.
14. Y. Kumagai, O. Takeuchi, and S. Akira, *J. Infect. Chemother.*, 2008, **14**, 86.
15. K. Takeda and S. Akira, *Curr. Protoc. Immunol.*, 2007, **Chapter 14**, Unit.
16. S. Akira, *Adv. Immunol.*, 1902, **78**, 1.
17. S. Akira, K. Takeda, and T. Kaisho, *Nature Immunol.*, 2001, **2**, 675.
18. A. Cottalorda, C. Verschelde, A. Marçais, M. Tomkowiak, P. Musette, S. Uematsu, S. Akira, J. Marvel, and N. Bonnefoy-Berard, *Eur. J. Immunol.*, 2006, **36**, 1684.
19. T. Kaisho and S. Akira, *Biochim. Biophys. Acta*, 2002, **1589**, 1.
20. G. R. Stark, I. M. Kerr, B. R. Williams, R. H. Silverman, and R. D. Schreiber, *Annu. Rev. Biochem.*, 1998, **67**, 227.
21. S. S. Diebold, T. Kaisho, H. Hemmi, S. Akira, and C. Reis e Sousa, *Science*, 2004, **303**, 1529.
22. A. D. Edwards, S. S. Diebold, E. M. Slack, H. Tomizawa, H. Hemmi, T. Kaisho, S. Akira, and Reis e Sousa, *Eur. J. Immunol.*, 2003, **33**, 827.
23. D. Rajagopal, C. Paturel, Y. Morel, S. Uematsu, S. Akira, and S. S. Diebold, *Blood*, 2010, **115**, 1949.
24. A. Krug, G. D. Luker, W. Barchet, D. A. Leib, S. Akira, and M. Colonna, *Blood*, 2004, **103**, 1433.
25. J. Lund, A. Sato, S. Akira, R. Medzhitov, and A. Iwasaki, *J. Exp. Med.*, 2003, **198**, 513.

26. A. ISAACS and J. Lindemann, *Proc. R. Soc. Lond B Biol. Sci.*, 1957, **147**, 258.
27. A. Le Bon, N. Etchart, C. Rossmann, M. Ashton, S. Hou, D. Gewert, P. Borrow, and D. F. Tough, *Nat. Immunol.*, 2003, **4**, 1009.
28. A. Le Bon, G. Schiavoni, G. D'Agostino, I. Gresser, F. Belardelli, and D. F. Tough, *Immunity.*, 2001, **14**, 461.
29. D. Durali, de Goer de Herve MG, J. Giron-Michel, B. Azzarone, J. F. Delfraissy, and Y. Taoufik, *Blood*, 2003, **102**, 4084.
30. F. E. Lund and T. D. Randall, *Nat. Rev. Immunol.*, 2010, **10**, 236.
31. de Goer de Herve MG, D. Durali, B. Dembele, M. Giuliani, T. A. Tran, B. Azzarone, P. Eid, M. Tardieu, J. F. Delfraissy, and Y. Taoufik, *PLoS ONE.*, 2011, **6**, e19366.
32. T. Sareneva, S. Matikainen, M. Kurimoto, and I. Julkunen, *J. Immunol.*, 1998, **160**, 3232.
33. J. D. Hood, H. J. Warshakoon, M. R. Kimbrell, N. M. Shukla, S. Malladi, X. Wang, and S. A. David, *Hum. Vaccin.*, 2010, **6**, 1.
34. N. M. Shukla, T. C. Lewis, T. P. Day, C. A. Mutz, R. Ukani, C. D. Hamilton, R. Balakrishna, and S. A. David, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 3232.
35. N. J. Horscroft, D. C. Pryde, and H. Bright, *J. Antimicrob. Chemother.*, 2012, **67**, 789.
36. S. J. Bergman, M. C. Ferguson, and C. Santanello, *Infect. Dis. Clin. North Am.*, 2011, **25**, 819.
37. B. J. Weigel, S. Cooley, T. Defor, D. J. Weisdorf, A. Panoskaltis-Mortari, W. Chen, B. R. Blazar, and J. S. Miller, *Am. J. Hematol.*, 2012.
38. J. F. Gerster, K. J. Lindstrom, R. L. Miller, M. A. Tomai, W. Birmachu, S. N. Bomersine, S. J. Gibson, L. M. Imbertson, J. R. Jacobson, R. T. Knafla, P. V. Maye, N. Nikolaidis, F. Y. Oneyemi, G. J. Parkhurst, S. E. Pecore, M. J. Reiter, L. S. Scribner, T. L. Testerman, N. J. Thompson, T. L. Wagner, C. E. Weeks, J. D. Andre, D. Lagain, Y. Bastard, and M. Lupu, *J. Med. Chem.*, 2005, **48**, 3481.
39. K. Hirota, K. Kazaoka, I. Niimoto, H. Kumihara, H. Sajiki, Y. Isobe, H. Takaku, M. Tobe, H. Ogita, T. Ogino, S. Ichii, A. Kurimoto, and H. Kawakami, *J. Med. Chem.*, 2002, **45**, 5419.
40. Y. Isobe, M. Tobe, H. Ogita, A. Kurimoto, T. Ogino, H. Kawakami, H. Takaku, H. Sajiki, K. Hirota, and H. Hayashi, *Bioorg. Med. Chem.*, 2003, **11**, 3641.
41. A. Kurimoto, T. Ogino, S. Ichii, Y. Isobe, M. Tobe, H. Ogita, H. Takaku, H. Sajiki, K. Hirota, and H. Kawakami, *Bioorg. Med. Chem.*, 2004, **12**, 1091.
42. A. Kurimoto, K. Hashimoto, T. Nakamura, K. Norimura, H. Ogita, H. Takaku, R. Bonnert, T. McNally, H. Wada, and Y. Isobe, *J. Med. Chem.*, 2010, **53**, 2964.
43. J. Lee, T. H. Chuang, V. Redecke, L. She, P. M. Pitha, D. A. Carson, E. Raz, and H. B. Cottam, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 6646.
44. A. X. Xiang, S. E. Webber, B. M. Kerr, E. J. Rueden, J. R. Lennox, G. J. Haley, T. Wang, J. S. Ng, M. R. Herbert, D. L. Clark, V. N. Banh, W. Li, S. P. Fletcher, K. R. Steffy, D. M. Bartkowski, L. I. Kirkovsky, L. A. Bauman, and D. R. Averett, *Nucleosides Nucleotides Nucleic Acids*, 2007, **26**, 635.
45. N. M. Shukla, M. R. Kimbrell, S. S. Malladi, and S. A. David, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 2211.
46. N. M. Shukla, S. S. Malladi, C. A. Mutz, R. Balakrishna, and S. A. David, *J. Med. Chem.*, 2010, **53**, 4450.
47. N. M. Shukla, S. S. Malladi, V. Day, and S. A. David, *Bioorg. Med. Chem.*, 2011, **19**, 3801.
48. N. M. Shukla, C. A. Mutz, R. Ukani, H. J. Warshakoon, D. S. Moore, and S. A. David, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 6384.
49. N. M. Shukla, C. A. Mutz, S. S. Malladi, H. J. Warshakoon, R. Balakrishna, and S. A. David, *J. Med. Chem.*, 2012, **55**, 1106.
50. E. Brummer, M. A. Antonysamy, L. Bythadka, G. W. Gullikson, and D. A. Stevens, *FEMS Immunol. Med. Microbiol.*, 2010, **59**, 81.
51. Lindstrom, K. J. and Nikolaidis, N. Imidazo[4,5-c]pyridin-4-amines. World patent. [WO 95/02597]. 1995.
52. Kshirsagar, T. A., Merrill, B. A., Langer, S. E., Lindstrom, K. J., Johannessen, S. C., Marszalek, G. J., Wurst, J. R., Manske, K. J., Niwas, S., Lundquist, G. D. Jr., Heppner, P. D., Griesgraber, G. W., and Danielson, M. E. Method of preferentially inducing the biosynthesis of interferon. World patent. WO 2006/091467 A2[1], 252. 2006.
53. Kshirsagar, T. A., Lundquist, G. D. Jr., Dellaria, J. F. Jr., Radmer, M. R., and Zimmermann, B. M. Oxime and hydroxylamine substituted imidazo[4,5-c] ring compounds and methods. World patent. [WO 2006/086634]. 2006.
54. Dellaria, J. F., Haraldson, G. A., Heppner, P. D., Lindstrom, K. J., and Merrill, B. A. Sulfonamido substituted imidazopyridines. [US 6,525,064 B1], 1-35. 2003.
55. J. D. Hood, H. J. Warshakoon, M. R. Kimbrell, N. M. Shukla, S. Malladi, X. Wang, and S. A. David, *Hum. Vaccin.*, 2010, **6**, 1.
56. A. T. Baviskar, C. Madaan, R. Preet, P. Mohapatra, V. Jain, A. Agarwal, S. K. Guchhait, C. N. Kundu, U. C. Banerjee, and P. V. Bharatam, *J. Med. Chem.*, 2011, **54**, 5013.
57. M. S. Jin and J. O. Lee, *Curr. Opin. Immunol.*, 2008, **20**, 414.
58. S. I. Yoon, O. Kurmasov, V. Natarajan, M. Hong, A. V. Gudkov, A. L. Osterman, and I. A. Wilson, *Science*, 2012, **335**, 859.
59. H. Hemmi, T. Kaisho, O. Takeuchi, S. Sato, H. Sanjo, K. Hoshino, T. Horiuchi, H. Tomizawa, K. Takeda, and S. Akira, *Nat. Immunol.*, 2002, **3**, 196.
60. T. Kaisho and S. Akira, *J. Allergy Clin. Immunol.*, 2006, **117**, 979.
61. K. B. Gorden, K. S. Gorski, S. J. Gibson, R. M. Kedl, W. C. Kieper, X. Qiu, M. A. Tomai, S. S. Alkan, and J. P. Vasilakos, *J. Immunol.*, 2005, **174**, 1259.
62. K. S. Gorski, E. L. Waller, J. Bjornton-Severson, J. A. Hanten, C. L. Riter, W. C. Kieper, K. B. Gorden, J. S. Miller, J. P. Vasilakos, M. A. Tomai, and S. S. Alkan, *Int. Immunol.*, 2006, **18**, 1115.
63. H. P. Kokatla, E. Yoo, D. B. Salunke, D. Sil, C. F. Ng, R. Balakrishna, S. S. Malladi, L. M. Fox, and S. A. David, *Org. Biomol. Chem.*, 2013, **11**, 1179.
64. S. A. David, M. S. Smith, G. Lopez, S. Mukherjee, S. Buch, and O. Narayan, *AIDS Res. Human Retrovir.*, 2001, **17**, 59.
65. M. R. Kimbrell, H. Warshakoon, J. R. Cromer, S. Malladi, J. D. Hood, R. Balakrishna, T. A. Scholdberg, and S. A. David, *Immunol. Lett.*, 2008, **118**, 132.