
This article was downloaded by: [University of Western Ontario]

On: 07 October 2014, At: 02:39 Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered

office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/Incn19

Synthesis and Biological Activity of 5-Azacytosine Nucleosides Derived from 4-Thio-2-Deoxy-L-threo-Pentofuranose and 4-Thio-2-Deoxy-D-erythro-Pentofuranose

Loredana Cappellacci $^{\rm a}$, Kamal N. Tiwari $^{\rm a}$, John A. Montgomery $^{\rm a}$ & John A. Secrist III $^{\rm a}$

^a Southern Research Institute, P.O. Box 55305, Birmingham, AL, 35255-5305, USA

Published online: 04 Oct 2006.

To cite this article: Loredana Cappellacci, Kamal N. Tiwari, John A. Montgomery & John A. Secrist III (1999) Synthesis and Biological Activity of 5-Azacytosine Nucleosides Derived from 4-Thio-2-Deoxy-L-threo-Pentofuranose and 4-Thio-2-Deoxy-D-erythro-Pentofuranose, Nucleosides and Nucleotides, 18:4-5, 613-614, DOI: 10.1080/15257779908041514

To link to this article: http://dx.doi.org/10.1080/15257779908041514

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

SYNTHESIS AND BIOLOGICAL ACTIVITY OF 5-AZACYTOSINE NUCLEOSIDES DERIVED FROM 4-THIO-2-DEOXY-L-threo-PENTOFURANOSE AND 4-THIO-2-DEOXY-D-erythro-PENTOFURANOSE

Loredana Cappellacci, Kamal N. Tiwari, John A. Montgomery, and John A. Secrist III Southern Research Institute, P.O. Box 55305, Birmingham, AL 35255-5305, USA

ABSTRACT: 1-O-Acetyl-2-deoxy-3,5-di-O-toluoyl-4-thio-D-*erythro*-pentofuranose and 2-deoxy-1,3,5-tri-O-acetyl-4-thio-L-*threo*-pentofuranose were coupled with 5-azacytosine to obtain α and β anomers of nucleosides.

For some years we have been pursuing the synthesis of a variety of 4'-thionucleosides as anticancer and anti-infective agents. In that regard we have prepared 4'-thionucleosides with a variety of different carbohydrates attached to both the normal purine and pyrimidine bases as well as analogs of those bases. One of our goals in this program is to gain a further understanding about the active site of deoxycytidine kinase, so that we are better able to prepare potential drugs that are activated by this enzyme.

As part of this program, we have prepared a series of 4'-thionucleosides incorporating both 5-azacytosine and 5,6-dihydro-5-azacytosine, in order to assess their anticancer activity, as well as to compare the saturated and unsaturated compounds in the same system. Specifically, we have prepared the α and β anomers of the nucleosides incorporating 5-azacytosine and 5,6-dihydro-5-azacytosine into 4-thio-2-deoxy-L-threo-pentofuranose and 4-thio-2-deoxy-D-erythro-pentofuranose. Chemical syntheses and biological data on these four compounds is presented.

The synthesis of 1-O-acetyl-2-deoxy-4-thio-3,5-di-O-p-toluoyl-D-erythro-pentofuranose 1 and 2-deoxy-1,3,5-O-acetyl-4-thio-L-threo-pentofuranose 6, as 1:1 mixture of anomers have been performed as previously reported.

Trimethylsilyl triflate catalyzed coupling of thiosugar 1 and 6 with silylated 5-azacytosine afforded the corresponding nucleosides (2 and 7) as anomeric mixtures (β : α

ratio $\sim 1:1.8$). Silica gel chromatography and fractional crystallization of 2 and 7 afforded pure anomers.

Reduction with sodium borohydride of compounds 2α , 2β , 7α , 7β , afforded 4α , 4β , 9α , 9β , respectively. Deprotection of compounds 2α , 2β , 4α , 4β , 7α , 7β , 9α , 9β , with sodium methoxide afforded 3α , 3β , 5α , 5β , 8α , 8β , 10α , 10β , respectively.

The assignment of the anomeric configurations of compounds were made by NOE difference spectroscopy. Only 2'-deoxy-4'-thio-5-azacytosine (3β) showed significant toxicity, as shown in Table 1. All the 2'-deoxy-4'-thiopyrimidine nucleoside analogues were evaluated against Hepatitis B virus (HBV) in vitro and no significant antiviral activity was observed.

Table 1. Cytotoxicity Data: IC50 (μM)					
Compound	CCRF-CEM (leukemia)	CAKI-1 (renal)	DLD-1 (colon)	NCI-H23 (lung)	SNB-7 (CNS)
3β	0.001	0.4	7	4	9

REFERENCES

- Tiwari, K. N.; Montgomery, J. A.; Secrist III, J. A. The Synthesis and Biological Activity of 1-(2-deoxy-4-thio-α-L-Threo-Pentofuranosyl)Thymine. Nucleosides Nucleotides, 1993, 12(8), 841-846; Secrist III, J. A., Tiwari, K. N., Riordan, J. M., Montgomery, J. A. Synthesis and Biological Activity of 2'-Deoxy-4'-thio Pyrimidine Nucleosides. J. Med. Chem. 1991, 34, 2361-2366.
- Secrist III, J. A.; Tiwari, K. N.; Shortnacy-Fowler, A. T.; Messini, L.; Riordan, J. M.; Montgomery, J. A. Synthesis and Biologic Activity of Certain 4'-Thio-Darabinofuranosyl Purine Nucleosides. J. Med. Chem. 1998, 41, 3865-3871.