Xenon difluoride-trimethylsilyl isocyanate-triflic acid as a new system for the amination of aromatic compounds

Namig Sh. Pirkuliev,^{a,b} Valery K. Brel,^{*a} Novruz G. Akhmedov,^b Nikolai S. Zefirov^{a,b} and Peter J. Stang^c

^a Institute of Physiologically Active Compounds, Russian Academy of Sciences, Chernogolovka, 142432 Moscow region, Russian Federation. Fax: +7 095 913 2113; e-mail: brel@ipac.ac.ru

^b Department of Chemistry, M. V. Lomonosov Moscow State University, 119899 Moscow, Russian Federation

^c Department of Chemistry, Unversity of Utah, Salt Lake City, UT 84112, USA

10.1070/MC2001v011n05ABEH001514

In the title system, $OCNXeOSO_2CF_3$ is formed, which readily oxidises iodobenzene to [PhI+-NCO -OTf]. The direct amination of aromatic substrates is possible with the use of XeF_2 -Me₃SiNCO-CF₃SO₃H.

Since the discovery of the first xenon compounds by Bartlett¹ in 1962, a large number of xenon derivatives with Xe–F, Xe–Cl, Xe–O, Xe–N, Xe–C and Xe–N bonds were prepared.² The methodology of Xe–element bond formation is based on the rupture of M–element bonds (where M = Bi, Sb or Si) with the aid of xenon fluorides. The Si–element fission bond is most promising and interesting for obtaining compounds with Xe–N bonds.^{2(a),3} Thus, D. D. DesMarteau^{3(c)} used this approach for the preparation of the relatively stable compound FXe(NSO₂F)₂. Recently, theoretical computations and product analysis of the reaction of XeF₂ with NaN₃ and NaOCN indicated the intermediate formation of FXeN₃ and FXeNCO.⁴

Previously, we found that the reaction of trimethylsilyl isocyanate with XeF₂ or FXeOSO₂CF₃ in the presence of olefins proceeds through the formation of FXeNCO and OCNXeOSO₂CF₃ intermediates, which are easily added at the double bond of an olefin. In a continuation of this work, we studied the interaction of trimethylsilyl isocyanate with XeF₂ or FXeOSO₂CF₃ in the presence of aromatic compounds. It is well known that, in this case, the reaction path strongly depends on the order in which the reactants were added. In particular, if finely dispersed XeF₂ in CH₂Cl₂ was transformed into FXeOTf⁵ and the latter was treated with trimethylsilyl isocyanate and then with iodobenzene, mixed iodonium sulfonate 3^6 was obtained.[†]

$$\begin{array}{ccc} XeF_2 + HOTf & \xrightarrow{CH_2Cl_2} & FXeOTf & \xrightarrow{Me_3SINCO} \\ & & & & \\ I & & \\ [Xe(OTf)(NCO)] & \xrightarrow{PhI} & TfO^- PhI^+-NCO \\ & & & \\ 2 & & & \\ Scheme & 1 & \end{array}$$

When trimethylsilyl isocyanate was added to a XeF_2 solution in CH_2Cl_2 and the resulting mixture was subsequently treated with triflic acid and iodobenzene, phenyl(*p*-iodophenyl)iodonium triflate **4** and iodoanilines **5**, **6** (as a mixture of *ortho* and *para* isomers) were isolated from the reaction mixture as a final products.[‡] Note that in this case the formation of compound **3** was not detected.

Consequently, the XeF_2 -Me₃SiNCO-CF₃SO₃H system acts as an aminating reagent for aromatic compounds.^{7,8}

Moreover, we found that with the use of other aromatic compounds such as benzene, toluene, chlorobenzene and *o*-xylene

[Isocyanato(trif)loxy)- λ^3 -iodo]benzene **3**. ¹H NMR (CD₃CN) δ : 8.3–7.5 (m). ¹³C NMR (CD₃CN) δ : 172.5 (CO), 136.6, 135.0, 132.9, 122.7 (Ph), 120.8. ¹⁹F NMR (CD₃CN) δ : –78.6 (CF₃). IR (CCl₄, ν /cm⁻¹): 2510, 1625 (NCO), 1258, 1178, 1023 (OTf).



instead of iodobenzene the formation of amines 7 and 8 takes place (Table 1).

Table 1 Amination of aromatic compounds with XeF2-Me3SiNCO-CF3SO3H.

Substrate	Reaction time/h	Yield ^a (%)	<i>ortho</i> -isomer	<i>para-</i> isomer	Yield ^{<i>a</i>} of 4 (%)
C ₆ H ₆	3	45	_		
MeC ₆ H ₅	3	40	54	46	
IC ₆ H ₅	4	38	50	44	44
ClC ₆ H ₅	6	35	45	48	
o-Xylene	3	43	_	_	
"Icolated wields based on VoE wood					

^aIsolated yields based on XeF₂ used.

* Reaction of iodobenzene with XeF₂-Me₃SiNCO-HOTf. XeF₂ (4.72 mmol) was dissolved in dry CH₂Cl₂ (15 ml)¹⁰ under argon; then, Me₃SiNCO (5.1 mmol) was added. Triflic acid (10 mmol) was slowly added dropwise at -78 °C to the solution of FXeNCO in dry CH2Cl2 and then the mixture was stirred at -30 °C for ~1 h. Then, a large excess of PhI (12 mmol) was added to the suspension at -78 °C, and the mixture was stirred until no more gas (Xe) was evolved (see Table 1 for total reaction time). The solution was heated to room temperature and poured into a dilute solution of HCl with ice. An excess of iodobenzene and phenyl(p-iodophenyl)iodonium triflate 4 were extracted with dichloromethane $(3 \times 15 \text{ ml})$. The volatile materials were then removed under reduced pressure to give an oily residue, which was dissolved in dry diethyl ether. The mixture was vigorously shaken for several minutes to precipitate slightly coloured crystals. Analytically pure samples were obtained by recrystallization from CH₂Cl₂-Et₂O. The aqueous layer was neutralised with a 30% sodium hydroxide solution to pH ~13. The mixture of o- and p-iodoanilines 5, 6 was then extracted with dichloromethane and dried with MgSO4. Products were isolated after evaporating the solvent. Isomeric mixtures were analysed by 1H NMR.

Phenyl(p-*iodophenyl*)*iodonium triflate* **4**: yield 44%, mp 146–147 °C (lit.,¹¹ 144–148 °C). ¹H NMR (CDCl₃) δ: 7.48 (m, 2H, Ph), 7.64 (m, 1H, Ph), 7.80 (m, 4H, C₆H₄), 8.10 (m, 2H, Ph). ¹⁹F NMR (CDCl₃) δ: -78.4 (CF₃SO₃).

[†] Typical procedure for the preparation of [PhI⁺NCO -OTf]. 5.52 mmol of Me₃SiNCO was added to a stirred suspension of 4.72 mmol of CF₃SO₂OXeF⁵ in 20 ml of CH₂Cl₂ at -78 °C under argon. The mixture was allowed to warm up to -40 °C and stirred until the formation of a yellow homogeneous solution. The solution was cooled to -78 °C; then, 4.72 mmol of iodobenzene was added using a syringe. The mixture was allowed to warm to -5 °C and stirred for ~2 h. The precipitate was filtered off under argon, washed with cold diethyl ether and dried *in vacuo* to yield the material of >98% purity. The analytically pure compound can be obtained by recrystallization from CH₂Cl₂–Et₂O.

These results indicate that the $XeF_2-Me_3SiNCO-CF_3SO_3H$ system makes it possible to perform conveniently electrophilic one-step amination of aromatic compounds with the formation of anilines in moderate yields. Highly deactivated aromatics slowly react under these conditions to form anilines in low yields. Thus, with nitrobenzene, the yields of nitroanilines were lower than 5%. In this case, the deactivation of a benzene ring probably occurs not only owing to the nitro group but also due to the further deactivation of the ring by the protonation of the nitro group in a triflic acid medium.⁹

It is believed that the amination of aromatic compounds[§] involves the formation of the reactive triflate $[H_2N^+=C=O -OTf]$ at the stage of the interaction of FXeNCO with triflic acid. This explanation was additionally supported by studies of the reactions with Me₃SiOTf in place of HOTf. In this case, the $[H_2N^+=C=O -OTf]$ species was not formed, apparently, due to the absence of hydrogen ions. Most likely, the formation of OCNXeOSO₂CF₃ occurs, and hence the reaction with iodobenzene leads to the formation of only iodonium sulfonate **3** (Scheme 1).

Thus, we found that in the case of the $XeF_2-CF_3SO_3H-Me_3SiNCO$ system the formation of OCNXeOSO₂CF₃ takes place, which readily oxidises iodobenzene to [PhI⁺-NCO ⁻OTf]. The direct amination of aromatic substrates with reasonable yields is possible with the use of $XeF_2-Me_3SiNCO-CF_3SO_3H$.

We are grateful to NIH and FIRCA (2RO3TW000437) for financial support.

References

- 1 N. Bartlett, Proc. Chem. Soc., 1962, 218.
- 2 (a) V. K. Brel, N. Sh. Pirkuliev and N. S. Zefirov, Usp. Khim., 2001, 70, 262 (Russ. Chem. Rev., 2001, 70, 231); (b) New Fluorinating Agents in Organic Synthesis, eds. L. German and S. Zemskov, Springer–Verlag, Berlin, 1989.
- 3 (a) D. D. DesMarteau, J. Am. Chem. Soc., 1978, 100, 6270; (b) D. D. DesMarteau, R. D. LeBlond, S. F. Hossain and D. Nothe, J. Am. Chem. Soc., 1981, 103, 7734; (c) J. F. Sawyer, G. J. Schrobilgen and S. J. Sutherland, J. Chem. Soc., Chem. Commun., 1982, 210; (d) A. A. A. Emara and G. J. Schrobilgen, Inorg. Chem., 1992, 31, 1323.
- 4 A. Schulz and T. M. Klapotke, Inorg. Chem., 1997, 36, 1929.
- 5 (a) T. M. Kasumov, N. Sh. Pirguliyev, V. K. Brel, Y. K. Grishin, N. S. Zefirov and P. J. Stang, *Tetrahedron*, 1997, **53**, 13139; (b) M. Wechsberg, P. A. Bulliner, F. O. Sladky, R. Mews and N. Bartlett, *Inorg. Chem.*, 1972, **11**, 3063.
- 6 V. V. Zhdankin, C. M. Critell, P. J. Stang and N. S. Zefirov, *Tetrahedron Lett.*, 1990, **31**, 4821.
- 7 (a) R. Schroter and E. Muller, Houben–Weyl, Methoden der Organischen Chemie, Thieme Verlag, Stuttgart, 1957, vol. XI/1, pp. 341–488; (b)
 P. Kovacic, Friedel–Crafts and Related Reactions, Interscience Publishers, New York, 1964, vol. II/2, p. 1493; (c) T. Sheradsky, The Chemistry of Amino, Nitroso and Nitro Compounds and Their Derivatives, John Wiley, New York, 1982, part 1, pp. 395–416.
- 8 (a) C. Graebe, *Ber. Dtsch. Chem. Ges.*, 1901, 34, 1778; (b) G. F. C. Jaubert and R. Hebd, *Seances Acad. Sci.*, 1901, 132, 841; (c) P. Kovacic, R. L. Russel and R. B. Bennett, *J. Am. Chem. Soc.*, 1964, 86, 1588; (d) P. Kovacic and R. B. Bennett, *J. Am. Chem. Soc.*, 1961, 83, 221; (e) G. A. Olah and T. D. Ernst, *J. Org. Chem.*, 1989, 54, 1203.
- 9 G. A. Olah, G. K. S. Prakash and J. Sommer, *Superacids*, John Wiley, New York, 1985.
- 10 D. D. Perrin, W. L. F. Armaredo and D. R. Perrin, *Purification of Laboratory Chemicals* (2nd edition), Pergamon Press, New York, 1980.
- 11 T. Kitamura, J.-I. Matsuyuki, K. Nagata, R. Furuki and H. Taniguchi, *Synthesis*, 1992, 945.

Received: 10th August 2001; Com. 01/1840

[§] General procedure for the amination of aromatic compounds. The reactions with other aromatics were carried out as described above for the interaction of iodobenzene with XeF_2 -Me₃SiNCO-HOTf system. Isomeric arylamines 7 and 8 were isolated in moderate yields (Table 1). Physico-chemical constants and spectral properties of 7 and 8 are in agreement with the published data.