

View Article Online View Journal

RSC Advances

This article can be cited before page numbers have been issued, to do this please use: F. Yang, H. Wang, L. Jiang, H. Yue, H. Zhang, Z. Wang and L. Wang, *RSC Adv.*, 2014, DOI: 10.1039/C4RA13272F.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This Accepted Manuscript will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/advances



Lipase-catalyzed synthesis of benzo[g]chromene derivatives

Cite this: DOI: 10.1039/c0xx00000x

Published on 11 December 2014. Downloaded by Cornell University Library on 15/12/2014 09:14:49.

ARTICLE TYPE

A green and one-pot synthesis of benzo[g]chromene derivatives through a multi-component reaction catalyzed by lipase

Fengjuan Yang,^{*a,b*} Haoran Wang,^{*a,b*} Liyan Jiang,^{*a,b*} Hong Yue,^{*a,b*} Hong Zhang,^{*a,c*} Zhi Wang^{**a,b*} and Lei Wang^{**a,b*}

s Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX DOI: 10.1039/b000000x

The synthesis of benzo[g]chromene derivatives through a multi-component reaction catalyzed by lipase was reported in the first time. This novel efficient method has the advantages of environmental friendliness, high yield and simple work-up. Moreover, this protocol extends the phenomenon of enzyme promiscuity.

A multi-component reaction is generally defined as a reaction in which three or more reactants combine in one pot to form a ¹⁵ single product that contains essentially all of the atoms of the starting materials (with the exception of condensation products, such as H₂O, HCl, or MeOH) [1-2]. Compared with the conventional chemical reactions, the multi-component reaction strategies can offer the advantage of simplicity and synthetic ²⁰ efficiency [3-4].

Enzyme catalytic promiscuity is the ability of an enzyme to catalyze more than one type of chemical transformation. As a "hidden skill" of enzyme, it can provide novel synthesis pathway that are currently not available [5,6] and can widen the ²⁵ application of enzyme. In this area, lipase is the most used enzyme due to its broad specificity and excellent stability in various media. Many elegant works of lipase catalytic promiscuity in the carbon–carbon bond-forming reactions (Aldol condensation, Morita–Baylis–Hillman reaction, Michael addition, ³⁰ Markovnikov addition, and Knoevenagel reaction, *et al.*) have

- been reported in the past few years [7-14]. As a part of our interest to explore the applications of lipase in this new area, we are focusing on the multi-component reaction of synthesis of benzo[g]chromene derivatives catalyzed by lipase in this study.
- ³⁵ Benzo[g]chromene derivatives have extensive bioactivities, such as antibacterial, antiproliferation and antitumor activities [15-20]. To the best of our knowledge, it is reported for the first time that the synthesis of benzo[g]chromene derivatives can be catalyzed by lipase (Scheme 1).



Scheme 1 Lipase-catalyzed synthesis of benzo[g]chromene derivatives

Initial studies were undertaken using benzaldehyde, malononitrile and 2-hydroxy-1,4-naphthoquinone as a model ⁴⁵ reaction. Several kinds of lipases were selected to catalyze this multi-component reaction and the results are listed in Table 1. It could be found that all the selected lipases can catalyze this multi-component reaction and the catalytic activities depend mainly on the lipase origin. When the denatured lipase or bovine ⁵⁰ serum albumin (BSA) was used as the catalyst, almost no product could be detected which suggested a special active conformation of enzyme play a crucial role in this multi-component reaction. Among of the selected lipases, *Candida sp.* lipase (CSL) exhibited the highest catalytic activity. Therefore, we chose CSL ⁵⁵ as the catalyst for the multi-component reaction. ‡

Table 1 The catalytic activities of different lipases in the synthesis of benzo[g]chromene derivatives ^a

Enter	Enguina	Isolated yield (9/)
Linuy		
1	Candida antarctica Lipase B	57
	(CALB)	
2	Porcine pancreas Lipase	79
	(PPL)	
3	Candida sp. Lipase	88
	(CSL)	
4	Pseudomonas fluorescens Lipase	54
	(PFL)	
5	Pseudomonas sp. Lipase	66
	(PSL)	
6	Bacillus subtillus Lipase	70
	(BSL2)	
7	C. rugosa Lipase	61
	(CRL)	
8	Bovine serum albumin	Trace
	(BSA)	
9	Candida sp. Lipase	Trace
-	(denatured) ^b	
10	No enzyme	Trace
	1.0 enzyme	11400

^a Reaction condition: 2-hydroxy-1,4-naphthoquinone (1 mmol), malononitrile (1 mmol) and benzaldehyde (1 mmol), ethanol (2mL),
⁶⁰ enzyme (20 mg, protein content), 55 °C, 12h. ^b CSL was denatured by heating it to 100 °C for 6 h in water before lyophilization.

Choosing a suitable solvent is of crucial importance for the enzyme catalytic performance [21]. Thus, seven organic solvents ⁶⁵ were screened for this reaction and the results are presented in Figure 1. Compared with other solvents, the highest yield (88%) could be obtained while ethanol was used as the reaction media. It's believed that the solvent can lead to a conformational change Published on 11 December 2014. Downloaded by Cornell University Library on 15/12/2014 09:14:49

26].

of the enzyme and then affect the enzyme activity [22].



Fig. 1 Solvent effect on the synthesis of benzo[g]chromene derivatives catalyzed by lipase

- 5 Reaction condition: 2-hydroxy-1,4-naphthoguinone (1 mmol). malononitrile (1 mmol) and benzaldehyde (1 mmol), solvent (2mL), CSL (20 mg, protein content), 55 °C, 12h.
- It's generally believed that reaction temperature is another 10 vital influence factor for all enzymatic reactions [23]. In this study, the reaction temperature was varied from 25 to 75 °C to investigate its effect. As shown in Figure 2, the yield increased as temperature was enhanced from 25 to 55 °C and dropped dramatically at higher temperature. The increased collision 15 chance between the enzyme and substrate at the elevated temperature may improve the reaction rate. Further increasing temperature may disrupt the enzyme conformation and decrease the enzyme activity. Since the yield was found to be the highest at 55 °C, the optimum temperature for this reaction was 55 °C.



Fig. 2 Temperature effect on the synthesis of benzo[g]chromene derivatives catalyzed by lipase

2-hydroxy-1,4-naphthoquinone (1 mmol), Reaction condition: malononitrile (1 mmol) and benzaldehyde (1 mmol), ethanol (2mL), CSL 25 (20 mg, protein content), 12h at different temperature.

To explore the scope and feasibility of this multi-component reaction, a series of benzo[g]chromene derivatives were synthesized under the optimum reaction conditions. The results in Table 2 demonstrated that the protocol could be applied to aromatic aldehydes either with electron-30 withdrawing groups (Entry 2-6) or electron-donating groups (Entry 7-9) with satisfied yields (from 81% to 93%). It is noteworthy that the electronic nature of substituents of the aromatic aldehyde has no distinct effect on the multi-component reaction.

The lipase-catalyzed Knoevenagel condensation and Michael addition 35 have been reported previously [10, 24]. According to these reports and our results in this study, a plausible mechanism was proposed in Scheme 2. The synthesis of benzo[g]chromene derivatives involves a Knoevenagel condensation, Michael addition, cyclization and isomerization, respectively. As shown in Scheme 2, lipase could catalyze 40 the steps of Knoevenagel condensation and Michael addition during the reaction process. Firstly, enzymatic Knoevenagel condensation of the aldehyde 1 to malononitrile 2 was occurred to produce an intermediate 5. Secondly, Michael addition of the 2-Hydroxy-1,4-naphthoquinone 3 on the intermediate 5 could be catalyzed by lipase to produce an intermediate 45 6. Finally, intramolecular cyclization and isomerization formed the product 4 automatically. It's important to note that the benzo[g]chromene derivatives obtained in this study were racemic when all the screened lipases were used as catalyst, which indicated that lipase didn't exhibit the stereoselectivity in the Michael addition of this reaction. This 50 phenomenon was in accordance with the recent reported literatures [25,



Scheme 2 Mechanism of the lipase-catalyzed synthesis of benzo[g]chromene derivatives

Table 2 Lipase-catalyzed synthesis of benzo[g]chromene derivatives with different aromatic aldehydes $^{\rm a}$



^a Reaction condition: 2-Hydroxy-1,4-naphthoquinone (1 mmol), malononitrile (1 mmol) and aromatic aldehyde (1 mmol), ethanol (2mL),
⁵ CSL (20 mg, protein content), 55 °C, 12h.

Conclusions

Published on 11 December 2014. Downloaded by Cornell University Library on 15/12/2014 09:14:49

In summary, an efficient and simple method for the synthesis of benzo[g]chromene derivatives catalyzed by lipase was reported for the first time. After thorough optimization of reaction 10 conditions, all the products could be obtained in high yields (from 81% to 93%). Compared with the reported methods [27-33], the notable features of this new synthetic route are not only atom economy, environmental friendliness and simple operational process, but more importantly, this work significantly expands 15 the utility of lipase in organic synthesis and encourages us to use

- the current tools of enzyme engineering and directed evolution to increase the catalytic performance of lipase. It's known that immobilization is a powerful tool to avoid the enzyme aggregation in organic solvent and recover and reuse of the
- ²⁰ enzyme with high remnant activity [34-37]. Further study of the immobilization enzyme on the lipase-catalyzed synthesis of benzo[g]chromene derivatives is now in progress in our laboratory.

25 Acknowledgements

We gratefully acknowledge the National Natural Science Foundation of China (No. 21172093 and 31070708), the Natural Science Foundation of Jilin Province of China (No. 20140101141JC) and the Scientific Research Fund of Jilin

³⁰ University (No. 450060326007 and 450060491559) for the financial support.

ARTICLE TYPE

Notes and references

- ³⁵ ^a Key Laboratory of Molecular Enzymology and Engineering of Ministry of Education, Jilin University, Changchun 130023, P R China; Email: wangzhi@jlu.edu.cn, w lei@jlu.edu.cn
 - ^b School of life sciences, Jilin University, Changchun 130023, P R China;
 - ^c College of Chemistry, Jilin University, Changchun 130023, P R China;
- ⁴⁰ † Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/
- ‡ A typical enzymatic procedure of the reaction: CSL (20 mg, protein 45 content) was added to a 25 mL round-bottom flask containing aromatic
- 45 content) was added to a 25 mil found-bottom mask containing atomatic aldehyde (1 mmol), malononitrile (1 mmol) and ethanol (2 mL). The suspension was maintained at 55 °C for 10 min. Then, 2-hydroxy-1,4naphthoquinone (1 mmol) was added to the reaction mixture. After completion of the reaction (after 12 h, monitored by TLC), the reaction
- 50 mixture was concentrated under vacuum. The residue was washed with water and cold diethyl ether three times to remove unreacted starting materials and other organic contaminations, and then the filter cake was recrystallized from 95% ethanol to give products 4 with high purity. The experiments were performed triplicate, and all data were obtained based 55 on the average values. The products were characterized by NMR and ESI-MS experiments.
- 1 E. Ruijter, R. Scheffelaar and R. V. A. Orru, *Angew. Chem. Int. Ed.*, 2011, **50**, 6234.
- 60 2 B. Ganem, Accounts Chem. Res., 2009, 42, 463.
- 3 H. Bienaymé, C. Hulme, G. Oddon and P. Schmitt, *Chem. Eur. J.*, 2000, 6, 3321.
- 4 R. M. Armstrong, A. P. Combs, P. A. Tempest, S. D. Brown and T. A. Keating, Acc. Chem. Res., 1996, 29, 123.
- 65 5 I. Nobeli, A. D. Favia and J. M. Thornton, Nat. Biotechnol., 2009, 27, 157.
- 6 L. L. Torres, A. Schließmann, M. Schmidt, N. Silva-Martin, J. A. Hermoso, J. Berenguer, U. T. Bornscheuer and A. Hidalgo, *Org. Biomol. Chem.*, 2012, **10**, 3388.
- 70 7 C. Branneby, P. Carlqvist, A. Magnusson, K. Hult, T. Brinck and P. Berglund, J. Am. Chem. Soc., 2003, 125, 874.
- 8 M. Svedendahl, K. Hult and P. Berglund, J. Am. Chem. Soc., 2005, 127, 17988.
- 9 M. T. Reetz, R. Mondiere and J. D. Carballeira, Tetrahedron Lett., 2007, 48, 1679.
- 10 W. B. Wu, N. Wang, J. M. Xu, Q. Wu and X. F. Lin, Chem. Commun., 2005, 2348.
- 11 Y. F. Lai, H. Zheng, S. J. Chai, P. F. Zhang and X. Z. Chen, *Green Chem.*, 2010, **12**, 1917.
- 80 12 H. R. Wang, Z. Wang, C. Y. Wang, F. J. Yang, H. Zhang, H. Yue and L. Wang, *RSC adv.*, 2014, **4**, 35686.
 - 13 Z. G. Le, L. T. Guo, G. F. Jiang, X. B. Yang and H. Q. Liu, *Green Chem. Lett. Rev.*, 2013, **6**, 277.
- 14 F. J. Yang, Z. Wang, H. R. Wang, H. Zhang, H. Yue and L. Wang, *RSC Adv.*, 2014, 4, 25633.
- 15 A. M. El-Agrody, M. H. El-Hakim, M. S. A. Abd El-Latif, A. H. Fakery, M. El-Sayed and K. A. El-Ghareab, *Acta Pharm.*, 2000, 50, 111.
- 16 J. Zamocka, E. Misikova and J. Durinda, Cesk Farm, 1992, 41, 170.
- 90 17 S. J. Mohr, M. A. Chirigos, F. S. Fuhrman and J. W. Pryor, *Cancer Res.*, 1975, **35**, 3750.

Published on 11 December 2014. Downloaded by Cornell University Library on 15/12/2014 09:14:49.

- 18 M. Brunavs, C. P. Dell and W. M. Owton, *J. Fluor. Chem.*, 1994, **68**, 201.
- 19 V. K. Tandon, M. Vaish, S. Jain, D. S. Bhakuni and R. C. Srimal, *Indian J. Pharm. Sci.*, 1991, **53**, 22.
- 5 20 P. Coudert, J. M. Coyquelet, J. Bastide, Y. Marion and J. Fialip, Ann. Pharm. Fr., 1988, 46, 91.
- 21Y. K. Huang, S. W. Tsai, Appl. Microbiol. Biotechnol., 2014, 98, 621.
- 22 Y. S. Lin, P. Y. Wang, A. C. Wu and S. W. Tsai, J. Mol. Catal. B-Enzym., 2011, 68, 245.
- 10 23 R. Tian, C. H. Yang, X. F. Wei, E. N. Xun, R. Wang, S. G. Cao, Z. Wang and L. Wang, *Biotechnol. Bioprocess Eng.*, 2011, 16, 337.
 - 24 Z. Wang, C. Y. Wang, H. R. Wang, H. Zhang, Y. L. Su, T. F. Ji and L. Wang, *Chin. Chem. Lett.*, 2014, **25**, 802.
- P. Steunenberg, M. Sijm, H. Zuilhof, J. P. M. Sanders, E. L. Scott and M. C. R. Franssen, *J. Org. Chem.*, 2013, **78**, 3802.
- 26 J. L. Wang, J. M. Xu, Q. Wu, D. S. Lv and X. F. Lin, *Tetrahedron*, 2009, **65**, 2531.
- 27 X. H. Wang, X. H. Zhang, S. J. Tu, F. Shi, X. Zou, S. Yan, Z. G. Han, W. J. Hao, X. D. Cao and S. S. Wu, J. Heterocyclic Chem., 2009, 46, 832.
- 28 K. Azizi and A. Heydari, RSC Adv., 2014, 4, 6508.
- 29 C. S. Yao, C. X. Yu, T. J. Li and S. J. Tu, *Chin. J. Chem.*, 2009, **27**, 1989.
- 30 Y. Yu, H. Y. Guo and X. J. Li, J. Heterocyclic Chem., 2010, 48, 1264.
- 25 31 M. G. Dekamin, M. Eslami and A. Maleki, *Tetrahedron*, 2013, 69, 1074.
 - 32 J. M. Khurana, B. Nand and P. Saluja, Tetrahedron, 2010, 66, 5637.
 - 33 J. M. Khurana, D. Magoo and A. Chaudhary, Synth. Comm., 2012, 42, 3211.
- 30 34 R. C. Rodrigues, C. Ortiz, A. Berenguer-Murcia, R. Torres and R. Fernández-Lafuente, *Chem. Soc. Rev.*, 2013, 42, 6290.
 - 35 E. N. Xun, X. L. Lv, W. Kang, J. X. Wang, H. Zhang, L. Wang and Z. Wang, Appl. Biochem. Biotechnol., 2012, 168, 697.
- 36 K. Hernandez and R. Fernández-Lafuente, *Enzyme Microb. Technol.*, 2011, **48**, 107.
 - 37 C. Garcia-Galan, A. Berenguer-Murcia, R. Fernández-Lafuente and R. C. Rodrigues, *Adv. Synth. Catal.*, 2011, **353**, 2885.