



Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: Y. Jia, T. Li, C. Yu, B. Jiang and C. Yao, *Org. Biomol. Chem.*, 2016, DOI: 10.1039/C5OB02336J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

A facile one-pot synthesis of 2,3-diarylated benzo[*b*]furans via relay NHC and palladium catalysis

Yanyan Jia, Tuanjie Li,* Chenxia Yu, Bo Jiang and Changsheng Yao*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

An efficient one-pot synthesis of 2,3-diarylated benzo[*b*]furans was realized through the relay catalysis of N-Heterocyclic Carbene (NHC) and palladium from substituted 2'-bromodiphenylbromomethanes and aryl aldehydes. The easy availability of the starting materials, good compatibility of catalysts, convergent assembly and concomitant modification of target scaffold, and potential utilization value of the products make this strategy attractive in organic synthesis.

Introduction

The wide distribution of benzo[*b*]furan skeleton in natural products^[1] (Figure 1) and synthetic chemical entities with biological relevance has drawn much attention to its rapid construction and facile modification, e.g., arylation due to the diverse bioactivity^[2] and electroluminescent property of arylated benzo[*b*]furans^[3]. To date, several strategies have been developed to successfully construct benzo[*b*]furan scaffold from acyclic precursors including aryloxyketones^[4], *o*-hydroxybenzophenones^[5], *o*-halogenated phenols^[6], *o*-alkynylphenols^[7], substituted benzylic alcohols^[8], *O*-aryl ketoximes^[9], and diazo compounds^[10] (Scheme 1, (a)~(h)). In addition, the effective introduction of an aryl group at C-2 or C-3 position of benzo[*b*]furan could also be achieved by transition-metal-catalyzed coupling reactions using unsubstituted benzo[*b*]furan as starting materials (Scheme 1, (i) and (j))^[11]. However, the harsh conditions, long reaction time, complexity of feedstocks and intricacy of operation associated with these protocols necessitate the development of new alternative for the convenient fabrication and decoration of benzo[*b*]furan framework.

Recently, N-Heterocyclic Carbenes (NHCs) have been proved to be versatile catalysts for many organic transformations^[12]. Apart from the traditional umpolung reaction of aldehydes (e.g., Benzoin condensation^[13] and Stetter reaction^[14]), numerous organic conversions including Michael addition^[15], transesterification^[16] and ring-opening polymerization^[17] could be promoted by NHCs efficiently. In addition, the potential of NHCs as ligands for transition metals has been pioneered^[18]. Interestingly, the compatibility of NHC

catalysts with a transition-metal catalyst has also been demonstrated in some coupling reactions^[19]. A recent report disclosed by Glorius et al showed that the NHC-catalyzed cross-coupling of aromatic aldehydes with activated alkyl halides delivered the aryl ketones readily (Scheme 2, (a))^[20], thus we envisioned that the cross-coupling of 2'-bromodiphenylbromomethanes with aryl aldehydes and the subsequent annulation catalysed by palladium may proceed in a one-pot manner without the isolation of the intermediates, which could pave a new avenue to benzo[*b*]furans (Scheme 2 (b)). To continue our work on NHCs-catalyzed cascade synthesis of heterocycles^[21], we shall herein report our recent work on sequential synthesis of benzo[*b*]furans through the relay catalysis of NHC/palladium.







Figure 1. Representatives of natural products containing benzo[*b*]furan core

School of Chemistry and Chemical Engineering,
Jiangsu Key Lab of Green Synthetic Chemistry For Functional Materials,
Jiangsu Normal University, Xuzhou, Jiangsu 221116, P R China.
†E-mail: ltj2008@jssu.edu.cn; csyao@jssu.edu.cn.
Electronic Supplementary Information (ESI) available: ¹H and ¹³C NMR spectra.
For ESI, See DOI: 10.1039/x0xx00000x

**Scheme 1.** Methods for the synthesis of diarylated benzo[b]furans**Scheme 2.** Previous work and this work

Results and discussion

According to the hypothesis described above, we chose 4-chlorobenzaldehyde **1a** and 1-bromo-2-(bromo(phenyl)methyl)benzene **2a** as model substrates to optimize reaction conditions (Table 1). Initially, the formation of desired product **3a** was not detected when the reaction was conducted at room temperature presented by Pd(OAc)₂ and PPh₃ in DMF. To our delight, product **3a** was obtained in 46% yield in the presence of **7** when the reaction temperature was elevated to 90 °C (Table 1, entries 1-4). Then assessment of palladium salts and ligands demonstrated that the combination of Pd(OAc)₂ and PPh₃ should be favourable (Table 1, entries 5-12 and entry 4) and the ratio of PPh₃ to Pd(OAc)₂ should be 5:1 (Table 1, entries 13-14 and entry 4). Next the optimization of solvents showed that the DMF was superior to other counterparts (Table 1, entries 15-17 and entry 13). The following investigation of the effect of base on this process illuminated that Cs₂CO₃ was better than K₂CO₃, tBuOK, DBU, Et₃N, K₃PO₄ (Table 1, entries 18-22 and entry 13). Finally, the screening of temperature showed that 95 °C should be preferable (Table 1, entries 23-24 and entry 13).


Table 1. Optimization of the reaction conditions

Entry	NHC/ [Pd]	Ligand (x mol%)	Base	Solvent	T (°C)	Yield (%) ^[a]
1	4/Pd(OAc) ₂	PPh ₃ (40)	Cs ₂ CO ₃	DMF	90	--
2	5/Pd(OAc) ₂	PPh ₃ (40)	Cs ₂ CO ₃	DMF	90	--
3	6/Pd(OAc) ₂	PPh ₃ (40)	Cs ₂ CO ₃	DMF	90	Trace
4	7/Pd(OAc) ₂	PPh ₃ (40)	Cs ₂ CO ₃	DMF	90	46
5	7/Pd(PPh ₃) ₄	--	Cs ₂ CO ₃	DMF	90	10
6	7/Pd(PPh ₃) ₂ Cl ₂	--	Cs ₂ CO ₃	DMF	90	Trace
7	7/Pd ₂ (dba) ₃	--	Cs ₂ CO ₃	DMF	90	Trace
8	7/PdCl ₂	PPh ₃ (40)	Cs ₂ CO ₃	DMF	90	--
9	7/Pd(OAc) ₂	P(<i>m</i> -tol) ₃ (40) ^[b]	Cs ₂ CO ₃	DMF	90	--
10	7/Pd(OAc) ₂	dppf (40) ^[b]	Cs ₂ CO ₃	DMF	90	Trace
11	7/Pd(OAc) ₂	dppe (40) ^[b]	Cs ₂ CO ₃	DMF	90	--
12	7/Pd(OAc) ₂	dppb (40) ^[b]	Cs ₂ CO ₃	DMF	90	--
13	7/Pd(OAc) ₂	PPh ₃ (50)	Cs ₂ CO ₃	DMF	90	78
14	7/Pd(OAc) ₂	PPh ₃ (60)	Cs ₂ CO ₃	DMF	90	53
15	7/Pd(OAc) ₂	PPh ₃ (50)	Cs ₂ CO ₃	THF	Ref.	-- ^[c]
16	7/Pd(OAc) ₂	PPh ₃ (50)	Cs ₂ CO ₃	DMSO	90	19
17	7/Pd(OAc) ₂	PPh ₃ (50)	Cs ₂ CO ₃	dioxane	Ref.	-- ^[c]
18	7/Pd(OAc) ₂	PPh ₃ (50)	K ₂ CO ₃	DMF	90	--
19	7/Pd(OAc) ₂	PPh ₃ (50)	tBuOK	DMF	90	--
20	7/Pd(OAc) ₂	PPh ₃ (50)	DBU	DMF	90	--
21	7/Pd(OAc) ₂	PPh ₃ (50)	Et ₃ N	DMF	90	--
22	7/Pd(OAc) ₂	PPh ₃ (50)	K ₃ PO ₄	DMF	90	--
23	7/Pd(OAc) ₂	PPh ₃ (50)	Cs ₂ CO ₃	DMF	95	81
24	7/Pd(OAc) ₂	PPh ₃ (50)	Cs ₂ CO ₃	DMF	100	75

[a]. Isolated yield [b]. P(*m*-tol)₃ = Tri(*m*-tolyl)phosphine; dppf = 1,1'-bis(diphenylphosphino)ferrocene; dppe = 1,2-Bis(diphenylphosphino)ethane; dppb = 1,4-bis(diphenylphosphino)butane [c]. Intermediate **3aa** (2-(2-bromophenyl)-1-(4-chlorophenyl)-2-phenylethanone) was isolated (See SI).

With the optimized reaction conditions in hand, a concise exploration of the substrate scope of this one-pot protocol was carried out. As shown in Table 2, the aryl aldehydes bearing the electron-donating or electron-withdrawing groups could participate in the reaction and gave the desired products in moderate to good yields (52–85 %, Table 2, entries 1–8), thus the electronic nature of the substituents had no obvious effect on this protocol. Besides, the heterocyclic aryl aldehydes were also tolerated and the expected products were isolated in satisfactory yields (78 and 75 %, Table 2, entries 9–10). To expand the substrate scope further, substituted 2'-bromodiphenylbromomethane was subject to this procedure and the corresponding benzo[*b*]furans (**3k–3n**) could be acquired in moderate yields under standard conditions (53–63 %, Table 2, entries 11–14). These results highlighted the wide application range of this method. All of the products were characterized by IR, NMR and HRMS and they were in good agreement with the reported data.


Table 2. Synthesis of benzo[*b*]furans

						
Entry	R ¹	R ²	R ³	Time (h)	Product	Yield (%) ^[a]
1	4-ClC ₆ H ₄	H	H	15	3a	81
2	4-FC ₆ H ₄	H	H	16	3b	75
3	C ₆ H ₅	H	H	18	3c	85
4	4-BrC ₆ H ₄	H	H	16	3d	52
5	4-CH ₃ C ₆ H ₄	H	H	24	3e	77
6	3-ClC ₆ H ₄	H	H	16	3f	64
7	3-CH ₃ C ₆ H ₄	H	H	16	3g	55
8	3,4-Cl ₂ C ₆ H ₃	H	H	16	3h	72
9	Thiophen-2-yl	H	H	16	3i	78
10	Furan-2-yl	H	H	16	3j	75
11	4-ClC ₆ H ₄	H	4-CH ₃	17	3k	62
12	4-ClC ₆ H ₄	5-Cl	H	17	3l	53
13	Thiophen-2-yl	5-Cl	H	16	3m	63
14	C ₆ H ₅	5-Cl	H	16	3n	60


[a]. Isolated yield

A mechanism to rationalize the formation of benzo[*b*]furans was presented in **Scheme 3**. The nucleophilic attack of Breslow intermediate **A**, generated from aldehydes **1** and NHC, on 2'-bromodiphenylbromomethanes **2** provided

intermediate **B**. The subsequent collapse of **B** afforded the key intermediate **C** and regenerated NHC catalyst^[12,20,21]. The following keto-enol tautomerization and intramolecular C-O bond formation assisted by palladium gave rise to the ultimate product **3**^[19].

**Scheme 3.** Plausible catalytic cycle

To shed some light on the reaction mechanism, a control experiment was carried out (**Scheme 4** and see SI). Intermediate **3aa** could undergo further intramolecular cyclization to furnish product **3a** under the catalysis of Pd. Thus, **3aa** should play a pivotal role in this one-pot synthesis of benzo[*b*]furan.

**Scheme 4.** Pd-catalyzed cyclization of intermediate **3aa**

Conclusions

In summary, we have developed an efficient alternative approach to 2,3-diarylated benzo[*b*]furans by virtue of sequential catalysis of NHC and palladium from substituted 2'-bromodiphenylbromomethanes and aryl aldehydes. This protocol featured operational simplicity, easy accessibility of the raw materials, good tolerance of catalysts, and straightforward assembly with simultaneous modification of target scaffold. Other study aimed at the expansion of the application of the relay catalytic system consisting of NHC and metal is underway in our lab.

Experimental

Typical procedure for synthesis of benzo[*b*]furans. An oven-dried 25-mL flask equipped with a magnetic stir bar was charged with the thiazolium salt **7** (18.5 mg, 0.05 mmol), Cs₂CO₃ (179.2 mg, 0.55 mmol), freshly distilled DMF (5 mL), aromatic aldehydes **1** (0.5

ARTICLE

Journal Name

mmol), 2'-bromodiphenylbromomethanes **2** (0.5 mmol), Pd(OAc)₂ (11.2 mg, 0.05 mmol) and PPh₃ (65.6 mg, 0.25 mmol). The mixture was stirred at 95 °C until completion (monitored by TLC). Deionized water (5 mL) was added and the mixture then extracted with EtOAc. The organic layer was dried over MgSO₄. The desired products **3** were obtained through filtration, concentration in vacuo and purification by column chromatography (silica gel, petroleum).

2-(4-chlorophenyl)-3-phenylbenzofuran (3a). 123 mg, 81 % yield; white solid; M.P.: 90 - 92 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.60 - 7.57 (m, 2H, ArH), 7.55 (d, *J* = 8.0 Hz, 1H, ArH), 7.50 - 7.40 (m, 6H, ArH), 7.36 - 7.32 (m, 1H, ArH), 7.30 - 7.22 (m, 3H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 149.3, 134.2, 132.5, 130.1, 129.7, 129.1, 129.1, 128.7, 128.2, 127.8, 124.9, 123.1, 120.1, 118.0, 111.1; IR (potassium bromide) (ν, cm⁻¹): 1618, 1608, 1582, 1560, 1483, 1449, 1402, 1253, 1201, 1089, 1064, 1014, 962, 840, 749, 695; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₀H₁₄ClO: 305.0733, found: 305.0723.

2-(4-fluorophenyl)-3-phenylbenzofuran (3b). 108 mg, 75 % yield; white solid; M.P.: 87 - 88 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.66 - 7.60 (m, 2H, ArH), 7.55 (d, *J* = 8.0 Hz, 1H, ArH), 7.50 - 7.39 (m, 6H, ArH), 7.35 - 7.31 (m, 1H, ArH), 7.26 - 7.22 (m, 1H, ArH) 7.04 - 6.98 (m, 2H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 162.7 (*J*_{CF} = 250.0 Hz), 153.9, 149.6, 132.6, 130.1, 129.7, 129.0, 128.9 (*J*_{CF} = 10.0 Hz), 127.7, 126.9 (*J*_{CF} = 10.0 Hz), 124.7, 123.0, 120.0, 117.2 (*J*_{CF} = 1.0 Hz), 115.5 (*J*_{CF} = 20.0 Hz), 111.1; IR (potassium bromide) (ν, cm⁻¹): 1621, 1602, 1510, 1450, 1407, 1373, 1253, 1219, 1203, 1164, 1065, 963, 840, 797, 751, 696; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₀H₁₄FO: 289.1023, found: 289.1023.

2,3-diphenylbenzofuran (3c). 114 mg, 85 % yield; white solid, M.P.: 115 - 116 °C (reported 122-123 °C)^[22a]; ¹H NMR (400 MHz, CDCl₃): δ 7.66 (d, *J* = 8.0 Hz, 2H, ArH), 7.57 - 7.39 (m, 7H, ArH), 7.35 - 7.29 (m, 4H, ArH), 7.25 - 7.22 (m, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 150.5, 132.8, 130.7, 130.2, 129.8, 129.0, 128.4, 128.3, 127.6, 127.0, 124.7, 122.9, 120.0, 117.5, 111.1; IR (potassium bromide) (ν, cm⁻¹): 1601, 1568, 1496, 1455, 1440, 1368, 1339, 1203, 1187, 1109, 1062, 1027, 960, 827, 747, 693; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₀H₁₅O: 271.1174, found: 271.1174.

2-(4-bromophenyl)-3-phenylbenzofuran (3d). 90 mg, 52 % yield; white solid; M.P.: 89 - 91 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.56 - 7.40 (m, 11H, ArH), 7.37 - 7.33 (m, 1H, ArH), 7.23 (d, *J* = 7.6 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 149.4, 132.5, 131.6, 130.1, 129.7, 129.6, 129.1, 128.4, 127.9, 125.0, 123.1, 122.5, 120.1, 118.1, 111.1; IR (potassium bromide) (ν, cm⁻¹): 1603, 1577, 1497, 1482, 1451, 1396, 1253, 1076, 1006, 962, 894, 825, 772, 747, 703; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₀H₁₄BrO: 349.0228, found: 349.0227.

3-phenyl-2-p-tolylbenzofuran (3e). 109 mg, 77 % yield; white solid; M.P.: 83 - 85 °C (reported 96 - 97 °C)^[22b]; ¹H NMR (400 MHz, CDCl₃): δ 7.55 (d, *J* = 8.0 Hz, 3H, ArH), 7.52 - 7.44 (m, 5H, ArH), 7.42 - 7.38 (m, 1H, ArH), 7.34 - 7.30 (m, 1H, ArH), 7.25 - 7.21 (m, 1H, ArH), 7.13 (d, *J* = 8.0 Hz, 2H, ArH), 2.35 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃):

δ 153.9, 150.8, 138.4, 133.0, 130.3, 129.8, 129.1, 128.9, 127.8, 127.5, 127.0, 124.4, 122.8, 119.9, 116.8, 111.0, 100.0, 21.4; IR (potassium bromide) (ν, cm⁻¹): 2916, 1604, 1512, 1453, 1442, 1369, 1255, 1202, 1181, 1110, 1076, 1065, 962, 817, 770, 739, 702; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₁H₁₇O: 285.1279, found: 285.1278.

2-(3-chlorophenyl)-3-phenylbenzofuran (3f). 194 mg, 64 % yield; white solid; M.P.: 74 - 76 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.70 (s, 1H, ArH), 7.53 (d, *J* = 8.0 Hz, 1H, ArH), 7.48 - 7.40 (m, 7H, ArH), 7.32 (t, *J* = 7.4 Hz, 1H, ArH), 7.24 - 7.19 (m, 2H, ArH), 7.16 (t, *J* = 7.8 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 148.8, 134.4, 132.3, 132.3, 130.0, 129.6, 129.6, 129.1, 128.2, 127.9, 126.7, 125.1, 124.9, 123.1, 120.2, 118.6, 111.1; IR (potassium bromide) (ν, cm⁻¹): 1598, 1560, 1497, 1476, 1452, 1370, 1257, 1207, 1114, 1080, 1062, 965, 843, 742, 699; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₀H₁₄ClO: 305.0733, found: 305.0733.

3-phenyl-2-m-tolylbenzofuran (3g). 78 mg, 55 % yield; white solid; M.P.: 68 - 70 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.55 (d, *J* = 8.0 Hz, 2H, ArH), 7.51 - 7.49 (m, 3H, ArH), 7.47 - 7.43 (m, 2H, ArH), 7.41 - 7.36 (m, 2H, ArH), 7.34 - 7.30 (m, 1H, ArH), 7.23 - 7.21 (m, 1H, ArH), 7.16 (t, *J* = 7.8 Hz, 1H, ArH), 7.09 (d, *J* = 7.6 Hz, 1H, ArH), 2.31 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 153.9, 150.7, 138.1, 132.9, 130.5, 130.2, 129.7, 129.1, 128.9, 128.2, 127.6, 127.6, 124.6, 124.3, 122.9, 120.0, 117.4, 111.1, 21.4; IR (potassium bromide) (ν, cm⁻¹): 2922, 1605, 1497, 1453, 1372, 1340, 1260, 1221, 1183, 1064, 1010, 965, 869, 769, 746, 699; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₁H₁₇O: 285.1279, found: 285.1273.

2-(3,4-dichlorophenyl)-3-phenylbenzofuran (3h). 122 mg, 72 % yield; white solid; M.P.: 116 - 118 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.81 (d, *J* = 2.0 Hz, 1H, ArH), 7.56 (d, *J* = 8.0 Hz, 1H, ArH), 7.52 - 7.42 (m, 7H, ArH), 7.40 - 7.38 (m, 1H, ArH), 7.35 (t, *J* = 8.4 Hz, 2H, ArH), 7.28 - 7.24 (m, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 147.9, 132.8, 132.2, 132.1, 130.6, 130.4, 130.0, 129.6, 129.2, 128.4, 128.1, 125.9, 125.4, 123.2, 120.3, 119.0, 111.2; IR (potassium bromide) (ν, cm⁻¹): 1595, 1548, 1449, 1369, 1274, 1261, 1206, 1133, 1071, 1028, 964, 887, 817, 768, 743, 698; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₂₀H₁₃Cl₂O: 339.0343, found: 339.0333.

3-phenyl-2-(thiophen-2-yl)benzofuran (3i). 108 mg, 78 % yield; white solid; M.P.: 139 - 140 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.57 - 7.49 (m, 5H, ArH), 7.46 - 7.42 (m, 2H, ArH), 7.34 - 7.30 (m, 2H, ArH), 7.25 (dd, *J*₁ = 4.8 Hz, *J*₂ = 1.2 Hz, 1H, ArH), 7.24 - 7.20 (m, 1H, ArH), 6.99 (dd, *J*₁ = 5.2 Hz, *J*₂ = 3.6 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 153.8, 146.6, 132.7, 132.0, 130.2, 130.0, 128.9, 128.0, 127.4, 126.0, 125.7, 124.7, 123.1, 119.8, 116.8, 111.0; IR (potassium bromide) (ν, cm⁻¹): 1604, 1452, 1374, 1337, 1220, 1196, 1177, 1109, 1080, 957, 931, 851, 831, 798, 771, 748, 707, 698; HRMS (APCI) *m/z*: Calcd. for [M+H]⁺ C₁₈H₁₃OS: 277.0687, found: 277.0684.

2-(furan-2-yl)-3-phenylbenzofuran (3j). 97 mg, 75 % yield; white solid; M.P.: 82 - 84 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.60 (d, *J* = 8.0 Hz, 2H, ArH), 7.56 - 7.48 (m, 4H, ArH), 7.44 - 7.40 (m, 2H, ArH), 7.34 (t, *J* = 7.8 Hz, 1H, ArH), 7.25 - 7.20 (m, 1H, ArH), 6.66 (d, *J* = 3.6 Hz,

¹H, ArH), 6.45–6.44 (m, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 146.3, 145.9, 144.3, 142.9, 131.7, 129.7, 128.6, 127.7, 124.9, 123.2, 120.1, 111.5, 111.2, 109.2; IR (potassium bromide) (ν, cm⁻¹): 1734, 1718, 1701, 1654, 1647, 1636, 1559, 1541, 1458, 1442, 1091, 1012, 740, 700; HRMS (APCI) m/z: Calcd. for [M+H]⁺ C₁₈H₁₃O₂: 261.0916, found: 261.0915.

2-(4-chlorophenyl)-3-p-tolylbenzofuran (3k). 92 mg, 62 % yield; white solid; M.P.: 110–112 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.61–7.58 (m, 2H, ArH), 7.53 (d, *J* = 8.0 Hz, 1H, ArH), 7.49–7.47 (m, 1H, ArH), 7.37–7.34 (m, 2H, ArH), 7.33–7.31 (m, 1H, ArH), 7.29–7.26 (m, 4H, ArH), 7.23–7.21 (m, 1H, ArH), 2.44 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 154.0, 149.2, 137.6, 134.0, 130.2, 129.8, 129.5, 129.4, 129.3, 128.7, 128.1, 124.9, 123.0, 120.2, 118.0, 111.1, 21.4; IR (potassium bromide) (ν, cm⁻¹): 2919, 1514, 1487, 1450, 1254, 1201, 1108, 1085, 1065, 1012, 965, 898, 831, 748; HRMS (APCI) m/z: Calcd. for [M+H]⁺ C₂₁H₁₆ClO: 319.0890, found: 319.0890.

5-chloro-2-(4-chlorophenyl)-3-phenylbenzofuran (3l). 89 mg, 53 % yield; white solid; M.P.: 79–81 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.58–7.57 (m, 1H, ArH), 7.56–7.55 (m, 1H, ArH), 7.49–7.47 (m, 2H, ArH), 7.45–7.44 (m, 5H, ArH), 7.30–7.29 (m, 2H, ArH), 7.28–7.27 (m, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 152.3, 150.8, 134.6, 131.8, 131.5, 129.6, 129.2, 128.8, 128.6, 128.2, 125.1, 119.7, 117.6, 112.1; IR (potassium bromide) (ν, cm⁻¹): 1610, 1558, 1541, 1507, 1498, 1486, 1447, 1402, 1367, 1258, 1206, 1087, 1058, 1014, 967, 833, 798; HRMS (APCI) m/z: Calcd. for [M+H]⁺ C₂₀H₁₃Cl₂O: 339.0338, found: 339.0326.

5-chloro-3-phenyl-2-(thiophen-2-yl)benzofuran (3m). 68 mg, 63 % yield; white solid; M.P.: 98–100 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.53–7.52 (m, 4H, ArH), 7.44 (d, *J* = 8.4 Hz, 1H, ArH), 7.38 (d, *J* = 2.0 Hz, 1H, ArH), 7.34 (dd, *J*₁ = 3.8 Hz, *J*₂ = 1.0 Hz, 1H, ArH), 7.30–7.27 (m, 2H, ArH), 7.25 (d, *J* = 2.4 Hz, 1H, ArH), 7.00 (dd, *J*₁ = 5.2 Hz, *J*₂ = 3.6 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 152.1, 148.0, 132.1, 131.6, 131.3, 129.9, 129.1, 128.8, 128.3, 127.5, 126.6, 126.2, 124.8, 119.4, 116.3, 112.0; IR (potassium bromide) (ν, cm⁻¹): 1616, 1577, 1507, 1473, 1448, 1327, 1256, 1218, 1202, 1058, 1019, 966, 870, 804, 716, 693; HRMS (APCI) m/z: Calcd. for [M+H]⁺ C₁₈H₁₁ClOS: 311.0297, found: 311.0298.

5-chloro-2,3-diphenylbenzofuran (3n). 91 mg, 63 % yield; white solid; M.P.: 101–102 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.65–7.63 (m, 2H, ArH), 7.48–7.45 (m, 6H, ArH), 7.33–7.29 (m, 4H, ArH), 7.27 (d, *J* = 2.0 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 152.3, 151.9, 132.1, 131.7, 130.2, 129.6, 129.1, 128.7, 128.6, 128.5, 127.9, 127.0, 124.8, 119.6, 117.1, 112.1; IR (potassium bromide) (ν, cm⁻¹): 1603, 1505, 1450, 1439, 1321, 1257, 1206, 1081, 1066, 1057, 1026, 966, 915, 803, 764; HRMS (APCI) m/z: Calcd. for [M+H]⁺ C₂₀H₁₄ClO: 305.0733, found: 305.0736.

Acknowledgements

We are grateful for financial support by National Natural Science Foundation of China (Nos. 21271091, 21372101), a project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institution.

Notes and references

- 1 For natural products containing the benzo[b]furan framework, see: a) P. Cagniant and D. Cagniant, *Adv. Heterocycl. Chem.*, 1975, **18**, 337; b) K.-S. Song and I. Raskin, *J. Nat. Prod.*, 2002, **65**, 76; c) B. Sritularak, K. Likhitwitayawuid, J. Conrad, B. Vogler, I. Klaiber and W. Kraus, *J. Nat. Prod.*, 2002, **65**, 589; d) T. Pacher, C. Seger, D. Engelmeier, S. Vajrodaya, O. Hofer and H. Greger, *J. Nat. Prod.*, 2002, **65**, 820; e) H. Kwiecien, M. Smist and M. Kowalewska, *Curr. Org. Synth.*, 2012, **9**, 529.
- 2 For biological properties, see: a) K. M. Zareba, *Drugs Today*, 2006, **42**, 75; b) K. M. Dawood, H. Abdel-Gawad, E. A. Rageb, M. Ellithy and H. A. Mohamed, *Bioorg. Med. Chem.*, 2006, **14**, 3672; c) A. Ladurner, A. G. Atanasov, E. H. Heiss, L. Baumgartner, S. Schwaiger, J. M. Rollinger, H. Stuppner and V. M. Dirsch, *Biochem. Pharmacol.*, 2012, **84**, 804; d) P. Yadav, P. Singh and A. K. Tewari, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 2251; e) T. Yempala, J. P. Sridevi, P. Yogeeswari, D. Sriram and S. Kantevari, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 5393; f) T. S. Chundawat, N. Sharma and S. Bhagat, *Medicinal Chemistry*, 2014, **10**, 409; g) G. S. Hassan, S. M. Abou-Seri, G. Kamel and M. M. Ali, *Eur. J. Med. Chem.*, 2014, **76**, 482.
- 3 a) S. Ikawa, S. Yagi, T. Maeda, H. Nakazumi, H. Fujiwara, S. Koseki and Y. Sakurai, *Inorg. Chem. Commun.*, 2013, **38**, 14; b) A. Tronnier, N. Nischan, S. Metz, G. Wagenblast, I. Muenster, T. Strassner, *Eur. J. Inorg. Chem.*, 2014, **2014**, 256; c) C. W. Lee, J.-A. Seo, M.-S. Gong, J. Y. Lee, *Chem. Eur. J.*, 2013, **19**, 1194.
- 4 a) I. Kim and J. Choi, *Org. Biomol. Chem.*, 2009, **7**, 2788; b) K.-S. Kim and I.-Y. Kim, *Org. Lett.*, 2010, **12**, 5314; c) J. H. Lee, M. Kim, and I. Kim, *J. Org. Chem.*, 2014, **79**, 6153; d) Y.-S. Bao, A. Bao, Z. Bao, M. Jia and M. Baiyin, *Org. Biomol. Chem.*, 2015, **13**, 4179.
- 5 a) F. Schevenels and I. E. Marko, *Org. Lett.*, 2012, **14**, 1298; b) M. J. Moure, R. SanMartin and E. Dominguez, *Angew. Chem., Int. Ed.*, 2012, **51**, 3220.
- 6 a) R.-P. Wang, S. Mo, Y.-Z. Lu and Z.-M. Shen, *Adv. Synth. Catal.*, 2011, **353**, 713; b) J.-R. Wang and K. Manabe, *J. Org. Chem.*, 2010, **75**, 5340; c) G. S. Gill, D. W. Grobely, J. H. Chaplin and B. L. Flynn, *J. Org. Chem.*, 2008, **73**, 1131; d) C. Eidamshaus and J. D. Burch, *Org. Lett.*, 2008, **10**, 4211; e) J.-P. Wan, H. Wang, Y. Liu, and H. Ding, *Org. Lett.*, 2014, **16**, 5160.
- 7 a) N. Matsuda, K. Hirano, T. Satoh and M. Miura, *J. Org. Chem.*, 2012, **77**, 617; b) C. Kanazawa, K. Goto and M. Terada, *Chem. Commun.*, 2009, 5248; c) C. Martínez, R. Álvarez and J. M. Aurecochea, *Org. Lett.*, 2009, **11**, 1083; d) Y. Liang, S. Tang, X.-D. Zhang, L.-Q. Mao, Y.-X. Xie and J.-H. Li, *Org. Lett.*, 2006, **8**, 3017; e) N. Isono and M. Lautens, *Org. Lett.*, 2009, **11**, 1329.
- 8 B. Anxionnat, D. Gomez Pardo, G. Ricci, K. Rossen, and J. Cossy, *Org. Lett.*, 2013, **15**, 3876.
- 9 a) A. A. Tabolin and S. L. Ioffe, *Chem. Rev.*, 2014, **114**, 5426; b) O. Miyata, N. Takeda and T. Naito, *Org. Lett.*, 2004, **6**, 1761; c) N. Takeda, O. Miyata and T. Naito, *Eur. J. Org. Chem.*, 2007, 1491.
- 10 a) M. P. Doyle, M. Ratnikov and Y. Liu, *Org. Biomol. Chem.*, 2011, **9**, 4007; b) X. Xu, M. O. Ratnikov, P. Y. Zavalij and M. P. Doyle, *Org. Lett.*, 2011, **13**, 6122; c) C. Soldi, K. N. Lamb, R. A.

- Squitieri, M. González-López, M. J. Di Maso and J. T. Shaw, *J. Am. Chem. Soc.*, 2014, **136**, 15142.
- 11 For direct C-3 arylation of benzofurans, see: a) M. Ionita, J. Roger and H. Doucet, *ChemSusChem*, 2010, **3**, 367; b) B. Liégault, I. Petrov, S. I. Gorelsky and K. Fagnou, *J. Org. Chem.*, 2010, **75**, 1047; c) A. Carrère, D. Brinet, J.-C. Florent, P. Rousselle and E. Bertounesque, *J. Org. Chem.*, 2012, **77**, 1316. For direct C-2 arylation of benzofurans, see: d) T. A. Dwight, N. R. Rue, D. Charyk, R. Josselyn and B. DeBoef, *Org. Lett.*, 2007, **9**, 3137; e) S. Matsuda, M. Takahashi, D. Monguchi and A. Mori, *Synlett*, 2009, 1941; f) T. Dao-Huy, M. Haider, F. Glatz, M. Schnüerch, and M. D. Mihovilovic, *Eur. J. Org. Chem.*, 2014, **2014**, 8119.
- 12 For selected reviews, please see: a) D. Enders, O. Niemeier and A. Henseler, *Chem. Rev.*, 2007, **107**, 5606; b) N. Marion, S. Diez-Gonzalez and S. P. Nolan, *Angew. Chem., Int. Ed.*, 2007, **46**, 2988; c) A. T. Biju, N. Kuhl and F. Glorius, *Acc. Chem. Res.*, 2011, **44**, 1182; d) X. Bugaut and F. Glorius, *Chem. Soc. Rev.*, 2012, **41**, 3511; e) A. Grossmann and D. Enders, *Angew. Chem., Int. Ed.*, 2012, **51**, 314; f) D. T. Cohen and K. A. Scheidt, *Chem. Sci.*, 2012, **3**, 53; g) X.-Y. Chen and S. Ye, *Synlett*, 2013, **24**, 1614; h) D. C. M. Albanese and N. Gaggero, *Eur. J. Org. Chem.*, 2014, **2014**, 5631; i) J. Mahatthananchai and J. W. Bode, *Acc. Chem. Res.*, 2014, **47**, 696. j) D. M. Flanagan, F. Romanov-Michailidis, N. A. White and T. Rovis, *Chem. Rev.*, 2015, **115**, 9307; k) R. S. Menon, A. T. Biju and V. Nair, *Chem. Soc. Rev.*, 2015, **44**, 5040.
- 13 a) Y. Cheng, J.-H. Peng, Y.-J. Li, X.-Y. Shi, M.-S. Tang and T.-Y. Tan, *J. Org. Chem.*, 2011, **76**, 1844; b) C. J. Collett, R. S. Massey, J. E. Taylor, O. R. Maguire, A. C. O'Donoghue and A. D. Smith, *Angew. Chem., Int. Ed.*, 2015, **54**, 6887.
- 14 a) D. A. DiRocco, E. L. Noey, K. N. Houk and T. Rovis, *Angew. Chem., Int. Ed.*, 2012, **51**, 2391; b) M.-Q. Jia and S.-L. You, *Chem. Commun.*, 2012, **48**, 6363; c) N. E. Wurz, C. G. Daniliuc and F. Glorius, *Chem. Eur. J.*, 2012, **18**, 16297; d) A. Patra, A. Bhunia and A. T. Biju, *Org. Lett.*, 2014, **16**, 4798.
- 15 a) T. Boddaert, Y. Coquerel and J. Rodriguez, *Adv. Synth. Catal.*, 2009, **351**, 1744; b) T. Boddaert, Y. Coquerel and J. Rodriguez, *Chem. Eur. J.*, 2011, **17**, 2266; c) E. M. Phillips, M. Riedrich and K. A. Scheidt, *J. Am. Chem. Soc.*, 2010, **132**, 13179; d) Q. Kang and Y. hang, *Org. Biomol. Chem.*, 2011, **9**, 6715; e) D. Enders, K. Lüttgen and A. A. Narine, *Synthesis*, 2007, 959; f) M. Hans, L. Delaude, J. Rodriguez, and Y. Coquerel, *J. Org. Chem.*, 2014, **79**, 2758.
- 16 a) G. A. Grasa, T. Güeveli, R. Singh and S. P. Nolan, *J. Org. Chem.*, 2003, **68**, 2812; b) J. A. M. Hepperle, R. C. Samanta and A. Studer, *Synlett*, 2013, **24**, 1233; c) I. Chiarotto, M. Feroci, G. Sotgiu and A. Inesi, *Eur. J. Org. Chem.*, 2013, **2013**, 326.
- 17 a) M. Li, H. Song and B. Wang, *Eur. J. Inorg. Chem.*, 2015, **2015**, 4055; b) P. Horeglad, M. Cybularczyk, B. Trzaskowski, G. Z. Zukowska, M. Dranka and J. Zachara, *Organometallics*, 2015, **34**, 3480.
- 18 a) O. Schuster, L. Yang, H. G. Raubenheimer and M. Albrecht, *Chem. Rev.*, 2009, **109**, 3445; b) E. Levin, E. Ivry, C. E. Diesendruck and N. G. Lemcoff, *Chem. Rev.*, 2015, **115**, 4607; c) A. V. Zhukhovitskiy, M. J. MacLeod and J. A. Johnson, *Chem. Rev.*, 2015, **115**, 11503; d) Z.-K. Xiao, H.-Y. Yin and L.-X. Shao, *Org. Lett.*, 2013, **15**, 1254; e) H.-Y. Yin, M.-Y. Liu and L.-X. Shao, *Org. Lett.*, 2013, **15**, 6042; f) S.-C. Yin, Q. Zhou, X.-Y. Zhao and L.-X. Shao, *J. Org. Chem.*, 2015, **80**, 8916.
- 19 a) R. Lebeuf, K. Hirano and F. Glorius, *Org. Lett.*, 2008, **10**, 4243; b) T. Nemoto, T. Fukuda and Y. Hamada, *Tetrahedron Lett.*, 2006, **47**, 4365; c) Z. Chen, X. Yu and J. Wu, *Chem. Commun.*, 2010, **46**, 6356; d) D. A. DiRocco and T. Rovis, *J. Am. Chem. Soc.*, 2012, **134**, 8094.
- 20 M. Padmanaban, A. T. Biju and F. Glorius, *Org. Lett.*, 2011, **13**, 98.
- 21 a) C. Yao, D. Wang, J. Lu, T. Li, W. Jiao and C. Yu, *Chem. Eur. J.*, 2012, **18**, 1914; b) Z. Xiao, C. Yu, T. Li, X.-S. Wang and C. Yao, *Org. Lett.*, 2014, **16**, 3632; c) Y. Xie, C. Yu, T. Li, S. Tu and C. Yao, *Chem. Eur. J.*, 2015, **21**, 5355; d) Y. Que, T. Li, C. Yu, X.-S. Wang and C. Yao, *J. Org. Chem.*, 2015, **80**, 3289; e) Y. Que, Y. Xie, T. Li, C. Yu, S. Tu and C. Yao, *Org. Lett.*, 2015, DOI: 10.1021/acs.orglett.5b03223.
- 22 a) F. Churruarín, R. San Martín, I. Tellitu and E. Domínguez, *Eur. J. Org. Chem.*, 2005, **12**, 2481; b) V. R. Veeramani, M. Pal and K. R. Yeleswarap, *Tetrahedron*, 2003, **59**, 3283.