



Cite this: *Chem. Commun.*, 2015, 51, 6226

Received 23rd January 2015,
Accepted 4th March 2015

DOI: 10.1039/c5cc00677e

www.rsc.org/chemcomm

Toward the discovery of dual inhibitors for botulinum neurotoxin A: concomitant targeting of endocytosis and light chain protease activity†

Hajime Seki,^a Song Xue,^a Mark S. Hixon,^b Sabine Pellett,^c Marek Remeš,^d
Eric A. Johnson^c and Kim D. Janda^{*ae}

Dyngo-4a™ has been found to be an endocytic inhibitor of BoNT/A neurotoxicity through dynamin inhibition. Herein, we demonstrate this molecule to have a previously unrecognized dual activity against BoNT/A, dynamin–protease inhibition. To establish the importance of this dual activity, detailed kinetic analysis of Dyngo-4a's inhibition of BoNT/A metalloprotease as well as cellular and animal toxicity studies have been described. The research presented is the first polypharmacological approach to counteract BoNT/A intoxication.

Clostridium botulinum, an anaerobic bacterium, produces botulinum neurotoxin (BoNT), which is considered to be the most lethal of known toxins. BoNT is a 150 kDa protein consisting of a 100 kDa heavy chain and a 50 kDa light chain.¹ The heavy chain is responsible for cellular surface recognition, toxin internalization and translocation of the light chain. The light chain (LC), a zinc metalloprotease, is the etiological agent responsible for neurotoxicity, namely, proteolysis of SNARE (soluble *N*-ethylmaleimide sensitive factor attachment protein receptor) proteins.² Upon the cleavage of SNAREs, vesicles containing acetylcholine are unable to fuse with the presynaptic neuromuscular junction. As a result, the release of acetylcholine is blocked, leading to flaccid paralysis (botulism) and potential death.

Among the eight serotypes for the BoNTs (A–H), A, B, and E cause botulism in humans.³ BoNT/A and E target SNAP-25

(synaptosome-associated protein-25 kDa), and B targets synaptobrevin. In particular, intoxication from BoNT/A persists up to several months and its potency is greater than any other serotypes (LD₅₀: 1 ng kg^{−1}).⁴ BoNT/A is sold under the trade name of BOTOX® and has been used for the treatment of medical conditions such as migraine and facial wrinkle reduction. Yet, it is also recognized as a potential bioweapon by Center for Disease Control and Prevention (CDC) as one of the six category A agents.⁵ Current treatments for BoNT/A intoxication are limited to antitoxin and vaccination, which can readily remove the toxin from circulation; however, such treatments become ineffective once cellular internalization of the toxin takes place.⁶

With increasing threats of terrorism in the last two decades, the discovery of therapeutic treatments for BoNT/A intoxication is of utmost importance and urgency, which cannot be overstated.⁷ Because BoNT/A's metalloprotease is the agent responsible for its neurotoxicity our laboratory as well as others have been actively investigating the development of LC inhibitors including active site and exosite inhibitors.⁸ In terms of active site inhibitors, we have developed hydroxamic acids, taking advantage of their strong coordinating nature to zinc.⁹ We have also discovered exosite inhibitors from natural product screens: chicoric acid and lomofungin (1 & 3, Fig. 1) are prototypical examples.¹⁰ Although, it has been a major challenge to translate potent active site enzyme inhibitor efficacy to cellular and ultimately animal lethality models, we recently demonstrated that a prodrug approach has enabled hydroxamic acids to gain traction in order to facilitate cellular uptake and attenuated LC activity.¹¹

An alternative rational approach to neutralize BoNT/A is to prevent its internalization from the cellular surface. In this regard, Meunier and co-workers reported how the dynamin inhibitor, Dyngo-4a™, thwarted BoNT/A internalization in hippocampal neurons and delayed the onset of botulism in an *in vivo* mouse model (4, Fig. 1).¹² Dyngo-4a is a pharmacologically refined analogue of Dynasore, which is a known inhibitor of dynamin.¹³ Dynamin is a mechanochemical GTPase responsible for the vesicle scission step in endocytosis.¹⁴ From the standpoint of BoNT/A neurotoxicity,

^a Departments of Chemistry and Immunology and Microbial Sciences, The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA. E-mail: kdjanda@scripps.edu; Tel: +1 858 784-2516

^b Modeling & Simulation Global DMPK, Takeda California, Inc., 10410 Science Center Drive, San Diego, CA 92121, USA

^c Department of Bacteriology, University of Wisconsin, 1550 Linden Drive, Madison, Wisconsin 53706, USA

^d Department of Organic Chemistry, Faculty of Science, Charles University, Hlavova 2030/8 128 43 Praha 2, Czech Republic

^e Worm Institute for Research and Medicine (WIRM), The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

† Electronic supplementary information (ESI) available: Compound synthesis and characterization, kinetics data, cell assay and animal study data. See DOI: 10.1039/c5cc00677e




Fig. 1 Structures and targets of exosite inhibitors (1–3) and endocytic inhibitor (4).

Dyngo-4a's mechanism of action was attributed to the fact that BoNT/A mainly utilizes a dynamin-dependent endocytic pathway to enter neuronal cells.¹²

In the effort to search for molecules that could inhibit the protease of BoNT/A, we turned our focus to the polyphenolic scaffold embedded within Dyngo-4a. We suspected that this Dyngo-4a's phenolic architecture could serve as an exosite LC inhibitor, based on its structural similarity to chicoric acid. Thus, Dyngo-4a was tested *in vitro* employing both FRET-based SNAPtide and LC/MS-based 66mer assays.¹⁵ As anticipated the compound was inactive in the SNAPtide test, while it showed inhibitory activity in the 66mer assay, indicating that Dyngo-4a's binding was exosite driven. To further define Dyngo-4a's inhibition profile, detailed kinetics of the compound was examined with varied concentrations of the 66mer substrate (Fig. 2). Dyngo-4a showed a competitive inhibition mechanism with $K_i = 0.32 \pm 0.05 \mu\text{M}$.

As a means to understand Dyngo-4a's mechanism of inhibition, we investigated Dyngo-4a's ability to access the α and β -exosites of BoNT/A's LC. To begin, a dual inhibition assay between Dyngo-4a and chicoric acid *i*-Pr ester (2, Fig. 1), an α -exosite inhibitor, was undertaken.¹⁶ As a frame of reference, 2 presents a competitive inhibition mechanism with $K_i = 1.8 \pm 0.3 \mu\text{M}$. Here we observed




Fig. 2 Global fit using a competitive inhibition mechanism.

Dyngo-4a vs ChA *i*-Pr ester 2




Fig. 3 Mutually exclusive fit for Dyngo-4a and chicoric acid *i*-Pr ester.

Dyngo-4a vs Lomofungin





Fig. 4 Mutually exclusive fit for Dyngo-4a and lomofungin.

mutually exclusive binding, indicating that the two molecules cannot bind simultaneously (Fig. 3). A parallel assay now engaging lomofungin, a presumed β -exosite inhibitor, also presented mutual exclusivity (Fig. 4). Moreover, Dyngo-4a demonstrated complete inhibition of the BoNT/A LC at 20 μM , by a competitive inhibition mechanism, whereas chicoric acid only displayed partial inhibition with a non-competitive profile.^{10a} In a final attempt to further define the BoNT/A LC and Dyngo-4a's interactions, a competitive assay between Dyngo-4a and an active site hydroxamate inhibitor, previously validated from crystallographic analysis, was conducted.^{9b,d} Here non-mutually exclusive binding was observed, *i.e.* both can bind simultaneously with an enhancement factor (α) of 1.6 ± 0.8 (ESI^\dagger). Although, we are currently unable to unambiguously assign the location within the BoNT/A LC where Dyngo-4a resides, our data obtained in total suggest that Dyngo-4a imparts itself within one or both of the exosite regions of the BoNT/A LC.

Augmenting these kinetic studies we conducted an SAR examination of Dyngo-4a to probe what functional groups were essential for its activity (Scheme 1). A variety of analogues were either purchased or synthesized, the latter through a condensation between a hydrazide and an aldehyde under acidic conditions (Scheme 1), and tested in the 66mer assay. As expected, the three phenolic groups were found to be important for the inhibitory activity. Interestingly, 6, having an altered phenolic substitution pattern exhibited virtually equal potency and a competitive inhibition mechanism with $K_i = 0.46 \pm 0.08 \mu\text{M}$ (ESI^\dagger). Moreover, Dyngo-4a



Scheme 1 SAR study examining Dyngo-4a as the inhibitor-scaffold and BoNT/A protease. The top panel presents the general synthetic route used to prepare Dyngo-4a analogues. All assays were conducted as previously described.¹⁵

analogues presenting only one/two hydroxyl groups were inactive. Strikingly, although the hydroxyl functionality displayed within the naphthyl moiety was not crucial as a point of the inhibitory action, naphthyl scaffolding was required for inhibition (**12** & **13**, Scheme 1).

Having established that Dyngo-4a is a promising BoNT/A protease inhibitor, we evaluated the compound in a cellular assay using human induced pluripotent stem cell (hiPSC)-derived neurons.¹⁷ This platform was chosen due to the high sensitivity to the BoNT, a steep dose-response curve, and species-relevant assay with a more pure and defined population of neurons than other cell-based assays. To distinguish between its endocytic and protease inhibitory activities, the assay was conducted in two different modes. In one assay, Dyngo-4a was added 1 h before the toxin exposure to examine its endocytic inhibition; in the other, the inhibitor was added 45 min after the toxin exposure to assess its protease inhibition. Dyngo-4a showed SNAP-25 protection, albeit at a relatively high concentration of 200 μM when it was added before toxin exposure. Unfortunately, a similar inhibition was not observed when added after toxin exposure. Post-exposure cellular rescue using protease inhibitors has been rare.^{8,11} We also hypothesize that a lack of inhibition might also be due to our inability to examine higher concentrations of Dyngo-4a. This is especially pertinent with the sensitive nature of hiPSC neurons to foreign substances, which has been noted, and indeed Dyngo-4a showed cellular toxicity above 250 μM.

With kinetic, SAR, and cellular studies unfolded, we summoned a mouse lethality model, hoping to further probe the importance of non-endocytic efficacy engendered within Dyngo-4a. Thus, Dyngo-4a was injected 2.5–3 h post intoxication where in the mice begin to labor (pinched abdomens and labored breathing), and survival time was monitored and is summarized in Fig. 5. Remarkably, we observed that multiple rodents survived the challenge. Although other mechanisms that might come into play are currently unknown, clearly Dyngo-4a has an effect on survival that would not be governed in this time frame by dynamin inhibition of BoNT/A internalization.

In summary, Dyngo-4a, a heretofore inhibitor of BoNT/A neurotoxicity previously thought to function solely *via* the blocking of heavy chain internalization through the inhibition




Fig. 5 Mouse lethality study of Dyngo-4a protection.

of dynamin, has now been found to be also a regulator of the BoNT/A protease. This unrecognized BoNT/A LC inhibition by Dyngo-4a engenders this molecule to now be the first reported pharmacological antagonist that can blunt multiple processes associated with BoNT/A neurotoxicity. Kinetic examination of Dyngo-4a and the BoNT/A LC though complex and challenging posits Dyngo-4a's mode of inhibition through the protease's exosite. Although the results from a cell-based assay using hiPSC neurons suggest predominant endocytic inhibition, the survival of mice was observed following a lethal injection of BoNT/A, implicating protease inhibition by Dyngo-4a.

While unintended off-target activities can lead to toxicity and jeopardize drug discovery efforts, in recent years, the number of drugs targeting multiple targets has been increased, and a number of polypharmacological strategies have been put forth to address challenging goals, as represented by the success of multikinase inhibitors.¹⁸ Making a case for Dyngo-4a is tantalizing in that a single pharmacophore, a polyphenol, targets two completely different cellular proteins, one endogenous and the other exogenous. We believe that a rational dual inhibition design strategy targeting endocytosis and the BoNT/A LC has enormous potential to combat one of the most potent toxins known to man. We are hopeful that this newly found dual inhibitor activity will stimulate the discovery of optimized new generation endocytic-protease inhibitors of not only BoNT/A but also of the other BoNT serotypes.

We acknowledge the National Institute of Health (AI080671) for funding, and Dr. Michael C. Goodnough (MetabioLogics, Madison, WI) for conducting the BoNT/A mouse lethality assay. The authors thank Dr. Peter Šilhár for the synthesis of chicoric acid i-Pr ester. M.R. acknowledges the Fulbright Commission for the financial support. This is manuscript #29038 from The Scripps Research Institute.

Notes and references

- (a) C. Montecucco and G. Schiavo, *Q. Rev. Biophys.*, 1995, **28**, 423; (b) K. Oguma, Y. Fujinaga and K. Inoue, *Microbiol. Immunol.*, 1995, **39**, 161.
- L. L. Simpson, *Annu. Rev. Pharmacol.*, 2004, **44**, 167.
- N. Dover, J. R. Barash, K. K. Hill, G. Xie and S. S. Arnon, *J. Infect. Dis.*, 2014, **209**, 192.
- E. J. Schantz and E. A. Johnson, *Microbiol. Rev.*, 1992, **56**, 80.
- (a) S. L. Osborne, C. F. Latham, P. J. Wen, S. Cavaignac, J. Fanning, P. G. Foran and F. A. Meunier, *J. Neurosci. Res.*, 2007, **85**, 1149;

- (b) S. S. Arnon, R. Schechter, T. V. Inglesby, D. A. Henderson, J. G. Bartlett, M. S. Ascher, E. Eitzen, A. D. Fine, J. Hauer, M. Layton, S. Lillibridge, M. T. Osterholm, T. O'Toole, G. Parker, T. M. Perl, P. K. Russell, D. L. Swerdlow, K. Tonat and W. G. C. Biodefense, *JAMA, J. Am. Med. Assoc.*, 2001, **285**, 1059.
- 6 J. M. Rusnak and L. A. Smith, *Hum. Vaccines*, 2009, **5**, 794.
- 7 J. C. Burnett, E. A. Henchal, A. L. Schmaljohn and S. Bavari, *Nat. Rev. Drug Discovery*, 2005, **4**, 281.
- 8 B. Willis, L. M. Eubanks, T. J. Dickerson and K. D. Janda, *Angew. Chem., Int. Ed.*, 2008, **47**, 8360.
- 9 (a) G. E. Boldt, J. P. Kennedy and K. D. Janda, *Org. Lett.*, 2006, **8**, 1729; (b) P. Šilhár, N. R. Silvaggi, S. Pellett, K. Čapková, E. A. Johnson, K. N. Allen and K. D. Janda, *Bioorg. Med. Chem.*, 2013, **21**, 1344; (c) G. N. Stowe, P. Šilhár, M. S. Hixon, N. R. Silvaggi, K. N. Allen, S. T. Moe, A. R. Jacobson, J. T. Barbieri and K. D. Janda, *Org. Lett.*, 2010, **12**, 756; (d) N. R. Silvaggi, G. E. Boldt, M. S. Hixon, J. P. Kennedy, S. Tzipori, K. D. Janda and K. N. Allen, *Chem. Biol.*, 2007, **14**, 533.
- 10 (a) P. Šilhár, K. Čapková, N. T. Salzameda, J. T. Barbieri, M. S. Hixon and K. D. Janda, *J. Am. Chem. Soc.*, 2010, **132**, 2868; (b) L. M. Eubanks, P. Šilhár, N. T. Salzameda, J. S. Zakhari, X. C. Feng, J. T. Barbieri, C. B. Shoemaker, M. S. Hixon and K. D. Janda, *ACS Med. Chem. Lett.*, 2010, **1**, 268.
- 11 (a) P. Šilhár, L. M. Eubanks, H. Seki, S. Pellett, S. Javor, W. H. Tepp, E. A. Johnson and K. D. Janda, *J. Med. Chem.*, 2013, **56**, 7870; (b) H. Seki, S. Pellett, P. Šilhár, G. N. Stowe, B. Blanco, M. A. Lardy, E. A. Johnson and K. D. Janda, *Bioorg. Med. Chem.*, 2014, **22**, 1208.
- 12 C. B. Harper, S. Martin, T. H. Nguyen, S. J. Daniels, N. A. Lavidis, M. R. Popoff, G. Hadzic, A. Mariana, N. Chau, A. McCluskey, P. J. Robinson and F. A. Meunier, *J. Biol. Chem.*, 2011, **286**, 35966.
- 13 (a) M. T. Howes, M. Kirkham, J. Riches, K. Cortese, P. J. Walser, F. Simpson, M. M. Hill, A. Jones, R. Lundmark, M. R. Lindsay, D. J. Hernandez-Deviez, G. Hadzic, A. McCluskey, R. Bashir, L. Liu, P. Pilch, H. McMahon, P. J. Robinson, J. F. Hancock, S. Mayor and R. G. Parton, *J. Cell Biol.*, 2010, **190**, 675; (b) E. Macia, M. Ehrlich, R. Massol, E. Boucrot, C. Brunner and T. Kirchhausen, *Dev. Cell*, 2006, **10**, 839; (c) A. McCluskey, J. A. Daniel, G. Hadzic, N. Chau, E. L. Clayton, A. Mariana, A. Whiting, N. N. Gorgani, J. Lloyd, A. Quan, L. Moshkanbaryans, S. Krishnan, S. Perera, M. Chircop, L. von Kleist, A. B. McGeachie, M. T. Howes, R. G. Parton, M. Campbell, J. A. Sakoff, X. Wang, J. Y. Sun, M. J. Robertson, F. M. Deane, T. H. Nguyen, F. A. Meunier, M. A. Cousin and P. J. Robinson, *Traffic*, 2013, **14**, 1272; (d) S. Lee, K. Y. Jung, J. Park, J. H. Cho, Y. C. Kim and S. Chang, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 4858.
- 14 (a) H. T. McMahon and E. Boucrot, *Nat. Rev. Mol. Cell Biol.*, 2011, **12**, 517; (b) A. J. Newton, T. Kirchhausen and V. N. Murthy, *Proc. Natl. Acad. Sci. USA*, 2006, **103**, 17955.
- 15 (a) L. M. Eubanks, M. S. Hixon, W. Jin, S. W. Hong, C. M. Clancy, W. H. Tepp, M. R. Baldwin, C. J. Malizio, M. C. Goodnough, J. T. Barbieri, E. A. Johnson, D. L. Boger, T. J. Dickerson and K. D. Janda, *Proc. Natl. Acad. Sci. USA*, 2007, **104**, 6490; (b) K. Čapková, M. S. Hixon, L. A. McAllister and K. D. Janda, *Chem. Commun.*, 2008, 3525.
- 16 Attempted dual inhibition assay between Dyno-4a and chicoric acid were found to be complex and difficult to interpret due to the partial inhibition character of chicoric acid. On the other hand, a similar assay using chicoric acid i-Pr ester showed workable kinetic data. A future report detailing synthesis and kinetic analysis of a series of chicoric acid derivatives will be reported in due course.
- 17 R. C. M. Whitemarsh, M. J. Strathman, W. H. Tepp, C. Stankewicz, L. G. Chase, E. A. Johnson and S. Pellett, *Toxicon*, 2013, **68**, 78.
- 18 (a) J. U. Peters, *J. Med. Chem.*, 2013, **56**, 8955; (b) P. Ciceri, S. Muller, A. O'Mahony, O. Fedorov, P. Filippakopoulos, J. P. Hunt, E. A. Lasater, G. Pallares, S. Picaud, C. Wells, S. Martin, L. M. Wodicka, N. P. Shah, D. K. Treiber and S. Knapp, *Nat. Chem. Biol.*, 2014, **10**, 305.