

Green Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. T. S. Lew, D. S. W. Lim and Y. Zhang, *Green Chem.*, 2015, DOI: 10.1039/C5GC01374G.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/greenchem

Tedrick Thomas Salim Lew, Diane Shu Wen Lim and Yugen Zhang*

Green Chemistry

COMMUNICATION



Copper(I)-Catalyzed Amidation Reaction of Organoboronic Esters and Isocyanates

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A simple and efficient methodology for the preparation of amides from easily available organoboronic esters and isocyanates has been accomplished using ligand-free copper(I) catalyst. The reaction system demonstrated broad substrate scope and provided convenient access to a wide variety of secondary amides.

Amide bonds are one of the most important chemical building blocks in nature. Not only do they constitute the backbone of peptides and proteins, but they are also found extensively in synthetic polymers, fine chemicals and pharmaceutically active molecules.¹ Despite their ubiquity, most amides are prepared through the traditional approach of condensation reaction between carboxylic acids and amine in the presence of stoichiometric amounts of coupling reagents.^{2,3} It is therefore not surprising that a catalytic and more atom-economical approach for amide formation was recently highlighted as the top challenge in green chemistry research.⁴ Aside from economic and environmental concerns, the preparation of sterically-hindered and electron-deficient amides from carboxylic acid-amine condensation is also widely recognized to be a long-standing challenge.⁵

In recent years, there have been significant efforts to explore new reaction pathways for amide synthesis from different starting materials using transition metals as the catalysts.^{3,6-12} More recently, the use of organoboron reagents in transition metal-catalyzed addition reactions with isocyanates to form secondary amides has been developed.¹³ These reactions are particularly useful for the synthesis of various functionalized amides because of the wide availability of organoboron reagents with different functional groups.¹⁴ In addition, organoborons are non-toxic and highly stable.¹⁵⁻¹⁶ However, the activation of organoboron reagents in amidation reactions has thus far been the domain of precious rhodium^{13a} and palladium^{13b} catalysts, thereby limiting its

Institute of Bioengineering and Nanotechnology, 31 Biopolis Way, The Nanos, Singapore 138669. E-mail: ygzhang@ibn.a-star.edu.sg Electronic Supplementary Information (ESI) available: Details of experimental applications. Furthermore, in Rh or Pd catalyst systems organoboron homo-coupling side reaction is unavoidable, leading to the use of a large excess of organoboron reagents for the Rh system and low amide yield in Pd system.¹³ The preparation of sterically-hindered and electron-deficient amides in syntheticallyuseful yields is also difficult to achieve with these precious metal catalyst systems. On the other hand, the use of cost-effective first row transition metals such as copper as catalysts for amidation reactions has attracted great attention.¹⁷ Herein we report an efficient copper-catalyzed system for an amidation reaction between arylboronates and isocyanates. Some of the notable features of our approach include: 1) the use of cheap and earthabundant copper catalysts, 2) broad substrate scope of both arylboronates and isocyanates, which includes sterically encumbered and electron-deficient substrates, and 3) suppression of organoboron homo-coupling side reaction.

We began our investigation by using phenylboronic 1,3propanediol ester (1a) and 3,5-dimethylphenyl isocyanate (2a) as the model substrates for the amidation reaction. It was encouraging to find that in the presence of IPrCuCl (10 mol%), NaOt-Bu (1 equiv.) and DMF (2 mL) at 120 °C, the desired product (**3a**) was detected after 16 hours, albeit in a low yield (Table 1, entry 1). Two of the notable side products which were observed are amidine 4a and the carbamate formed from the reaction between 2a and NaOt-Bu. As the reaction time was extended to 48 h, the carbamate intermediate was consumed and this was accompanied by a substantial increase in the amide yield. When the reaction temperature was increased to 140 °C, the yield of amide 3a increased significantly (Table 1, entry 2) while there was only trace amount of carbamate detected. This suggests that the carbamat formation is reversible and that it decomposes at extended reaction time or higher temperature to give 2a and the base. However, at temperatures above 140 °C (Table 1, entry 3), isocyanate 2a decomposes through reaction with DMF.¹⁸ No phenylboronic ester homo-coupling side product (biphenyl) was observed in the reaction mixture.

Various copper catalysts were then screened. To our delight, ligand-free CuI catalyst was found to be the most efficient catalyst for this reaction and the formation of amidine **4a** was significantl suppressed (Table 1, entries 4 - 6). Other Cu(I) catalysts were found to be less effective than CuI, while the use of Cu(II) catalysts led to

procedures and product characterizations. See DOI: 10.1039/x0xx00000x

Journal Name

significant formation of undesirable biphenyl side product, thereby resulting in lower amide yields (Table 1, entries 5-10). Further optimization also revealed that LiOt-Bu gave the highest yield of **3a** among the different bases screened (Table 1, entries 6, 11-17). A slight excess of **1a** (1.5 equivalents) was found to further improve the yield of **3a** to 82% by compensating for the loss of arylboronate ester to background protodeboronation. The use of other organoboron reagents in place of **1a** gave the product in a lower yield under the same reaction conditions (Table S1 in the ESI). Both the Cu catalyst and the base were crucial for the formation of the amide as no product was formed in the absence of either reagent (Table 1, entries 19 and 20).

COMMUNICATION

Table 1 Optimization of reaction conditions for Cu-catalyzed amidation of
organoboronic ester and isocyanate

	+ Cu c base	atalyst		
1a	2a		3a	4a
Entry	Catalyst (mol%)	Base	<i>T</i> (°C)	Yield 3a (%) ^b
1	(IPr)CuCl (10)	NaOt-Bu	120	16
2	(IPr)CuCl (10)	NaOt-Bu	140	34
3	(IPr)CuCl (10)	NaOt-Bu	160	23
4	CuCl (10)	NaOt-Bu	140	40
5	CuCl (20)	NaOt-Bu	140	53
6	Cul (20)	NaOt-Bu	140	60
7	CuCN (20)	NaOt-Bu	140	52
8	CuOAc (20)	NaOt-Bu	140	34
9	CuCl ₂ (20)	NaOt-Bu	140	35
10	CuBr ₂ (20)	NaOt-Bu	140	15
11	Cul (20)	LiOt-Bu	140	64
12	Cul (20)	KOt-Bu	140	-
13	Cul (20)	LiOMe	140	45
14	Cul (20)	CsF	140	17
15	Cul (20)	KF	140	42
16	Cul (20)	Cs_2CO_3	140	-
17	Cul (20)	K_3PO_4	140	-
18 [°]	Cul (20)	LiOt-Bu	140	82
19	Cul (10)	LiOMe	140	55
20	Cul (20)	-	140	-
21	-	LiOt-Bu	140	-

^{*a*} Reaction conditions: **1a** (0.5 mmol), **2a** (0.5 mmol), base (0.5 mmol), DMF (2 mL), 16 h, Ar atmosphere, unless otherwise noted. ^{*b*} Yields were determined by ¹H NMR spectroscopy using 1,3,5-trimethoxybenzene as internal standard. ^{*c*} Reaction was carried out using 1.5 equivalents (0.75 mmol) of **1a**.

With the optimized reaction conditions, the scope of arylboronate esters was investigated (Table 2). The reaction proceeded well with sterically-hindered arylboronate esters, giving the corresponding amides in moderate to excellent yields (**3b-3c**). A range of arylboronate esters bearing electron-donating groups (e.g.

OMe), as well as electron-withdrawing substituents, (e.g., the lides, CN, CF₃), could provide the amide product the good states (**3d-3j**). However, yields were decreased for arylboronate esters with very strong withdrawing groups, such as nitro group (13% yield, SI). We were particularly pleased to find that **3i** could be efficiently prepared by this protocol as trifluoromethyl-substituted amides serve as important building blocks in the preparation of a wide variety of agrochemicals and pharmaceuticals.¹⁹ Importantly, the Cu(I)-catalyzed system was not restricted to aromatic substrates. Heteroaromatic substrates, such as pyridyl and thienyl derivatives, also furnished moderate yields of the desired amides (**3k-3I**). Although a lower yield of 40% was initially obtained from the alkenyl boronic ester (**3m**), extending the reaction time to 48 hours resulted in a modest yield of 56%.

 Table 2
 Substrate scope of arylboronate esters 1 for Cu-catalyzed ami

 dation reaction with isocyanates
 a,b



^a Conditions: 1 (0.75 mmol), 2a (0.5 mmol), Cul (20 mol%), LiOt-Bu (0.5 mmol), DMF (2 mL), 140 °C, 16 h, Ar atmosphere, unless otherwise noted.
 ^b Isolated yields. ^c 48 h reaction time.

With good reactivity for organoboronate esters in hand, the substrate scope of the isocyanate was then screened (Table 3). Sterically-hindered substrates and aryl isocyanates bearing an electron-donating group readily underwent amidation, resulting it excellent product yields (**3o-3p**). In the condensation reaction conventionally employed for amide synthesis, the preparation of amides derived from electron-deficient amines is often difficult because of the decreased nucleophilicity of the amines. On the other hand, isocyanates bearing electron-withdrawing halide groups were efficiently transformed into the desired amide products through this Cu(I)-catalyzed reaction system (**3q-3r**). Various functional groups such as cyano and nitro substituents were also well tolerated under the reaction conditions and the desired products were obtained in modest yields (**3s-3t**). Notably, the bulk adamantanyl isocyanate also proved to be an excellent substrate to

COMMUNICATION

Journal Name

provide the corresponding sterically-hindered amide in high yield (**3u**), highlighting the versatility and efficacy of Cu(I)-catalytic system for amide formation. In addition, gram scale synthesis of 3a was also carried and 78% of yield was obtained (SI).

Table 3 Substrate scope of isocyanates **2** for Cu-catalyzed amidation rea-ction with phenylboronic 1,3-propanediol ester



^a Conditions: 1a (0.75 mmol), 2 (0.5 mmol), Cul (20 mol%), LiOt-Bu (0.5 mmol), DMF (2 mL), 140 °C, 16 h, Ar atmosphere, unless otherwise noted.
 ^b Isolated yields.

The amidation reaction may follow a similar mechanism to that of the Cu-catalyzed carboxylation of organoboronic esters with CO_2 reported by Hou and co-workers,²⁰ with the major difference being isocyanate as the carbon source instead of CO_2 . The arylboronate ester will first undergo a transmetalation reaction with the alkoxide complex **5** formed by the metathesis reaction between CuI and LiOt-Bu (Scheme 1). The resulting organocopper complex would then participate in an addition reaction with the electrophilic isocyanate to form intermediate **6**, which subsequently undergoes metathesis reaction with LiOt-Bu to regenerate the complex **5**. Simultaneously, this metathesis reaction produces lithium salt **7** which, upon protonation, gives the desired amide product.



Scheme 1 Proposed catalytic cycle for amidation reaction between organo-boronic esters and isocyanates.

In summary, we have demonstrated an efficient coppercatalyzed system for the amidation reaction between widely available organoboronate esters and isocyanates_{view}Thacleboad substrate scope showcases the versatilityDahdobHtyCofCopper catalysts to mediate this type of synthetic transformation which has previously been achieved only by the use of precious metal catalysts. Furthermore, this approach also provides an alternative route to access amides which are difficult to synthesize through a conventional condensation reaction. This new method would be of significant use in the amide preparation for pharmaceuticals, agrochemicals and materials synthesis.

Acknowledgements

This work was supported by the Institute of Bioengineering and Nanotechnology (Biomedical Research Council, Agency for Science, Technology and Research (A*STAR), Singapore).

Notes and references

- (a) J. Zabicky, *The Chemistry of Amides*, Wiley-Interscience, New York, 1970; (b) A. Greenberg, C. M. Breneman and J. F Liebman, *The Amide Linkage: Structural Significance in Chemistry, Biochemistry and Materials Science*, John Wiley & Sons: New York, NY, 2000; (c) J. S. Carey, D. Laffan, C. Thomson and M. T. Williams, *Org. Biomol. Chem.*, 2006, 4, 2337-2347; (d) T. J. Deming, *Prog. Polym. Sci.*, 2007, 32, 858-875.
- 2 (a) E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, **38**, 606-631; (b) S. D. Roughley and A. M. Jordan, *J. Med. Chem.*, 2011, **54**, 3451-3479.
- 3 (a) C. L. Allen and J. M. J. Williams, *Chem. Soc. Rev.*, 2011, 40, 3405-3415. (b) R. M. Lanigan and T. D. Sheppard, *Eur. J. Org. Chem.*, 2013, 7453-7465; (c) H. Lundberg, F. Tinnis, N. Selander and H. Adolfsson, *Chem. Soc. Rev.*, 2014, 43, 2714 2742.
- D. J. C. Constable, P. J. Dunn, J. D. Hayler, G. R. Humphrey, J. L. Leazer, Jr., R. J. Linderman, K. Lorenz, J. Manley, B. A. Pearlman, A. Wells, A. Zaks and T. Y. Zhang, *Green Chem.*, 2007, 9, 411-420.
- 5 (a) V. R. Pattabiraman and J. W. Bode, *Nature*, 2011, 480, 471-479; (b) G. Schafer, C. Matthey and J. W. Bode, *Angew. Chem.*, *Int. Ed.*, 2012, 51, 9173-9175.
- 6 Representative examples for amidation of aldehydes: (a) A. Tillack, I. Rudloff and M. Beller, *Eur. J. Org. Chem.*, 2001, 523-528; (b) J. W. W. Chang and P. W. H. Chan, *Angew. Chem.*, *Int. Ed.*, 2008, **47**, 1138-1140; (c) S. Y. Seo, T. J. Marks, *Org. Lett.*, 2008, **10**, 317-319; (d) A. M. Whittaker, V. M. Dong, *Angew. Chem., Int. Ed.*, 2015, *54*, 1312-1315; (e) B. Zhou, J. Du, Y. Yang and Y. Li, *Org. Lett.*, 2013, **15**, 2934-2937; (f) W. J. Yoo, C.-J. Li, *J. Am. Chem. Soc.*, 2006, **128**, 13064-13065.
- 7 Representative examples for amidation of alcohols: (a) C. Gunanathan, Y. Ben-David, D. Milstein, *Science*, 2007, **317**, 790-792; (b) C. Chen, Y. Zhang and S. H. Hong, *J. Org. Chem.*, 2011, **76**, 10005-10010; (c) A. J. A. Watson, A. C. Maxwell and J. M. J. Williams, *Org. Lett.*, 2009, **11**, 2667-2670; (d) T. Zweifel, J.-V. Naubron and H. Gruetzmacher, *Angew. Chem.*, *Int. Ed.*, 2009, **48**, 559-563; (e) N. Ambreen, T. Wirth, *Euro. J. Org. Chem.*, 2014, **34**, 7590; (f) L. U. Nordstrom, H. Vogt, R. Madsen, *J. Am. Chem. Soc.*, 2008, **130**, 17672; (g) Y. Wang, D. Zhu, L. Tang, S. Wang, Z. Wang, *Angew. Chem.*, *Int. Ed.*, 2011 **50**, 8917-8921.

- COMMUNICATION
- 8 Representative examples for amidation of alkenes: (a) R. Shi, H. Zhang, L. Lu, P. Gan, Y. Sha, H. Zhang, Q. Liu, M. Beller and A. Lei, *Chem. Commun.*, 2015, **51**, 3247-3250; (b) X. H. Yang, K. Li, R. J. Song and J. H. Li, *Eur. J. Org. Chem.*, 2014, 616-623; (c) Z. M. Deng, J. L. Wei, L. H. Liao, H. Y. Huang, X. D. Zhao, *Org. Lett.*, 2015, **17**, 1834-1837; (d) K. Shin, H. Kim, S. Chang, *Acc. Chem. Res.*, 2015, **48**, 1040-1052.
- 9 Representative examples for amidation of alkynes: (a) Y. Uenoyama, T. Fukuyama, O. Nobuta, H. Matsubara and I. Ryu, Angew. Chem., Int. Ed., 2005, 44, 1075-1078; (b) J. H. Park, S. Y. Kim, S. M. Kim and Y. K. Chung, Org. Lett., 2007, 9, 2465-2468; (c) M. Kumar, S. Devari, A. Kumar, S. Sultan, Q. N. Ahmed, M. Rizvi, B. A. Shah, Asian J. Org. Chem., 2015, 4, 438-441; (d) T. Hamada, X. Ye and S. S. Stahl, J. Am. Chem. Soc., 2008, 130, 833-835; (e) H. Liu, G. P. S. Lau and P. J. Dyson, J. Org. Chem., 2015, 80, 386-391.
- Representative examples for amidation of aryl halides: (a) J.
 R. Martinelli, T. P. Clark, D. A. Watson, R. H. Munday and S. L. Buchwald, Angew. Chem., Int. Ed., 2007, 46, 8460-8463; (b)
 S. Kathiravan, S. Ghosh, G. Hogarth, Ian A. Nicholls, Chem. Commun., 2015, 51, 4834-4837; (c) X.-F. Wu, H. Neumann and M. Beller, Chem.-Asian J., 2010, 5, 2168-2172; (d) N. Saito, T. Taniguchi, N. Hoshiya, S. Shuto, M. Arisawa and Y. Sato, Green Chem., 2015, 17, 2358-2361.
- Representative examples for amidation of nitriles: (a) E. Tomas-Mendivil, F. J. Suarez, J. Diez and V. Cadierno, *Chem. Commun.*, 2014, **50**, 9661-9664. (b) R. S. Ramon, N. Marion and S. P. Nolan, *Chem. Eur. J.*, 2009, **15**, 8695-8697; (c) K. Yasuda, Y. Obora, *J. Organomet. Chem.*, 2015, **775**, 33-38; (d) Y.-M. Liu, L. He, M.-M. Wang, Y. Cao, H.-Y. He and K.-N. Fan, *ChemSusChem*, 2012, **5**, 1392-1396.
- 12 Representative examples for amidation of isocyanates: (a) X.-Y. Shi, K.-Y. Liu, J. Fan, X.-F. Dong, J.-F. Wei and C.-J. Li, *Chem.-Eur. J.*, 2015, **21**, 1900-1903; (b) X.-Y. Shi, A. Renzetti, S. Kundu and C.-J. Li, *Adv. Synth. Catal.*, 2014, **356**, 723-728. (c) K. D. Hesp, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2011, **133**, 11430-11433; (d) Y. Kuninobu, Y. Tokunaga, A. Kawata and K. Takai, *J. Am. Chem. Soc.*, 2006, **128**, 202-209; (e) K. Muralirajan, K. Parthasarathy and C.-H. Cheng, *Org. Lett.*, 2012, **14**, 4262-4265; (f) J. R. Hummel and J. A. Ellman, *Org. Lett.*, 2015, **17**, 2400–2403; (g) J. Li and L. Ackermann, *Angew. Chem., Int. Ed.*, 2015, **54**, 1-5; (h) C. Gurtler, K. Danielmeier, *Tetrahedron Lett.*, 2004, **45**, 2515-2521.
- 13 (a) T. Miura, Y. Takahashi and M. Murakami, *Chem. Commun.*, 2007, 3577-3579; (b) E. Kianmehr, A. Rajabi and M. Ghanbari, *Tetrahedron Lett.*, 2009, **50**, 1687-1688.
- Selected examples of organoboron preparations without Grignard or organolithium reagents: (a) T. Ishiyama, M. Murata and N. Miyaura, J. Org. Chem., 1995, 60, 7508-7510; (b) G. A. Molander, S. L. J. Trice and S. D. Dreher, J. Am. Chem. Soc., 2010, 132, 17701-17703; (c) J. Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka and M. R. Smith, Science, 2002, 295, 305-308; (d) S. Shimada, A. S. Batsanov, J. A. K. Howard and T. B. Marder, Angew. Chem., Int. Ed., 2001, 40, 2168-2171.
- 15 D. G. Hall, Boronic Acids, Wiley-VCH, Weinheim, 2005.
- 16 For reviews see: (a) T. Hayashi and K. Yamasaki, *Chem. Rev.*, 2003, **103**, 2829-2844; (b) T. Miura and M. Murakami, *Chem. Commun.*, 2007, 217-224.
- (a) C. Zhang, N. Jiao, J. Am. Chem. Soc., 2010, 132, 28-29; (b)
 Q. Kang, Y. Zhang, Green Chem., 2012, 14, 1016-1019; (c) S.

C. Ghosh, J. S. Y. Ngiam, A. M. Seayad, D. T. Tuan, C. J. L. Chai, A. Chen, J. Org. Chem., 2012, **72**, 8007.680155 (d) 1874D3 Grigg, J. W. Rigoli, R. V. Hoveln, S. Neale, J. M. Schomaker, Chem. Eur. J., 2012, **18**, 9391-9396.

- 18 M. L. Weiner, J. Org. Chem., 1960, 25, 2245-2247.
- 19 (a) F. Leroux, P. Jeschke and M. Schlosser, *Chem. Rev.*, 2005,
 105, 827-856; (b) K. Mueller, C. Faeh and F. Diederich, *Science*, 2007, 317, 1881-1886; (c) M. Shimizu and T. Hiyama, *Angew. Chem., Int. Ed.*, 2005, 44, 214-231; (d) P. Jeschke, *ChemBioChem*, 2004, 5, 570-589
- 20 T. Ohishi, M. Nishiura and Z. Hou, *Angew. Chem., Int. Ed.,* 2008, **47**, 5792-5795.

^{4 |} Green Chem., 2015, 00, 1-3



A simple and efficient methodology for the preparation of amides from easily available organoboronic esters and isocyanates with ligand-free copper(I) catalyst.