Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: J. Quibell, G. Perry, D. Cannas and I. Larrosa, *Chem. Sci.*, 2018, DOI: 10.1039/C8SC01016A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/chemical-science

Dpen Access Article. Published on 26 March 2018. Downloaded on 26/03/2018 07:18:35.

Journal Name

ARTICLE

Received 00th January 20xx, Accepted 00th January 20xx Jacob M. Quibell,^a† Gregory J. P. Perry^{a,b}†, D. M. Cannas^a and Igor Larrosa^{*,a}

DOI: 10.1039/x0xx00000x

www.rsc.org/

Methods for the conversion of *aliphatic* acids to alkyl halides have progressed significantly over the past century, however, the analogous decarboxylative bromination of *aromatic* acids has remained a longstanding challenge. The development of efficient methods for the synthesis of aryl bromides is of great importance as they are versatile reagents in synthesis and are present in many functional molecules. Herein we report a transition metal-free decarboxylative bromination of aromatic acids. The reaction is applicable to many electron-rich aromatic and heteroaromatic acids which have previously proved poor substrates for Hunsdiecker-type reactions. In addition, our preliminary mechanistic study suggests that radical intermediates are not involved in this reaction, which is in contrast to classical Hunsdiecker-type reactivity. Overall, the process demonstrates a useful method for producing valuable reagents from inexpensive and abundant starting materials.

Transition-Metal-Free Decarboxylative Bromination of Aromatic

Introduction

Aryl bromides are the substrate of choice when performing transition metal-catalysed cross-coupling reactions¹ or preparing Grignard and organolithium reagents.² They are also used in a variety of other transformations, such as nucleophilic substitution and HalEx reactions,³ and are the core structures in many natural products and dyes.⁴ Consequently, developing efficient methods for the synthesis of aryl bromides remains an important objective.⁵ The ability to directly substitute a carboxyl group with a bromo group has interested the synthetic community for many years. This transformation was first demonstrated by Borodine over a century ago, but came to bear the name "The Hunsdiecker Reaction" during the 1940's and is now a fundamental reaction in organic synthesis.^{6,7} The reaction involves the mixing of an aliphatic carboxylic acid with bromine in the presence of a silver salt to produce the desired alkyl halide. Various developments in this area were made during the latter half of the 20th century,⁸ however, the applicability of all these methods was limited due to the requirement of stoichiometric transition metal salts and/or poor generality. It is only recently that significant progress has been made in the decarboxylative bromination of aliphatic acids (Scheme 1A). Namely, the groups of Glorius⁹ and Li¹⁰ have demonstrated that primary, secondary and tertiary carboxylic acids can be converted into the

Carboxylic Acids

corresponding alkyl bromides in the presence of either an iridium or silver catalyst.

Conversely, the decarboxylative bromination of aromatic acids has suffered from a slower pace of development (Scheme 1B). Early findings demonstrated that the bromination of aromatic acids could indeed proceed, however, the reaction with electron-rich substrates was unselective and electron-deficient aromatics gave varying yields (Scheme 2).¹¹ Many have looked to solve these issues, but current procedures still suffer as they (a) require stoichiometric transition metals, (b) are of poor generality, and/or (c) are unselective.¹² This is disappointing as decarboxylative bromination holds potential as an economic route for aryl halide formation;¹³ benzoic acids are inexpensive and abundant, and the bromide group is a handle for selective transformations. We were eager to reinvestigate the aromatic Hunsdiecker reaction in order to establish a more efficient and selective route for aryl bromide formation. Here we reveal the development of a transition metal-free decarboxylative bromination that is applicable to a variety of electron-rich aromatic acids. In addition, preliminary investigations begin to shed light on the mechanism of this reaction.

Results and discussion

We have recently reported a transition-metal-free decarboxylative iodination of aromatic acids.¹⁴ The success of this procedure lies in the ability to prepare a variety of aryl iodides, simply by heating a benzoic acid in the presence of I₂. We therefore questioned whether a similarly efficient and low-cost decarboxylative bromination could be developed. We began our investigation by exposing the benzoic acid **1a** to our previous conditions, but switching the halogen source from I₂ to Br₂ (Table 1). Unfortunately, this resulted in the undesired

^{a.} School of Chemistry, University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom. Email: igor.larrosa@manchester.ac.uk

^{b.} Institute of Transformative Bio-Molecules (WPI-ITbM) and Graduate School of Science, Nagoya University, Chikusa, Nagoya 464-8602, Japan.

⁺ G.J.P.P and J.M.Q contributed equally to this work.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here].

Dpen Access Article. Published on 26 March 2018. Downloaded on 26/03/2018 07:18:35.

DOI: 10.1039/C8SC01016A Journal Name



Scheme 1: Timeline of the contrasting progress in A) aliphatic vs B) aromatic decarboxylative bromination.



Scheme 2: Current status of the aromatic Hunsdiecker reaction.

formation of the brominated acid 1a' and dibrominated product 2a' and none of the desired product 2a (Table 1, Entry 1). This reactivity is comparable with the aromatic Hunsdiecker reaction shown in Scheme 2 and demonstrates the challenges for achieving a selective bromination. By lowering the equivalents of Br₂ the selectivity could be improved, however, a large amount of the brominated acid 2a' was still produced (Entry 2). We then investigated less electrophilic bromine sources, such as NBS (N-bromosuccinimide) and DBH (1,3dibromo-5,5-dimethylhydantoin), however a mixture of products was still obtained and the desired product was formed in low yield (Entries 3 and 4). We then turned to the use of tribromide reagents as bromine sources for this transformation. Although pyridinium tribromide performed poorly in this reaction (entry 5), we found that tetraalkyl ammonium tribromide salts, $N(Me_4)Br_3$ and $N(^nBu_4)Br_3$, displayed good reactivity and high selectivity for the desired decarboxylative bromination (entries 6 and 7). $N(^{n}Bu_{4})Br_{3}$ was chosen as the brominating reagent of choice, allowing the product to be isolated in 90% yield. Further control experiments revealed that the reaction does not proceed in the absence of a base (Entry 8), but that performing the reaction in the dark or adding one equivalent of water had little effect on the reaction (Entries 9 and 10).¹⁵

Having demonstrated an efficient transition metal-free decarboxylative bromination, we then turned to exploring the scope of this reaction (Scheme 3). Previously, the decarboxylative bromination of 4-methoxybenzoic acid under Hunsdiecker-type conditions resulted in a mixture of products (Scheme 2), therefore we were impressed to observe the formation of the aryl bromide **2b** in high yield and high selectivity. This clearly demonstrates the advantages of our procedure over previous techniques. The desired product, **2c**, was not observed when using 3-methoxybenzoic acid,

suggesting the position of decarboxylation must be sufficiently nucleophilic for the decarboxylative bromination to occur. Other highly electron-rich substrates (**1d-1h**) could also undergo the desired transformation, including non-*ortho*substituted benzoic acids (**1b**, **1h**), which are generally unreactive in transition metal-mediated decarboxylative functionalisations. The procedure can also be applied on a large scale, thus, 5.5 g of the brominated product **2e** was prepared using the standard conditions on the bench top at room temperature, without requiring column

Table	1.	Optimisation	of	the	Transition	Metal	Free
Decarb	оху	lative Brominat	ion.				

ecarboxylative Bromination.								
OMe			OM	le	OMe			
1a	CO ₂ H K + [Br] M (1.0 equiv) 1(₃ PO ₄ (1.0 equ eCN (0.2 M) 00 °C , 4 h		Br +	Br			
				1a', 2a',	R = CO ₂ H R = Br			
Entry	[Br]	1a	2a	1a'	2a'			
1^b	Br ₂	trace	0	23	72			
2	Br ₂	8	26	56	3			
3	NBS ^c	58	21	9	trace			
4	DBH^{d}	6	40	32	7			
5	PyHBr ₃ ^e	39	6	54	0			
6	N(Me ₄)Br ₃	6	85	5	2			
7	N(ⁿ Bu ₄)Br ₃	5	91 (90) ^f	1	trace			
8 ^{<i>g</i>}	N(ⁿ Bu ₄)Br ₃	100	0	0	0			
9 ^{<i>h</i>}	N(ⁿ Bu ₄)Br ₃	11	84	2	trace			
10 ⁱ	N([″] Bu₄)Br₃	10	85	trace	3			

^a Reaction conditions: **1a** (0.2 mmol), [Br] (0.2 mmol, 1.0 equiv), K₃PO₄ (0.2 mmol, 1.0 equiv), MeCN (1.0 mL), 100 °C, 4 h. ^b Br₂ (0.6 mmol, 3.0 equiv). ^c *N*-bromosuccinimide. ^d 1,3-dibromo-5,5dimethylhydantoin. ^e pyridinium tribromide ^f Yield in parenthesis is of isolated material. Isolated as a mixture with **2a' (2a:2a'**, >150:1). ^g No K₃PO₄ added. ^h Performed in the dark. ⁱ 1.0 equiv H₂O added.

Dpen Access Article. Published on 26 March 2018. Downloaded on 26/03/2018 07:18:35.



Scheme 3: Scope of the decarboxylative bromination of aromatic acids. Reactions carried out on a 0.5 mmol scale. ^a Ratios in brackets indicate mono:dibrominated material by NMR analysis before separation. Asterisk indicates position of dibromination. ^b Isolated as mixture. ^c Room temperature. ^d Yields determined by NMR analysis. ^e N(ⁿBu₄)Br₃ (4.0 equiv).

chromatography. Polymethylated benzoic acids are poorly reactive substrates in transition metal-mediated decarboxylations, but they, and even simple toluic acid, showed good reactivity under our conditions (2i-2i). The procedure could also be applied to napthoic acids, despite a slight loss in selectivity (2m, 2n). The position of dibromination is indicated in both Schemes 3 and 4 by a red asterisk. A range of halogenated and trifluoromethylated benzoic acids were successfully decarboxylated, however, the presence of a methoxy group was necessary to maintain efficient reactivity (**2o-2y**). Benzoic acids that do not bear electron-donating substituents were unreactive under these conditions (**2z-2ac**). In light of this, we were surprised to observe good reactivity with electron-poor polyfluorinated benzoic acids (**2ad-2af**). This goes against the general trend of reactivity in this reaction and we are currently investigating the cause of this unique behaviour.

The procedure could also be applied to a range of heteroaromatic acids (Scheme 4). The bromination of heteroaromatic acids is highly limited as elevated temperatures (>160 °C) and stoichiometric transition metals are generally necessary.¹⁶ Our conditions were applicable to indoles (4a), benzothiophenes (4b, 4c) and benzofurans (4d). Unfortunately, under standard reaction conditions benzothiophene-(4e) and, to an extent, benzofuran-2carboxylic acid (4f) afforded undesirable levels of dihalogenation. This side-reaction, namely the β -bromination of benzo-fused five-membered heterocycles, is well-known to proceed readily at room-temperature with a variety of brominating agents; thus representing a limitation of our methodology. A range of 5-membered heterocycles (4g-4j) as well as pyridine (4k), chromone (4l) and cinnamic acid (4m) derivatives all underwent the desired decarboxylative bromination selectively.



Scheme 4: Scope of the decarboxylative bromination of heteroaromatic acids. Reactions carried out on a 0.5 mmol scale ^a Ratios in brackets indicate mono:dibrominated material by NMR analysis before separation. Asterisk indicates position of dibromination. ^b Isolated as mixture. ^c 50 °C.

Having established an efficient protocol for the decarboxylative bromination of aromatic acids, we began a preliminary mechanistic investigation. Our initial experiment involved the use of the oxyallyl-substituted benzoic acid **1A** as a radical clock (Figure 1a). The formation of cyclised products, via attack of an aryl radical on the pendent allyl chain of this

ARTICLE

compound, is an extremely fast process (k = $8 \times 10^9 \text{ s}^{-1}$),¹⁷ therefore, if cyclised products were to be observed in this reaction a radical mechanism could be suggested. Upon exposing oxyallyl-substituted benzoic acid 1A to our standard reaction conditions we only observed the formation of the aryl bromide 2A and none of the cyclised product. In light of this, we can suggest that either the reaction does not proceed through a radical mechanism or, if radicals are involved, then the rate at which the product is formed from the radical intermediate is an exceptionally fast process. This is an interesting observation as similar experiments that have previously been conducted on Hunsdiecker-type decarboxylations of aliphatic carboxylic acids have strongly supported a radical mechanism.¹⁸ Likewise, previous Hunsdiecker-type decarboxylations of aromatic acids have also been proposed to proceed through radical intermediates.^{12k-I}, Overall, although the above result does not definitively rule out a radical mechanism, it calls for a more thorough evaluation of Hunsdiecker-type reactivity.

Our previous work on decarboxylative iodinations established a concerted decarboxylation-iodination process, via a 4-membered transition state, as a possible non-radical pathway for decarboxylative halogenations.¹⁴ Following a similar protocol (Figure 1b), our DFT study found a pathway for decarboxylative bromination proceeding through an analogous concerted decarboxylation-bromination transition state. TS-II. Thus, our current mechanistic hypothesis is as follows: the benzoic acid is initially transformed into the hypobromite species I upon exposure to K_3PO_4 and $N(^nBu_4)Br_3$.¹⁹ The hypobromite then undergoes decarboxylation via a 4membered transition state (TS-II), to provide the product 2e with concomitant loss of CO_2 . The barrier for this transformation was calculated to be 19.2 kcal mol⁻¹, which is consistent with a process that proceeds at room temperature.20

We further probed the mechanism of this reaction by conducting ${}^{13}C/{}^{12}C$ KIE experiments. Heavy atom isotope effects have been widely used as a means to study decarboxylation events in chemical and enzymatic processes.²¹ The KIE at the C1 position for decarboxylation processes can be accurately determined at natural abundance by measuring the isotopic composition of the evolved CO2 through massspectroscopy technique.²² Unfortunately, with this method no information is gained on the other carbon atoms. Exploiting quantitative ¹³C-NMR, Singleton and co-workers have devised a useful procedure for determining intermolecular competitive ¹³C/¹²C KIEs at all positions at natural abundance.²³ Over the course of the reaction, the starting material is progressively enriched in the slowest reacting isotopologues. By evaluating the isotopic composition in the starting material before and after the reaction the KIEs can be determined. This has proved a powerful tool in elucidating reaction mechanisms and we were eager to test its value on our system. Two independent experiments were performed on 2,6-dimethoxybenzoic acid (1e) under standard reaction conditions at 30 °C for 70 minutes (Figure 1c). Remarkably, a primary KIE was observed at both C1 and C2 positions: KIEs of 1.014 \pm 0.004 and 1.009 \pm 0.006 were obtained for C1, while larger KIEs of 1.028 ± 0.004

Figure 1. Mechanistic investigations.





b) DFT study of the proposed path



c) Experimental and calculated ¹³C-KIEs

OMe O	C1	C2	C3	C4	C5
5 3 OMe	1.014(4)	1.028(4)	0.996(2)	1.000*	1.001(5)
	1.009(6)	1.025(5)	0.992(2)	1.000*	0.997(4)
	<mark>1.013</mark>	1.023	<mark>0.997</mark>	1.000	<mark>1.001</mark>

a) Standard reaction conditions with benzoic acid **1A** at 50 °C for 30 min. b) Energies calculated in acetonitrile (B3LYP-D3BJ/6-31+G(d)).²⁴ Gibbs free energies (G) in kcal mol⁻¹. c) Experimental KIEs (in black), the uncertainty on the last figure is reported in brackets. For C4 a KIE of 1.000 was assumed.²⁵ Calculated KIEs (in red) for the proposed path.²⁶

and 1.025 ± 0.005 were measured for C2. This values are consistent with the proposed pathway where a concerted decarboxylation-bromination transition state is involved in the product determing step (Figure 1b). The lower KIE for C1 in comparison to C2 suggests either an early transition state,²⁷ or that another kinetically relevant step is occuring prior to the product determing step.Error! Bookmark not defined.^e Examination of the reaction path by DFT (TS-II to 2e) revealed an early formation of C1-Br bond, resulting in an "hidden" Wheeland intermediate.²⁸ Extrusion of CO₂ from this transient species took place late along the reaction coordinates, thus in agreement with the experimental observations.²⁹ To further probe our mechanistic hypothesis, the KIE values for the proposed path were calculated (Figure 1c). Computed and experimental values were found to be in excellent agreement, lending strong support to the proposed concerted decarboxylation-bromination pathway.

We have conducted a preliminary mechanistic study of the developed decarboxylative bromination of aromatic acids. At present, we have strong evidence that excludes the intermediacy of aryl radicals (Figure 1a). By measuring the ¹³C/¹²C KIEs and performing DFT calculations we identified a concerted decarboxylation-bromination as possible pathway for this transformation (Figure 1b-c). This represents an alternative mechanism for hunsdiecker-type reactivity, as radicals are usually considered key intermediates in similar processes. While further investigations are necessary to better

Journal Name

establish the mechanism of this reaction, we believe that these initial studies highlight previously unrealised features of our system. We hope that these results inspire future studies that may greatly impact this and related procedures, and lead to the development of more efficient decarboxylative technologies.

Conclusions

Due to slow progress and issues with selectivity, the utility of the aromatic Hunsdiecker reaction has previously failed to be fully realised. In this report we have detailed the successful development of a high yielding aromatic Hunsdiecker-type reaction. This has led to the development of a decarboxylative bromination of electron-rich aromatic acids using low-cost and abundant starting materials. The avoidance of transition metals and the ability to scale-up the reaction make the process attractive for its simplicity and low cost. The Hunsdiecker reaction is commonly proposed to proceed via a radical pathway, however, our combined experimental and theoretical mechanistic study has suggested an alternative mechanism that does not involve radical intermediates. These results directly challenge a long-held view of Hunsdiecker-type reactivity. Further studies are necessary, but we hope that future investigations will better elucidate this mechanism. Overall, we believe that this report demonstrates the potential of decarboxylative halogenation as an efficient route to valueadded chemical commodities.

Conflicts of interest

"There are no conflicts to declare".

Acknowledgements

We gratefully acknowledge the Engineering and Physical Sciences Research Council for funding (EP/L014017/2 and EP/K039547/1). We thank Dr. Ralph Adams (UoM) for the quantitative NMR experiments.

References

1996. (d) N. Sotomayor and E. Lete, *Curr. Org. Chem.* 2003, **7**, 275–300.

³ For nucleophilic substitution see (a) J. F. Burnett and R. E. Zahler, *Chem. Rev.* 1951, **49**, 273–412. (b) J. F. Hartwig, *Synlett* 2006, 1283– 1294. *Halogen Exchange* see: (c) A. Klapars and S. L. Buchwald, *J. Am. Chem. Soc.* 2002, **124**, 14844–14845. d) M. Chen, S. Ichikawa and S. L. Buchwald, *Angew. Chem. Int. Ed.* 2015, **54**, 263–266. e) T. D. Sheppard, *Org. Biomol. Chem.* 2009, **7**, 1043–1052. *aryl radical formation* see (f) W. P. Neumann, *Synthesis* 1987, 665–683. (g) H. Kim and C. Lee, *Angew. Chem. Int. Ed.* 2012, **51**, 12303–12306. (h) I. Ghosh, T. Ghosh, J. I. Bardagi and B. Konig, *Science* 2014, **346**, 725– 728. *aryne generation* see (i) H. Pellissier and M. Santelli, *Tetrahedron* 2003, **59**, 701–730. (j) R. A. Moss, M. S. Platz and M. Jones, Jr., In *Reactive Intermediate Chemistry*; Winkler, M.; Wenk, H. H.; Sander, W., Eds.; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2005; pp 741–794.

⁴ (a) A. Butler and J. V. Walker, *Chem. Rev.* 1993, **93**, 1937–1944. (b)
 J. L. Wolk and A. A. Frimer, *Molecules* 2010, **15**, 5473–5580. (c) J. L.
 Wolk and A. A. Frimer, *Molecules* 2010, **15**, 5561–5580. (d) G. W.
 Gribble, *J. Chem. Educ.* 2004, **81**, 1441–1449.

⁵ For selected approaches for the preparation of aryl bromides see Ref 3e and (a) R. Taylor, *Electrophilic Aromatic Substitution*; John Wiley: New York, 1990. (b) V. Snieckus, *Chem. Rev.* 1990, **90**, 879– 933. (c) A. Roglans, A. Pla-Quintana and M. Moreno-Mañas, *Chem. Rev.* 2006, **106**, 4622–4643. (d) D. Kalyani, A. R. Dick, W. Q. Anani and M. A. Sanford, *Org. Lett.* 2006, **8**, 2523–2526. (e) C. J. Teskey, A. Y. W. Lui and M. F. Greaney, *Angew. Chem. Int. Ed.* 2015, **54**, 11677–11680.

^b (a) A. Borodine, *Justus Liebigs Ann. Chem.* 1861, **119**, 121–123. (b) H. Hunsdiecker and C. Hunsdiecker, *Ber. Dtsch. Chem. Ges.* 1942, **75B**, 291–297. (c) C. Wilson, in *Org. React.*, John Wiley & Sons, Inc., Hoboken, NJ, USA, 2011, pp. 332–387.

⁷ (a) D. Crich, In *Comprehensive Organic Synthesis*; B. M. Trost, I. Fleming, Eds.; Pergamon Press, 1991; Vol. 7, pp 717–734. (b) D. Crich and K. Sasaki, in *Comprehensive Organic Synthesis II*. P. Knochel, G. A. Molander, Eds.; Elsevier, 2014, pp. 818–836. (c) J. J. Li, In *Name Reactions. A Collection of Detailed Reaction Mechanisms, 3rd Edition.*; Springer-Verlag Berlin Heidelberg, 2006, pp. 310–311. d) L. Kürti and B. Czakó, In *Strategic Applications of Named Reactions in Organic Synthesis*; Elsevier, California, SD, USA, 2005, pp. 218–219.

⁸ (a) S. Cristol and W. J. Firth, Jr., J. Org. Chem. 1961, 26, 280. (b) J.
K. Kochi, J. Am. Chem. Soc. 1965, 87, 2500–2502. (c) D. H. R. Barton,
H. P. Faro, E. P. Serebryakov and N. F. Woolsey, J. Chem. Soc. 1965,
2438–2444. (d D. H. R. Barton, D. Crich and W. B. Motherwell,
Tetrahedron Lett. 1983, 24, 4979–4982. (e) D. H. R. Barton, D. Crich and W. B. Motherwell, J. Chem. Soc., Chem. Commun. 1983, 939–
941. (f) J. I. Concepcion, C. G. Francisco, R. Freire, R. Hernandez, J. A.
Salazar and E. Suarez, J. Org. Chem. 1986, 51, 402–404. (g) P.
Camps, A. E. Lukach, X. Pujol and S. Vázquez, Tetrahedron 2000, 56, 2703–2707.

⁹ L. Candish, E. A. Standley, A. Gómez-Suárez, S. Mukherjee and F. Glorius, *Chem. Eur. J.* 2016, **22**, 9971–9974.

¹⁰ X. Tan, T. Song, Z. Wang, H. Chen, L Cui and C. Li, *Org. Lett.* 2017, 19, 1634–1637.

¹¹ (a) W. G. Dauben and H. Tilles, *J. Am. Chem. Soc.* 1950, **72**, 3185–3187. (b) R. A. Barnes and R. J. Prochaska, *J. Am. Chem. Soc.* 1950, **72**, 3188–3191. (c) J. W. H. Oldham, *J. Chem. Soc.* 1950, 100–108.

² For reports that require stoichiometric transition metals see

¹ J. Hassan, M. Sévignon, C. Gozzi, E. Schulz and M. Lemaire, *Chem. Rev.* 2002, **102**, 1359–1470. (b) C. C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot and V. Snieckus, *Angew. Chem. Int. Ed.* 2012, **51**, 5062–5085. (c) C. Bolm, J. P. Hildebrand, K. Muniz and N. Hermanns, *Angew. Chem. Int. Ed.* 2001, **40**, 3284–3308.

² (a) P. Knochel, W. Dohle, N. Gommermann, F. F. Kneisel, F. Kopp, T. Korn, I. Sapountzis and V. A. Vu, *Angew. Chem. Int. Ed.* 2003, **42**, 4302–4320. (b) D. Tilly, F. Chevallier, F. Mongin and P. C. Gros, *Chem. Rev.* 2014, **114**, 1207–1257. (c) *Handbook of Grignard Reagents*; G. S. Silverman, P. E. Rakita, Eds.; Dekker: New York, NY,

Dpen Access Article. Published on 26 March 2018. Downloaded on 26/03/2018 07:18:35.

Journal Name

reference 11 and: (a) J. S. Kiely, L. L. Nelson and P. J. Boudjouk, Org. Chem. 1977, 42, 1480. (b) S. Uemura, S. Tanaka, M. Okano and M. Hamana, J. Org. Chem. 1983, 48, 3297-3301. (c) Y. Luo, X. Pan and J. Wu, Tetrahedron Lett. 2010, 51, 6646-6648. (d) J. Cornella, M. Rosillo-Lopez and I. Larrosa, Adv. Synth. Catal. 2011, 353, 1359-1366. (e) X. Peng, X.-F. Shao and Z.-Q. Liu, Tetrahedron Lett. 2013, 54, 3079–3081. (f) Z. Fu, Z. Li, Y. Song, R. Yang, Y. Liu and H. Cai, J. Org. Chem. 2016, 81, 2794-2803. For reports that show limited scope and/or poor selectivity see : (g) D. H. R. Barton, B. Lacher and S. Z. Zard, Tetrahedron Lett. 1985, 26, 5939-5942. (h) D. H. R. Barton, B. Lacher and S. Z. Zard, Tetrahedron 1987, 43, 4321-4328. (i) B.-S. Koo, E.-H. Kim and K.-J. Lee, Synth. Commun. 2002, 32, 2275–2286. (j) H. Hamamoto, S. Hattori, K. Takemaru and Y. Miki, Synlett 2011, 2011, 1563-1566. (k) Z. Li, K. Wang and Z.-Q. Liu, Synlett 2014, 25, 2508–2512. (I) J. Fang, D. Wang, G.-J. Deng and H. Gong, Tetrahedron Lett. 2017, 58, 4503-4506.

¹³ For selected reviews on decarboxylative activation see: (a) L. J. Gooßen, K. Gooßen, N. Rodríguez, M. Blanchot, C. Linder and B. Zimmermann, *Pure Appl. Chem.* 2008, **80**, 1725–1733. (b) L. J. Gooßen, N. Rodríguez and K. Gooßen, *Angew. Chem. Int. Ed.* 2008, **47**, 3100–3120. (c) L. J. Gooßen, F. Collet and K. Gooßen, *Isr. J. Chem.* 2010, **50**, 617–629. (d) R. Shang and L. Liu, *Sci. China Chem.* 2011, **54**, 1670–1687. (e) N. Rodríguez and L. J. Gooßen, *Chem. Soc. Rev.* 2011, **40**, 5030–5048. (f) W. I. Dzik, P. P. Lange and L. J. Gooßen, *Chem. Sci.* 2012, **3**, 2671–2678. (g) J. Cornella and I. Larrosa, *Synthesis* 2012, **44**, 653–676. (h) L. J. Gooßen and K. Gooßen, Decarboxylative Coupling Reactions; In *Topics in Organometallic Chemistry*; L. J. Gooßen, Eds.; Springer-Verlag Berlin Heidelberg: 2013; Vol. 44 pp. 121–142. (i) G. J. P. Perry and I. Larrosa, *Eur. J. Org. Chem.* 2017, **2017**, 3517–3527.

¹⁴ G. J. P. Perry, J. M. Quibell, A. Panigrahi and I. Larrosa, *J. Am. Chem. Soc.* 2017, **139**, 11527–11536.

¹⁵ Previous decarboxylative halogenation procedures have required irradiation See ref. 8f-g and: (a) A. I. Meyers and M. P. Fleming, *J. Org. Chem.* 1979, **44**, 3405–3406. (b) K. Kulbitski, G. Nisnevich and M. Gandelman, *Adv. Synth. Catal.* 2011, **353**, 1438–1442.

¹⁶ See references 12b and 12d-f.

¹⁷ (a) L. J. Johnston, J. Lusztyk, D. D. M. Wayner, A. N. Abeywickreyma, A. L. J. Beckwith, J. C. Scaiano and K. U. Ingold, *J. Am. Chem. Soc.* 1985, **107**, 4594–4596. (b) M. Newcomb, *Encyclopedia of Radicals in Chemistry, Biology and Materials*; John Wiley & Sons, Ltd.: Chichester, UK, 2012; pp 1–19.

¹⁸ See reference 6c and R. G. Johnson and R. K. Ingham, *Chem. Rev.* 1956, **56**, 219–269.

¹⁹ Attempts to establish benzoyl hypobromite I as a definite intermediate were unsuccessful. However, the formation of such acyl hypohalites are reported in the literature, and it is therefore reasonable to suggest its presence in this system. See: (a) J. J. Kleinberg, *Chem. Educ.* 1946, **23**, 559–562. (b) R. A. Zingaro, J. E. Goodrich, J. Kleinberg, A. J. Vanderwerf, *J. Am. Chem. Soc.* 1949, **71**, 575–576. (c) D. D. Tanner, N. J. Bunce, In *Acyl Halides*; S. Patai, Eds; John Wiley & Sons, Ltd.: Chichester, UK, 1972; pp 455–500. (d) D. H. R. Barton, H. P. Faro, E. P. Serebryakov, N. F. J. Woolsey, *J. Chem. Soc.* 1965, 2438–2444. (e) P. C. Srivastava, P. Singh, M. Tangiri, A. Sinha, S.J. Bajpai, *Indian Chem. Soc.* 1997, **74**, 443–445.

²⁰ C. Colletto, S. Islam, F. Julia-Hernandez and I. Larrosa *J. Am. Chem. Soc.* 2016, **138**, 1677–1683.

²¹ (a) L. Melander and W. H. Saunders, *Reaction Rates of Isotopic Molecules*, John Wiley & Sons, Ltd.: New York, NY, 1980. (b) M.

Wolsfberg, W. A. Van Hook, P. Paneth and L. P. N. Rebelo, *Isotope effects in the Chemical, Geological, and Bio Sciences;* Springer, Dodrecht, 2010.

²² (a) B. W. Bromley, G. D. Hegeman and W. Meinschein, *Anal. Biochem.* 1982, **126**, 436–446. (b) J. F. Marlier and M. H. O'Leary, *J. Am. Chem. Soc.* 1984, **106**, 5054–5057. (c) J. F. Marlier and M. H. O'Leary, *J. Am. Chem. Soc.* 1986, **108**, 4896–4899. (d) *Molecules* 2013, **18**, 9278–9292. (e) A. A. Vandersteen, G. W. Howe, B. S. Lollar and R. Kluger, *J. Am. Chem. Soc.* 2017, **139**, 15049–15053.

²³ (a) D. A. Singleton, A. A. Thomas, J. Am. Chem. Soc. 1995, 117, 9357–9358. For applications see ref 20 and: (b) A. E. Keating, S. R. Merrigan, D. A. Singleton, K. N. Houk, J. Am. Chem. Soc. 1999, 121, 3933–3938. (c) D. A. Singleton, S. R. Merrigan, B. J. Kim, P. Beak, L. M. Phillips and J. K. Lee, J. Am. Chem. Soc. 2000, 122, 3926–3300. (d) D. A. Singleton, Y. Wang, H. W. Yang and D. Romo, Angew. Chem. Int. Ed. 2002, 41, 1572–1575.

²⁴ For information on the solvent correction model (SMD) see: (a) V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B* 2009, **113**, 6378-6396. For the dispersion correction (DFT-D3(BJ)) see: (b) S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.* **2010**, *132*, 154104-154119. (c) S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.* **2011**, *32*, 1456-1465. See SI for further details.

²⁵ The signal of C4 and C6 was used as "internal standard" in the determination of the isotopic composition of the starting material before and after the reaction. For further information see SI and ref 23a.

²⁶ Kinetic isotope effects were calculated using ISOEFF98 according to the Bigeleisen equation. (a) V. Anisinov, P. Paneth, *Journal of Mathematical Chemistry* 1999, **26**, 75–86. (b) V. Anisinov, P. Paneth, ISOEFF98, Lodz, Poland, 1998. Tunnelling correction was applied using Pyquiver according to Bell's infinite-parabola model. The effect was found to be negligible (c) T. L. Anderson and E. E. Kwan, *PyQuiver* 2016. (d) R. P. Bell, *Chem. Soc. Rev.* 1974, **3**, 513–544. For further information see SI

²⁷ In decarboxylation reactions the KIE on C1 is found to increase as the transition state becomes more product-like. See references
22b-e and (a) J. Bigeleisen, J. Chem. Phys. 1949, 17, 675–678. (b) C. Lewis, P. Paneth, M. H. O'Leary and D. Hilvert, J. Am. Chem. Soc.
1993, 115, 1410–1413. (c) G. G. B. Tcherkez, G. D. Farquhar, T. J. Andrews, Proc. Natl Acad. Sci. USA 2006, 103, 7246–7251.

²⁸ After the C1–Br bond is formed the gradient drops approaching 0, giving rise to a so-called "hidden intermediate". For information see: (a) E. Kraka and D. Cremer, *Acc. Chem. Res.* 2010, **43**, 591–601. (b) H. S. Rzepa and C. Wentrup, *J. Org. Chem.* 2013, **78**, 7565–7574. (c) D. Roca-Lopez, V. Polo, T. Tejero and P. Merino *J. Org. Chem.* 2015, **80**, 4076–4083.

²⁹ See SI for further details.

Transition-Metal-Free Decarboxylative Bromination of Aromatic Carboxylic Acids

Jacob M. Quibell, Gregory J. P. Perry, D. M. Cannas and Igor Larrosa

Aromatic acids are converted into aryl bromides simply and efficiently *via* decarboxylation providing new depth and insight into Hunsdiecker-type reactivity

