## Chemical Science

#### Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: X. Hu, F. Liu, X. Zhang, Z. Zhao and S. Liu, *Chem. Sci.*, 2020, DOI: 10.1039/D0SC00409J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the <u>Information for Authors</u>.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/chemical-science

View Article Online

View Journal

#### ARTICLE

### Expected and unexpected photoreactions of 9-(10-)substituted anthracene derivatives in cucurbit[n]uril hosts

Xianchen Hu,<sup>a</sup> Fengbo Liu,<sup>a</sup> Xiongzhi Zhang,<sup>a,b</sup> Zhiyong Zhao<sup>a,b</sup> and Simin Liu\*<sup>a,b</sup>

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

By arranging substrates in a "reaction ready" state through noncovalent interactions, supramolecular nanoreactors/catalysts show high selectivity and/or rate acceleration features. Herein, we report the host–guest complexation of 9-(10-)substituted anthracene derivatives (**G1–G3**) with cucurbit[*n*]uril (CB[*n*], *n* = 8, 10), and the photoreactions of these derivatives in the presence of CB[*n*] hosts. Both CB[10] and CB[8] showed no obvious effects on the photoreaction of 9,10-disubstituted derivative **G1**. For **G2** and **G3**, CB[10] operated as either a nanoreactor or catalyst (10%) for the photodimerization of two compounds with high selectivity and high yield. However, although CB[8] formed a 1:2 complex with **G2**, as also observed with CB[10], the photosolvolysis product (9-anthracenemethanol) was obtained quantitatively after photoirradiation of the CB[8]·2**G2** complex. This unexpected photosolvolysis was rationalized by a plausible catalytic cycle in which anthracene acts as a photoremovable protecting group (PPG) and the carbonium ion intermediate is stabilized by CB[8].

#### Introduction

Inspired bv natural enzymes, supramolecular nanoreactors/catalysts have been developed to achieve reactions with high selectivity and/or rate acceleration.<sup>1,2</sup> In host-guest chemistry, many types of macrocyclic host molecules have been explored as supramolecular nanoreactors/catalysts, such as cyclodextrins,<sup>3</sup> calixarenes,<sup>4</sup> pillararenes,<sup>5</sup> and nanocages, among others.<sup>6–9</sup> Photochemical reactions of anthracene and its derivatives, including anthracene dimerization and other unimolecular photoreactions, have been widely investigated over the last 100 years owing to the unique photo-responsive properties of these compounds.<sup>10,11</sup> However, the anthracene group acted as photoremovable protecting group (PPG) in only one example.<sup>12,13</sup> In supramolecular chemistry, host-guest interactions have been introduced to regulate the photodimerization of anthracene derivatives.<sup>14–18</sup> Furthermore, anthracene dimerization has been further used in the design of host-guest-related supramolecular polymers and optical materials/devices.19-25

Cucurbit[*n*]urils (CB[*n*]s, n = 5-8, 10) are a family of pumpkinlike macrocyclic hosts. CB[*n*] compounds bear a hydrophobic cavity and two identical portals surrounded by negative carbonyl groups as structural features, and have potential applications in many fields.<sup>26–29</sup> Similar to other hosts, CB[*n*]s





have also been employed to promote various organic reactions, especially photodimerization reactions,<sup>30–32</sup> but have acted as supramolecular catalysts in only a few examples because the product is usually "trapped" in the CB[n] cavity, inhibiting the catalytic process.<sup>33,34</sup>

In this work, we investigated host–guest complexation between CB[n] (n = 8, 10) and 9-(10-)substituted anthracene derivatives (**G1–G3**) (Fig. 1) in aqueous solution, and the CB[n]-mediated (n = 8, 10) photoreaction of these guests. Notably,

<sup>&</sup>lt;sup>a.</sup> The State Key Laboratory of Refractories and Metallurgy, School of Chemistry and Chemical Engineering, Wuhan University of Science and Technology, Wuhan 430081, China. E-mail: <u>liusimin@wust.edu.cn</u>.

<sup>&</sup>lt;sup>b.</sup> Institute of Advanced Materials and Nanotechnology, Wuhan University of Science and Technology, Wuhan 430081, China.

<sup>&</sup>lt;sup>+</sup> Electronic supplementary information (ESI) available: experimental details, binding <sup>1</sup>H NMR spectra, UV/Vis spectra of guests with CB[*n*], and relevant COSY, ESI-MS, <sup>13</sup>C NMR spectra, kinetics fitting. See DOI: 10.1039/x0xx00000x

emical Science Accep

#### Journal Name

# This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence Dpen Access Article. Published on 24 April 2020. Downloaded on 4/26/2020 3:20:40 AM

ARTICLE

#### photoreactions of macrocyclic host-promoted 9-(10-)substituted anthracene derivatives have not been reported previously.<sup>35–39</sup> Our results showed that CB[10] and CB[8] had no obvious effects on the photoreaction of G1, despite encapsulating G1. However, for G2 and G3, CB[10] operated as either a nanoreactor or catalyst (10%) for the photodimerization of two compounds with high selectivity and high yield. Although CB[8] formed a 1:2 complex with G2, as also observed with CB[10], photosolvolysis of G2 was unexpectedly observed instead of dimerization after photoirradiation of the CB[8]·2G2 complex (Scheme 1). A plausible reaction mechanism, referring to anthracene as a PPG is provided.



Scheme 1 Schematic illustration of CB[n]-mediated photoreactions of 9substituted anthracene derivative G2.

#### **Results and Discussion**

#### Complexation of G1 with CB[n] and photoreactions of complexes

Initially, the host–guest complexation of **G1** with CB[n] (n = 8, 10) was investigated by NMR, UV/Vis, and fluorescence spectroscopy, and ESI-MS analysis. <sup>1</sup>H NMR titration experiments clearly demonstrated the formation of host-guest complexes CB[10]·G1 and CB[10]·2G1. As shown in Fig. 2b, binding between G1 and CB[10] exhibited slow exchange kinetics on the <sup>1</sup>H NMR time scale, because both free and bound proton signals were observed. The slow exchange kinetics also allowed the binding ratio to be calculated as 1:2 (host/guest) from the integrals for peaks of bound guest and host. When the amount of CB[10] added was increased to 0.5 equiv. (Fig. 2c), a new set of bound guest peaks (marked as '&') was observed, along with a tiny amount of free guest (marked as '\*'). Clearly the new inclusion complex also exhibited slow exchange kinetics on the <sup>1</sup>H NMR time scale. When excess CB[10] was added to a solution of G1 (free CB[10] is insoluble in water and complexation with the guest usually allows CB[10] to enter the aqueous phase<sup>40</sup>), we calculated the binding ratio of CB[10] with **G1** to be 1:1, according to <sup>1</sup>H NMR peak integration (Fig. 2d). ESI-MS also indicated the existence of two inclusion complexes, CB[10]·G1 and CB[10]·2G1. As shown in Fig. S1a, ion peaks at m/z 577.5, 769.6, and 781.6 were observed, corresponding to



Fig. 2 <sup>1</sup>H NMR spectra (600 MHz, D<sub>2</sub>O, 298 K): of (a) free G1 (2.5 mM); (b) 4:1 mixture of G1 and CB[10]; (c) 2:1 mixture of G1 and CB[10]; (d) G1 and excess CB[10] (resonances of free G1 are marked with '\*'; resonances of CB[10] 2G1 are marked with '#', resonances of CB[10] G1 are marked with '&'; proton signals of impurities are marked with 'x').

1:2 complex CB[10]·2G1 ([CB[10] + 2G1<sup>4+</sup> - 4H<sup>+</sup>]<sup>4+</sup> = 577.5;  $[CB[10] + 2G1^{4+} - 5H^{+}]^{3+} = 769.6; [CB[10] + 2G1^{4+} - 4H^{+} + CI^{-}]^{3+}$ = 781.6). The ion peak at m/z 996.4, corresponding to 1:1 complex CB[10]·G1, was also observed ([CB[10] + G1<sup>4+</sup> – 2H<sup>+</sup>]<sup>2+</sup> = 996.4) (Fig. S1).

UV/Vis and fluorescence titrations were used to examine the complexation of CB[10] with G1 in aqueous solution. With the addition of CB[10], the long-wavelength absorption of free G1 at  $\lambda_{max}$  = 373 nm showed a bathochromic shift (Fig. S2). In the absence of CB[10], G1 showed the anthracene emission with a maximum wavelength of 422 nm. The fluorescence intensity of G1 changed when CB[10] was added continuously, owing to host-guest complexation of G1 with the CB[10] host (Fig. 3).41 Furthermore, before the amount of CB[10] was increased to ~0.5 equiv., the fluorescence intensity of G1 was observed to



Fig. 3 Fluorescence emission spectra recorded for G1 (31.25  $\mu\text{M})$  upon titration with CB[10]. Inset: Emission of G1 vs. CB[10] equivalent ( $\lambda_{ex}$  = 250 nm,  $\lambda_{em}$  = 422 nm).

#### Journal Name

ARTICLE



continuously decrease. However, at 1.0 equiv., the intensity increased about two-fold compared with that of G1 itself, with a slight redshift. Combined with the NMR data, we speculated that CB[10] initially increased  $\pi-\pi$  stacking of **G1** by encapsulating two guest molecules in the cavity, causing fluorescence quenching through formation of the H-dimer of G1.42 The further formation of 1:1 inclusion complex CB[10]·G1 prevented G1 from stacking and exterior aqueous environment, resulting in a stronger **G1** emission.<sup>41</sup> Meanwhile, host-guest complexation between CB[8] and G1 was examined by <sup>1</sup>H NMR (Fig. S3) and ESI-MS (Fig. S4). Broadening of all proton signals, including those of CB[8], was observed when the amount of CB[8] was 0.5 equiv. (Fig. S3c), indicating that binding between CB[8] and G1 underwent intermediate exchange on the <sup>1</sup>H NMR time scale.<sup>43</sup> ESI-MS showed one ion peak at m/z 826.2, corresponding to 1:1 complex CB[8]·G1 ([CB[8] + G1<sup>4+</sup> - 2H<sup>+</sup>]<sup>2+</sup> = 826.2).

With the binding results in hand, we checked the photoreactions of G1 in the absence and presence of CB[n] (n = 8, 10). Prior to UV irradiation at room temperature, all samples (in D<sub>2</sub>O or pure water) were degassed with nitrogen for 15 min ([G] = 2.5 mM), and the reaction was monitored by  $^{1}$ H NMR and UV/Vis spectroscopy. As shown in Fig. S5a, the photoproducts of G1 were complicated and unidentified. In the presence of 0.25 or 0.5 equiv. of CB[10], ratios which benefitted the dimerization reaction, the dimerization product was not observed as expected (Figs. S5b & S5c). Molecular modeling (MMFF) suggested two G1 molecules could adopt an "X" shape in the CB[10]·2G1 complex owing to electrostatic repulsion between the 9,10-substituted positive groups (Fig. 4),<sup>43</sup> which was not a "reaction ready" state for dimerization. Similar results were obtained for the photoreaction of G1 with various contents of CB[8]. Therefore, guest G2 containing one substituent at the 9-position was designed.

Complexation of G2 with CB[n] and photoreactions of complexes

Similar to G1, we first investigated the host-guest complexation of **G2** with CB[n] hosts, and then conducted the photoreaction of **G2** with/without CB[n] (n = 8, 10). <sup>1</sup>H NMR titrations showed that the aromatic proton and H<sub>a</sub> signals of G2 were shifted upfield ( $\Delta\delta$  = 0.97, 0.87, 0.97, 1.28, 1.05, and 1.63 ppm for protons  $H_1-H_5$  and  $H_a$ , respectively) more than the  $H_b$  and  $H_c$ proton signals ( $\Delta\delta$  = 0.47 and 0.28 ppm, respectively) by at least 0.40 ppm, suggesting that the anthracene moiety had advanced deeper and that the substituent was located around the portals of CB[10] (Fig. S6). Broadening of the guest signals again implied that the binding of CB[10] and G2 underwent intermediate exchange on the <sup>1</sup>H NMR time scale.<sup>43</sup> Upon adding excess CB[10], the ratio of CB[10] to G2 was calculated to be 1:2 based on the integrals of the host and guest signals (Fig. S6c). ESI-MS data further confirmed the formation of 1:2 complex CB[10]·2G2. As shown in Fig. S7, an ion peak was observed at m/z 1081.9, which corresponded to the 1:2 complex ([CB[10] +  $2\mathbf{G2}^{2+} - 2\mathbf{H}^+]^{2+} = 1081.9$ ). UV/Vis and fluorescence titration experiments also corroborated the host-guest interaction of G2 and CB[10] in the aqueous phase. The fluorescence emission intensity of G2 decreased upon adding 0.5 equiv. of CB[10] (Fig. S8), suggesting the existence of  $\pi$ - $\pi$  stacking between the encapsulated anthracene units of G2, similar to that of G1. compared **G1**, the Interestingly. with binding of monosubstituted G2 with CB[8] exhibited slow exchange kinetics on the <sup>1</sup>H NMR time scale, allowing the binding ratio to be calculated as 1:2 (host/guest) through integration (Fig. S9). This 1:2 binding was further verified by ESI-MS (Fig. S10). The differentiation of proton signals on the left and right phenyl rings in G2, and no splits in the proton signals of CB[8] under slow exchange kinetics, suggested that the anthracene moieties were inserted into the cavity of CB[8] with the two incorporated G2 molecules adopting a head-to-tail (h-t) orientation. Indeed, the MMFF-minimized model of complex CB[8]·2G2 showed partial aromatic rings of two G2 molecules located in the CB[8] cavity. The dihedral angle between the plane of the included



**Fig. 5** <sup>1</sup>H NMR spectra (600 MHz,  $D_2O$ , 298 K) of CB[10]-2**G2** ([**G2**] = 2.5 mM) solution with various UV irradiation times: (a) 0, (b) 10, (c) 25 min, and (d) free dimerization product **DG2**.

anthracene group and the equatorial plane of CB[8] was about 59° (Figs. 4 & S11). The distance between two included parallel anthracene groups was about 3.42 Å, suggesting the presence of  $\pi$ - $\pi$  stacking.

Under the same conditions as for G1, photoreactions of G2 were conducted in the absence and presence of CB[n] (n = 8, 10). As shown in Fig. S12, after UV irradiation for 6 h, h-t dimer product DG2 (purified and identified by irradiating complex CB[10]·2G2) was observed with unidentified side products and unreacted G2. In comparison, in the presence of 0.5 equiv. of CB[10], almost 100% of G2 was converted into a single product within 25 min. As shown in Fig. 5b, under UV irradiation for 10 min, G2 was partially converted into presumed new photoproduct(s). G2 was consumed after 25 min, with only one set of CB[10] host and bound product(s) peaks remaining (Fig. 5c). After the appropriate amount of 3,5-dimethylamantadine hydrochloride (3,5-DMADA) was added to the irradiated solution of CB[10]·2G2 to displace the product(s) from the CB[10] cavity (CB[10]·2(3,5-DMADA) is insoluble in water44), only one exclusive product, characterized as h-t dimer DG2, was obtained in almost quantitative yield (characterization data for DG2 is shown in Figs. S13-S15 in SI). By monitoring absorbance changes of **G2** at  $\lambda$  = 370 nm, we calculated the yields of DG2 with various irradiation times. As "the intracomplex photodimerization is unimolecular in nature",18 an apparent rate constant  $(k_1)$  of 0.1158 ± 0.0079 min<sup>-1</sup> for the dimerization reaction of G2 with 50% CB[10] was calculated with first-order kinetics (Fig S16). The half-conversion time ( $t_{1/2}$ , time for conversion of half the substrate) was used because the rate constant could not be accurately measured in some cases. Clearly, in the presence of 50% CB[10], the dimerization reaction of G2 was accelerated tremendously (Table 1). The high conversion, high selectivity, and rate acceleration of this photoreaction of G2 within CB[10] suggested that two G2 molecules were preorganized by CB[10] to adopt a "reaction ready" h-t orientation, which was in good agreement with the MMFF-minimized model of complex CB[10]·2G2 in Fig. 4.

Page 4 of 8

View Article Online Table 1. Half-conversion time (t<sub>1/2</sub>) for guests ທີ່ເຖິ/ໃຫ້ເກີດຢີໃດອີໂກງໃກ່<sup>ດ</sup>ອີ, 10) in water at 298 K.

Host	t <sub>1/2</sub> (G2)	t <sub>1/2</sub> (G3)
None	1.7 hª	7.5 hª
CB[10] (50%)	6 min⁵	37 min <sup>ь</sup>
CB[10] (10%)	44 min <sup>b</sup>	110 min <sup>b</sup>
CB[8] (50%)	8 min <sup>c</sup>	d
CB[8] (5%)	35 min <sup>c</sup>	d

<sup>a</sup> Estimated from NMR integrals (Figs. S12 & S32). <sup>b</sup> Calculated by fitting equation (Figs. S16, S18 & S39). <sup>c</sup> Estimated from diagrams of product yields (Fig. S26). <sup>d</sup> Not determined.

Inspired by a previous sample,<sup>33</sup> we investigated whether CB[10] could act as a supramolecular catalyst to convert **G2** into **DG2** exclusively. In the presence of 0.1 equiv. of CB[10] (10%), almost 100% of **G2** was converted into photoproduct **DG2** with trace amounts of side products within 120 min (Fig. S17). The  $t_{1/2}$  of 44 min was calculated from the first-order kinetics equation (Fig. S18). In comparison, the  $t_{1/2}$  of **G2** without CB[10] was estimated to be 1.7 hours (Table 1). All data suggested that CB[10] operated as the supramolecular catalyst in this case, although the photoreaction took longer to achieve than that with 0.5 equiv. of CB[10] (Scheme 1). As mentioned above, effective replacement of the product (**DG2**) with the starting material (**G2**) was necessary for host (CB[10]) to operate as a catalyst. To verify this, <sup>1</sup>H NMR competition experiments between **DG2** and **G2** with CB[10] were performed.

As predicted, the mixture of **G2**, **DG2**, and CB[10] with a 2:1:1 ratio resulted in most CB[10] being occupied by **G2** (Fig. S19c). An approximate equilibrium constant ( $K_{eq}$ ) was calculated as 8.8 × 10<sup>5</sup> M<sup>-1</sup> (Figs. S20 & S21), suggesting that CB[10] operated as a supramolecular catalyst owing to the spontaneous replacement of product **DG2** with starting material **G2**. The results showed that, as either the supramolecular nanoreactor or catalyst, CB[10] not only shortened the photoreaction time, but also tremendously improved the reaction selectivity.

When the host molecule was switched to CB[8], an unexpected photoreaction of G2 occurred, although CB[8] bound G2 with the same binding ratio as CB[10]. In the presence of 0.5 equiv. of CB[8], precipitate formation was observed when the G2 solution was irradiated with UV light. With extended irradiation time, additional yellow precipitate was generated. Therefore, the solutions after centrifugation were carefully monitored by <sup>1</sup>H NMR. By analyzing the stacking NMR spectra (Fig. 6) and the precipitate, the bound peaks of G2 were found to fully disappear in 60 min, while a singlet peak at 3.1 ppm appeared and remained, and the yellow precipitate contained no CB[8] and was soluble in DMSO and CDCl<sub>3</sub>. First, we suspected that the precipitate was anthraquinone, because anthracene derivatives can easily be oxidized, 45,46 despite the photoreaction being conducted under N<sub>2</sub> atmosphere. However, the NMR signals for the precipitate did not match those of anthraquinone. Further hypotheses included the photosolvolysis of G2. Eventually, the yellow precipitate was verified to be 9-anthracenemethanol by EI-MS and <sup>1</sup>H NMR

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence

Dpen Access Article. Published on 24 April 2020. Downloaded on 4/26/2020 3:20:40 AM

#### EDA k) 60 min j) 50 min i) 40 min h) 30 min g) 20 min Bound G2 f) 10 min e) 8 min d) 6 min C) 4 min b) 2 min a) 0 min 9 8 7 6 5 4 3



(Figs. S22a & S23). When irradiated in air, G2 first generated 9anthracenemethanol as a precipitate (suspended), which was then converted to anthraquinone (Fig. S22b), further confirming the occurrence of photosolvolysis prior to photooxidation. The singlet peak in the aqueous phase represented the ethylene proton signal of ethylenediamine (EDA). And the binding of protonated EDA in CB[8] helped solubilize CB[8] in water at high concentrations (maximum solubility of CB[8] in water is about 0.2 mM). Indeed, NMR titration experiments proved that small EDA molecules could be encapsulated by CB[8] (Fig. S24). Furthermore, CB[8] was tested to determine whether it could act as a supramolecular catalyst in the photosolvolysis of G2. In the presence of 0.05 equiv. of CB[8] (5%), G2 was almost all converted to 9-anthracenemethanol within 4 h (Fig. S25). In comparison, no precipitate was observed without CB[8]. The  $t_{1/2}$ of G2 in the presence of CB[8] was estimated because the data could not be ideally fitted with first-order kinetics (Table 1 & Fig. S26). CB[8] doubtless operated as a supramolecular catalyst for the photosolvolysis of G2. Therefore, why photosolvolysis occurred when G2 itself did not show the same reactivity required further investigation.

After a broad search of the photosolvolysis literature, we noted that a "photoremovable protecting group (PPG)" could be involved in this reaction.<sup>12,47</sup> To date, only one example of an anthracene unit acting as a PPG has been reported, with the plausible mechanism involving heterolysis and/or homolysis of the CH<sub>2</sub>–X bond in PPG–CH<sub>2</sub>–X (X, leaving group) to produce a carbonium ion intermediate. CB[*n*] hosts are known to be capable of stabilizing active carbocations.<sup>48,49,34</sup> Therefore, a plausible mechanism for the photosolvolysis of **G2** with CB[8] as the supramolecular catalyst in aqueous solution was proposed. As shown in Scheme 2, the initial step is light-induced heterolytic cleavage of the C–N bond of encapsulated **G2**, and/or homolytic cleavage of the C–N bond followed by rapid electron transfer (ET), to give the carbonium ion

intermediate and EDA. The CB[8]-stabilized carbonium jon is then attacked by solvent molecule  $H_2O$  to is weak binding with CB[8] and insolubility, 9-anthracenemethanol is displaced by **G2** or EDA and precipitated out of solution, allowing the catalytic process to continue. Compared with the observation of radical side products (caused by homolysis) in most reported examples,<sup>12</sup> the generation of a single product with quantitative yield in this case suggested that, with the assistance of CB[8], either C–N bond heterolysis was favored over homolysis or the electron transfer step after homolysis was accelerated.

The high conversion and high selectivity of this photosolvolysis of **G2** suggested that the orientation of the two **G2** molecules in CB[8] was not only unfavorable for dimerization, but also favorable for stabilization of the carbonium ion intermediate by CB[8]. In addition to acting as the supramolecular catalyst, the most interesting role of CB[8] in this case was efficiently switching the reaction pathway of **G2** from familiar photodimerization to unusual photosolvolysis.



Scheme 2 Plausible reaction mechanism for the photosolvolysis reaction of G2 in the presence of CB[8] host (5%).

#### Complexation of G3 with CB[n] and photoreactions of complexes

Guest molecule **G3** was synthesized to explore the effect of different substituent groups on the photochemical reaction. The host–guest complexation of **G3** with CB[n] (n = 8, 10) was investigated by <sup>1</sup>H NMR, UV/Vis, and fluorescence spectroscopy, and ESI-MS analysis (Figs. S27–S31). The ESI-MS data indicated that the complexation ratio of CB[n] with **G3** was 1:2 (ion peaks at m/z 1100.9 and 934.8 corresponding to 1:2 complexes CB[10]·2**G3** and CB[8]·2**G3**, respectively) (Figs. S29 & S31). Similar to **G2**, the photoreaction of **G3** slowly produced h–t dimerization product **DG3** and unidentified side products (Fig. S32). Furthermore, the dimerization of **G3** with CB[10] operating as a supramolecular nanoreactor and catalyst, exclusively gave **DG3** (Figs. S33–S39). No bound peaks of **DG3** were observed in the NMR competition experiments, indicating that the binding of **G3** with CB[10] was much stronger than that

ence

#### Journal Name

ARTICLE

of **DG3** (Fig. S38). The dimerization reaction of **G3** was also greatly accelerated with assistance from CB[10] (Table 1 & Fig. S39). However, in the presence of CB[8], no precipitate and an extremely complicated <sup>1</sup>H NMR spectrum of photoproducts were observed, suggesting the negative effect of CB[8] on the photoreaction of **G3** (Fig. S40). It is unclear whether the lack of photosolvolysis of **G3** was due to the deactivating effect of the pyridinium substituent group or the unfavorable orientation of **G3** in CB[8].

#### Conclusions

In summary, we have reported the effects of CB[n] (n = 8, 10) hosts on the photoreaction of 9-(10-)anthracene derivatives. Both CB[8]and CB[10] operated as supramolecular nanoreactors and catalysts in the photoreaction of 9-substituted anthracene derivative **G2**. However, CB[10] promoted the photodimerization of **G2**, which occurred slowly with lower selectivity without a host. In contrast, CB[8] exclusively induced the photosolvolysis of **G2**, which did not occur without a host. Furthermore, the photoreactions of two anthracene derivatives, **G1** and **G3**, were investigated in the presence of CB[n] to compare the effect of different CB[n] on the reactivity of anthracene derivatives with various substituents. The results showed that small differences in the host structures could cause large different effects on the host-involved reaction selectivity.

As the first example of CB[10] operating as a supramolecular nanoreactor/catalyst, we anticipate that more reactions, involving large-sized or more than two substrate molecules, can be promoted by the large cavity of CB[10]. The selectivity for photosolvolysis of aromatic substrates with a PPG is also expected to be improved by the ability of CB[*n*] to stabilize cations. These finding will further benefit CB[*n*] chemistry and supramolecular catalysis.

#### **Conflicts of interest**

There are no conflicts to declare.

#### Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (21871216, 21472143).

#### Notes and references

- 1 P. W. N. M. van Leeuwen, in *Supramolecular catalysis*, Wiley-VCH, 2008.
- 2 J. Meeuwissen and J. N. H. Reek, *Nat. Chem.*, 2010, **2**, 615–621.
- 3 U. H. Brinker and J.-L. Mieusset, in *Molecular Encapsulation Organic Reactions in Constrained Systems*, Wiley, 2010, pp. 43–115.
- 4 P. Neri, J. L. Sessler and M.-X. Wang, in *Calixarenes and beyond*, Springer, 2016, pp. 691–742.
- 5 E. H. Wanderlind, D. G. Liz, A. P. Gerola, R. F. Affeldt, V. Nascimento, L. C. Bretanha, R. Montecinos, L. Garcia-Rio, H. D. Fiedler and F. Nome, ACS Catal., 2018, 8, 3343–3347.
- 6 M. Yoshizawa, J. K. Klosterman and M. Fujita, *Angew. Chem. Int. Ed.*, 2009, **48**, 3418–3438.

- 7 S. Zarra, D. M. Wood, D. A. Roberts and J. R. Nitsche, Chem. Soc. New Article Online Rev., 2015, 44, 419–432. DOI: 10.1039/D0SC00409J
- A. Galan and P. Ballester, Chem. Soc. Rev., 2016, 45, 1720–1737.
- 9 S. Sadjadi, in Organic Nanoreactors: From Molecular to Supramolecular Organic Compounds, Academic Press, 2016.
- 10 H. D. Becker, Chem. Rev., 1993, 93, 145-172.
- 11 H. Bouas-Laurent, A. Castellan, J.-P. Desvergne and R. Lapouyade, *Chem. Soc. Rev.*, 2000, **29**, 43–55.
- P. Klán, T. Šolomek, C. G. Bochet, A. Blanc, R. Givens, M. Rubina, V. Popik, A. Kostikov and J. Wirz, *Chem. Rev.*, 2013, **113**, 119–191.
- 13 A. K. Singh and P. K. Khade, *Tetrahedron Lett.*, 2005, **46**, 5563–5566.
- 14 H. Wang, F. Liu, R. C. Helgeson and K. N. Houk, *Angew. Chem. Int. Ed.*, 2013, **52**, 655–659.
- 15 A. Tron, H.-P. Jacquot de Rouville, A. Ducrot, J. H. R. Tucker, M. Baroncini, A. Credi and N. D. McClenaghan, *Chem. Commun.*, 2015, **51**, 2810–2813.
- 16 J.-C. Gui, Z.-Q. Yan, Y. Peng, J.-G. Yi, D.-Y. Zhou, D. Su, Z.-H. Zhong, G.-W. Gao, W.-H. Wu and C. Yang, *Chin. Chem. Lett.*, 2016, 27, 1017–1021.
- 17 X. Wei, W. Wu, R. Matsushita, Z. Yan, D. Zhou, J. J. Chruma, M. Nishijima, G. Fukuhara, T. Mori, Y. Inoue and C. Yang, *J. Am. Chem. Soc.*, 2018, **140**, 3959–3974.
- 18 J. Ji, W. Wu, W. Liang, G. Cheng, R. Matsushita, Z. Yan, X. Wei, M. Rao, D.-Q. Yuan, G. Fukuhara, T. Mori, Y. Inoue and C. Yang, *J. Am. Chem. Soc.*, 2019, **141**, 9225–9238.
- 19 J.-F. Xu, Y.-Z. Chen, L.-Z. Wu, C.-H. Tung and Q.-Z. Yang, Org. Lett., 2013, 15, 6148–6151.
- 20 P. Wei, X. Yan and F. Huang, *Chem. Commun.*, 2014, **50**, 14105–14108.
- 21 X. Zhang, Y. Gao, Y. Lin, J. Hu and Y. Ju, *Polym. Chem.*, 2015, **6**, 4162–4166.
- 22 W. Guan, G. Wang, J. Ding, B. Li and L. Wu, Chem. Commun., 2019, 55, 10788–10791.
- 23 Q. Zhou, B. Zhang, D. Han, R. Chen, F. Qiu, J. Wu and H. Jiang, *Chem. Commun.*, 2015, **51**, 3124–3126.
- 24 W. Zhou, Y. Chen, Q. Yu, P. Li, X. Chen and Y. Liu, *Chem. Sci.*, 2019, 10, 3346–3352.
- 25 M. Tu, H. Reinsch, S. Rodríguez-Hermida, R. Verbeke, T. Stassin, W. Egger, M. Dickmann, B. Dieu, J. Hofkens, I. F. J. Vankelecom, N. Stock and R. Ameloot, *Angew. Chem. Int. Ed.*, 2019, **58**, 2423–2427.
- 26 J. Lagona, P. Mukhopadhyay, S. Chakrabarti and L. Isaacs, Angew. Chem., Int. Ed., 2005, 44, 4844–4870.
- 27 E. Masson, X. Ling, R. Joseph, L. Kyeremeh-Mensah and X. Lu, *RSC Adv.*, 2012, **2**, 1213–1247.
- 28 S. J. Barrow, S. Kasera, M. J. Rowland, J. del Barrio and O. A. Scherman, *Chem. Rev.*, 2015, **115**, 12320–12406.
- 29 X. Yang, F. Liu, Z. Zhao, F. Liang, H. Zhang and S. Liu, Chin. Chem. Lett., 2018, 29, 1560–1566.
- 30 K. I. Assaf and W. M. Nau, Chem. Soc. Rev., 2015, 44, 394–418.
- 31 S. Sadjadi, in *Organic Nanoreactors: From Molecular to Supramolecular Organic Compounds*, Elsevier, 2016, pp. 43–84.
- 32 K. Kim, in *Cucurbiturils and Related Macrocycles*, The Royla Society of Chemistry, 2019, pp. 86–120.
- 33 Y. Kang, X. Tang, H. Yu, Z. Cai, Z. Huang, D. Wang, J.-F. Xu and X. Zhang, *Chem. Sci.*, 2017, 8, 8357–8361.
- 34 L. Scorsin, J. A. Roehrs, R. R. Campedelli, G. F. Caramori, A. O. Ortolan, R. L. T. Parreira, H. D. Fiedler, A. Acuña, L. García-Río and F. Nome, ACS Catal., 2018, 8, 12067–12079
- 35 C. Yang, T. Mori, Y. Origane, Y. H. Ko, N. Selvapalam, K. Kim and Y. Inoue, J. Am. Chem. Soc., 2008, 130, 8574–8575.
- F. Biedermann, I. Ross and O. A. Scherman, *Polym. Chem.*, 2014, 5, 5375–5382.
- 37 C. P. Carvalho, Z. Domínguez, J. P. Da Silva and U. Pischel, *Chem. Commun.*, 2015, **51**, 2698–2701.
- 38 C.-H. Tung and J.-Q. Guan, J. Org. Chem., 1998, 63, 5857-5862.
- 39 D.-Y. Wu, B. Chen, X.-G. Fu, L.-Z. Wu, L.-P. Zhang and C.-H. Tung,

Journal Name

Org. Lett., 2003, 5, 1075–1077.

- 40 W. Gong, X. Yang, P. Y. Zavalij, L. Isaacs, Z. Zhao and S. Liu, *Chem. Eur. J.*, 2016, **22**, 17612–17618.
- 41 R. N. Dsouza, U. Pischel and W. M. Nau, *Chem. Rev.*, 2011, **111**, 7941–7980.
- 42 N. Barooah, J. Mohanty and A. C. Bhasikuttan, *Chem. Commun.*, 2015, **51**, 13225–13228.
- 43 X. Zhao, F. Liu, Z. Zhao, H. Karoui, D. Bardelang, O. Ouari and S. Liu, *Org. Biomol. Chem.*, 2018, **16**, 3809–3815.
- 44 X. Yang, Z. Zhao, X. Zhang and S. Liu, Sci. China Chem., 2018, 61, 787–791
- 45 K. I. Assaf, M. A. Alnajjar and W. M. Nau, *Chem. Commun.* 2018 54, 1734–1737. DOI: 10.1039/DOSC00409J
- 46 P. Anzenbacher, T. Niwa, L. M. Tolbert, S. R. Sirimanne and F. P. Guengerich, *Biochemistry*, 1996, **35**, 2512–2520.
- 47 P. Wang, Asian J. Org. Chem., 2013, 2, 452-464.
- 48 R. Wang and D. H. Macartney, *Tetrahedron Lett.*, 2008, 49, 311–314.
- 49 N. Basilio, L. García-Río, J. A. Moreira and M. Pessêgo, J. Org. Chem., 2010, 75, 848–855.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence

Open Access Article. Published on 24 April 2020. Downloaded on 4/26/2020 3:20:40 AM.

**Chemical Science Accepted Manuscript** 

#### Expected and unexpected photoreactions of 9-(10-)substituted anthracene yiew Article Online derivatives in cucurbit[*n*]uril hosts

Xianchen Hu, Fengbo Liu, Xiongzhi Zhang, Zhiyong Zhao and Simin Liu

Cucurbit[*n*]uril (CB[*n*], n = 8, 10) operated as a supramolecular nanoreactor and catalyst for the photoreaction of 9-substituted anthracene derivative with high selectivity and high yield, in which CB[10] promoted the photodimerization while CB[8] exclusively induced the occurrence of photosolvolysis of the anthracene derivative.

