


SYNTHESIS OF TRANS-4,8,12,15,15-PENTAMETHYLBICYCLO[9,3,1]PENTADEC-3E,7Z,12Z-TRIENE, A GEOMETRICAL ISOMER OF ANHYDROVERTICILLOL¹

Takashi KUMAGAI, Fumiaki ISE, Tadao UYEHARA, and Tadahiro KATO*
 Department of Chemistry, Faculty of Science, Tohoku University,
 Sendai 980

A 7Z-isomer of anhydroverticillol skeleton was constructed in 15% yield when the 7E allyl bromide having secoverticillol framework was treated with $\text{LiN}(\text{TMS})_2$. Reductive removal of the SO_2Ph group furnished a geometrical isomer of anhydroverticillol.

Verticillol (1)², a constituent of conifer wood of *Sciadopitys verticillata* Sieb. et Zucc. represents a class of structurally fascinating diterpenes that have not yet yielded to chemical synthesis. Its carbon skeleton is biogenetically envisaged as correlating with that of cembreoid such as neocembrene (2)³. Both may be derived from a common precursor (3) arisen from geranylgeraniol in the conifer wood.

After achievement of the synthesis⁴ of cembrene type compounds, our effort has been expended to develop the construction method of the former skeleton⁵. A strategy of our synthesis is based on the biogenetical consideration on its formation and a key feature is to apply the regioselective coupling reaction of properly functionalized allyl chloride (6) with the terminal double bond of phenyl prenyl sulfones (7a and b) by virtue of SnCl_4 followed by ring closure to make seco-verticillol



only one cis (Z) olefinic methyl at 14.5 ppm. The anhydroverticillol (5b, W = H) shows two signals at 15.2 and 15.7 ppm due to the cis olefinic methyls. This fact suggests that the geometrical isomerization took place at C₇=C₈ bond to afford an equilibrium mixture of 7E and 7Z isomers of 10b, the latter of which might be converted into 5a (W = SO₂Ph) under the basic conditions. In fact, it was revealed that the geometrically pure model compound (12) gave a 5:1 mixture of 12 and its isomers (13 plus 14) when treated with LiN(TMS)₂ in THF at -78°C for 1 hr¹⁹.

The ester group of the cis isomer (10a, X = CO₂Me) was similarly transformed in 95% overall yield into allyl bromide (10a, X = CH₂Br), from which construction of anhydroverticillol skeleton corresponding to 4 was attempted by the same procedure. It was, however, in marked contrast to the reaction of the cis allyl bromide where no C-C bond formation occurred entirely under a variety of conditions including LiN(TMS)₂ in THF.

Our present experiments have led to the conclusion that cis-trans isomerization takes place at the allyl bromide moiety of 10b and presumably of 10a prior to the requisite C-C-bond formation when treated with LiN(TMS)₂ and that only 7Z isomer of 10b can make the C-C bond, resulting in the formation of a geometrical isomer of anhydroverticillol skeleton.

References

1. Constitutes part 35 of the series, Cyclization of Polyenes. Part 33, T. Kato, M. Suzuki, T. Kobayashi, and Barry P. Moore, J. Org. Chem., 45, 1126 (1980); Part 34, Chem. Commun., in the press (1980).
2. B. Karlsson, Anne-M. Pilotti, Anne-C. Soderholm, T. Norin, S. Sundin, and M. Sumimoto, Tetrahedron, 34, 2349 (1978).
3. Concerning the synthesis of 2, see ref. 1.
4. H. Takayanagi, T. Uyehara, and T. Kato, Chem. Commun., 359 (1978) and references cited therein.
5. T. Kato, H. Takayanagi, T. Suzuki, and T. Uyehara, Tetrahedron Lett., 1201 (1978).
6. The phenyl prenyl sulfones (7a and b) were prepared from the corresponding alcohols by the sequential treatments with CBr₄ and PPh₃ followed by NaSO₂Ph.
7. For the preparation of 6, see ref. 4.
8. Dehydrochlorination of 8a with silica gel⁴ afforded exclusively the isopropenyl derivative.
9. 9b was also prepared from the isolated 8a in 70% yield by the action of SnCl₄ at -30°C for 2 hr.
8a PMR (CCl₄) δ 1.51 and 1.54 (each 3H, sharp s, CClMe₂), 1.66, 1.74 and 1.81 (each 3H, bs, C=CMe), 3.64 (2H, d, 7 Hz, CH₂SO₂Ph), 5.08 (t, 7 Hz), 5.17 (t, 7 Hz), and 6.63 (m) (each 1H, C=CH) ppm.
10. All the silica gel column chromatographies were carried out using a mixed solvent of hexane:AcOEt in the ratio of 5:1 ~ 3:1.
11. From the mixture of the cyclic chlorides was isolated pure 9a by high pressure LC. 9a was converted into 10a in 81% yield with LiCl in DMF at 100°C.
9a PMR δ 0.82, 0.98, 1.61, 1.63, 1.82 (each 3H, s, Mes), 3.07 (dd, 3 and 15 Hz) and 3.24 (dd, 4 and 15 Hz) (each 1H, CH₂SO₂Ph), 5.11 (bt, 7 Hz) and 6.61 (m) (each

- 1H, C=CH) ppm: CMR δ Mes (q) 12.4, 16.1, 17.1, 28.6, 34.9 and 51.7; CH₂ (t) 23.3, 27.2, 29.0, 38.4, 43.2, and 56.6; CH (d) 48.5, 50.5, 124.8, 127.7, 129.4, 133.6, and 142.2; quart. C (s) 38.7, 75.0, 135.0, 141.1, and 141.8; CO₂Me 170.3 ppm.
12. Physical evidence supports the structures. 10a (X = CO₂Me) PMR δ 0.69 and 0.90 (each 3H, s, C₁₅-Mes), 1.58, 1.73, and 1.81 (each 3H, bs, C=CMes), 3.00 (dd, 4 and 15 Hz) and 3.09 (dd, 3 and 15 Hz) (each 1H, CH₂SO₂Ph), 5.07 (m), 5.35 (bs), and 6.59 (m) (each 1H, C=CHs) ppm: CMR δ Mes (q) 12.4, 16.0, 17.4, 22.4, 26.8, 51.7; CH₂ (t) 27.1, 28.0, 28.5, 38.3, 56.0; CH (d) 43.8, 44.0, 123.3, 124.7, 128.2, 129.3, 133.6, 142.1; quart. C (s) 36.1, 127.5, 132.6, 134.7, 141.8; CO₂Me 170.2 ppm. 10b (X = CO₂Me) PMR δ 0.87 and 0.98 (each 3H, s, C₁₅-Mes), 1.61, 1.67, and 1.80 (each 3H, C=CMes), 2.78 and 3.19 (dd, 4 and 15 Hz) (each 1H, CH₂SO₂Ph), 5.09 (bt, 7 Hz), 5.23 (bs), and 6.65 (m) (each 1H, C=CHs) ppm: CMR δ Mes (q) 12.5, 16.1, 21.5, 22.5, 25.5, 51.8; CH₂ (t) 27.2, 28.0, 29.4, 38.5, 59.2; CH (d) 38.5, 45.3, 122.4, 124.7, 128.3, 129.6, 133.9, 142.5; quart. C (s) 35.4, 127.8, 134.8, 135.6, 140.9; CO₂Me 162.5 ppm.
13. The numbering of 9 and 10 is based tentatively on that of anhydroverticillol.
14. S. Torii, K. Uneyama, and S. Matsumoto, J. Org. Chem., 45, 16 (1980). The stereochemistry of 11a was unequivocally confirmed by an X-ray analysis by Dr. C. Kabuto (Tohoku University), to whom we appreciate. We are indebted to professors S. Torii and K. Uneyama for their kind supply of 11a and b.
15. The isomeric compound corresponding to 5a (W = SO₂Ph) could not be obtained at all.
16. Physical evidence supports the structure of 5a. 5a (W = SO₂Ph) PMR δ 0.80 and 0.82 (each 3H, s, gem-Mes), 1.67 (3H) and 1.80 (6H) (C=CMes), 4.93 (bd, 7 Hz), 5.31 (bt, 7 Hz), and 5.55 (bs) (each 1H, C=CHs) ppm. 5a (W = H) PMR δ 0.73 and 0.83 (each 3H, s, gem-Mes), 1.60, 1.62, and 1.69 (each 3H, bs, C=CMes), 5.03 (bd, 9 Hz), 5.39 (t, 9 Hz), and 5.44 (bs) (each 1H, C=CHs) ppm: CMR Mes (q) 14.5, 21.8, 22.8, 22.9, 26.6; CH₂ (t) 22.0, 24.6, 26.4, 28.5, 32.2, 34.9; CH (d) 40.9, 54.1, 123.1, 125.8, 127.9; quart. C (s) 34.2, 134.0, 134.4, and 135.4 ppm.
17. Dehydration of verticillol (1), kindly provided by professor M. Sumimoto, with several reagents described in the literatures²⁰ afforded a mixture of 5a (W = H) and the exocyclic isomer. Refluxing the mixture in toluene with iodine converted the latter into the former exclusively. Anhydroverticillol (5a, W = H) PMR δ 0.73 and 0.80 (each 3H, s, gem-Mes), 1.49, 1.57, and 1.76 (each 3H, bs, C=CMes), 4.75 (bd, 9 Hz), 5.22 (bd, 14 Hz), and 5.30 (bs) (each 1H, C=CHs) ppm: CMR δ Mes (q) 15.2, 15.7, 23.0, 23.7, 27.1; CH₂ (t) 21.5, 26.7, 30.8, 34.1, 39.6, 41.0; CH (d) 38.0, 42.5, 121.6, 124.7, 129.9; quart. C (s) 35.7, 132.7, 132.9, 135.9 ppm.
18. GLC was carried out using a Golay column Z-45 with column temperature at 200°C.
19. The isomers (13 plus 14) were separated by high pressure LC with hexane:AcOEt = 30:1.
20. a, H. Erdtman, T. Norin, M. Sumimoto, and A. Morrison, Tetrahedron Lett., 3879 (1964); b, C. Kaneko, S. Hayashi, and M. Ishikawa, Chem. Pharm. Bull. Jpn., 12, 1510 (1964).

(Received October 23, 1980)