

A NEW STEREOSELECTIVE ROUTE TO γ -BUTYROLACTONES: ASYMMETRIC SYNTHESSES OF (+)-*trans*-WHISKY AND (+)-*trans*-COGNAC LACTONES


Okiko MIYATA,^a Tetsuro SHINADA,^a Nanako KAWAKAMI,^a Kyoko TAJI,^a Ichiya NINOMIYA,^a Takeaki NAITO,^{*,a} Tadamasa DATE,^b and Kimio OKAMURA^b

Kobe Women's College of Pharmacy, ^aMotoyamakita, Higashinada, Kobe 658, Japan and Research Laboratory, Tanabe Seiyaku, Co., Ltd., ^bKawagishi, Toda, Saitama 335, Japan

A new stereoselective route to γ -butyrolactones such as (+)-*trans*-whisky and (+)-*trans*-cognac lactones (**1a**, **b**) has been developed by a combination of three key reactions: diastereoselective nucleophilic addition of thiophenol, cleavage of chiral N-acylsultam *via* thioester, and intramolecular displacement of the sulfonium group with carboxylate anion.

KEYWORDS asymmetric synthesis; whisky lactone; cognac lactone; butyrolactone; nucleophilic addition; thiophenol; (+)-sultam; ate complex


Structurally simple γ -butyrolactones are widespread naturally occurring substances found not only as sex pheromones¹⁾ but also as key flavor components.²⁾ The biological activities of these substances are strictly dependent on the absolute configuration of the chiral C4-carbon atom which is attached by oxygen to the lactone ring.³⁾ We have now provided a new method for the construction of two contiguous stereogenic centers at the C3- and C4-positions of γ -butyrolactones by demonstrating asymmetric syntheses of (+)-*trans*-whisky and (+)-*trans*-cognac lactones (**1a**, **b**), which are key components for the flavors of whisky, wine, and cognac.⁴⁻⁶⁾ Our synthetic strategy consists of two crucial reactions, diastereoselective nucleophilic addition of thiophenol to α,β -unsaturated acid derivatives (**2a**, **b**) having (+)-sultam and stereoselective displacement reaction of the corresponding sulfonium group with carboxylate anion.



Acylation of (+)-sultam, Oppolzer's reagent,⁷⁾ with β -(*n*-butyl)methacryloyl chloride in the presence of sodium hydride gave the chiral olefin **2a** in 90% yield, which was then treated with 10 eq of thiophenol in the presence of 0.3 eq of lithium thiophenoxide in THF at 0 °C^{8,9)} to give the chiral adduct **3a** in 83% yield in addition to small amounts of unidentified diastereoisomers. The stereostructure of the adduct **3a** was firmly established by its X-ray crystallography, which supported the nature of the addition reaction as *anti*-addition of thiophenol as in the case of our previous work.¹⁰⁾ The proposed mechanism for the diastereoselective *anti*-addition of thiophenol is as follows. The starting chiral olefin **2a** takes *s-trans*-conformation **A** due to steric hindrance between the vinyl methyl group and the C7-methylene part of the chiral auxiliary found in the corresponding *s-cis*-conformation. Lithium thiophenoxide would attack from

a convex α -face to form the enolate **B**, which is then protonated from β -face due to the stereoelectronic effect of the newly introduced sulfur group to give the (2*S*,3*S*)-adduct **3a**.

Since the application of conventional methods (LiOH ,⁷ $\text{LiOH-H}_2\text{O}_2$ ¹¹) to the cleavage of *N*-acylcamphor sultam was unsuccessful, we have developed a new method for its cleavage without loss of the induced chirality and with virtually complete recovery of the chiral auxiliary. Treatment of the adduct **3a** with aluminum thiobenzoyloxy "ate" complex, prepared from *n*-butyl lithium, benzyl mercaptan, and trimethylaluminum, gave the thioester **4a** in 84% yield. Both lithium benzylthiophenoxide and aluminum thiophenoxide "ate" complex were ineffective for the cleavage of *N*-acylcamphor sultam. Hydrolysis of the thioester **4a** in the presence of silver perchlorate proceeded smoothly to give the corresponding acid **5a** in 94% yield, which was then subjected to Arndt-Eistert reaction to afford the homologous acid **6a** in 51% yield with three steps from **5a**. *S*-Alkylation of the sulfide **6a** with methyl iodide in the presence of silver perchlorate followed by treatment of the resulting sulfonium salt with cesium carbonate underwent smooth lactonization^{12,13} by intramolecular substitution reaction with the naked carboxylate anion.¹⁴



Products were found to be a diastereomeric mixture (98:2) by the NMR spectrum and carefully purified by preparative TLC. The major lactone **1a** thus prepared was identical with the authentic sample of (+)-*trans*-whisky lactone upon comparison of their spectral data and optical rotations ($[\alpha]_D +79^\circ$ (c=1.22, MeOH) (lit.⁴) $[\alpha]_D +79^\circ$ (c=1.04, MeOH)).

Similarly, asymmetric synthesis of (+)-*trans*-cognac lactone (**1b**) was accomplished by the same reaction sequence from β -(n-pentyl)methacryloyl chloride. The product **1b** was also identical with the authentic sample of (+)-*trans*-cognac lactone upon comparison of their spectral data and optical rotations ($[\alpha]_D +78^\circ$ (c=1.12, CH₂Cl₂) (lit.¹⁵) $[\alpha]_D +79.5^\circ$ (CH₂Cl₂)).

ACKNOWLEDGEMENT We are grateful to Dr. T. Ebata, Japan Tobacco Inc., for providing the spectral data of authentic samples. Thanks are also extended to the Ministry of Education, Science, and Culture (Japan) for a research grant.

REFERENCES

- 1) G. Ohloff, "Progress in the Chemistry of Organic Natural Products," Springer-Verlag, New York, Vol. 35, 1978; Tse-Lok Ho, *Synth. Commun.*, **13**, 341 (1983).
- 2) J. M. Brand, J. C. Joung, and R. M. Silverstein, "Progress in the Chemistry of Organic Natural Products," Springer-Verlag, New York, Vol. 37, 1979; K. Mori, T. Ebata, and S. Takechi, *Tetrahedron*, **40**, 1761 (1984).
- 3) D. L. Wood, L. E. Browne, B. Ewing, K. Lindahl, W. D. Bedard, P. E. Tilden, K. Mori, G. B. Pitman, and P. R. Hughes, *Science*, **192**, 896 (1976); L. Friedman and J. G. Miller, *Science*, **172**, 1044 (1971).
- 4) M. Masuda and K. Nishimura, *Chemistry Lett.*, 1981, 1333.
- 5) R. M. Ortuno, R. Merce, and J. Font, *Tetrahedron*, **43**, 4497 (1987).
- 6) K. Nishimura, *Chemistry Today*, **189**, 30 (1986).
- 7) W. Oppolzer, *Tetrahedron*, **43**, 1969 (1987).
- 8) O. Miyata, T. Shinada, T. Naito, and I. Ninomiya, *Chem. Pharm. Bull.*, **37**, 3158 (1989).
- 9) O. Miyata, T. Shinada, I. Ninomiya, T. Naito, T. Date, K. Okamura, and S. Inagaki, *J. Org. Chem.*, **56**, 6556 (1991).
- 10) O. Miyata, T. Shinada, I. Ninomiya, and T. Naito, *Tetrahedron Lett.*, **32**, 3519 (1991).
- 11) W. Oppolzer and A. J. Kingma, *Helv. Chim. Acta*, **72**, 1337 (1989).
- 12) M. Casey, A. C. Manage, and P. J. Murphy, *Tetrahedron Lett.*, **33**, 965 (1992).
- 13) P. Bravo, G. Resnati, F. Viani, and A. Arnone, *Tetrahedron*, **43**, 4635 (1987).
- 14) H. Matsuyama, T. Nakamura, and N. Kamigata, *J. Org. Chem.*, **54**, 5218 (1989); T. Nakamura, H. Matsuyama, M. Takahashi, and N. Kamigata, Abstract of Papers, The 18th Symposium on Heteroatom Chemistry, Chiba, Dec. 1990, p. 5.
- 15) T. Ebata, K. Matsumoto, H. Yoshikoshi, K. Koseki, H. Kawakami, and H. Matsushita, *Heterocycles*, **31**, 1585 (1990).

(Received July 3, 1992)