View Article Online View Journal



# Organic & Biomolecular Chemistry

## Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. Ikawa, H. Kaneko, S. Masuda, E. Ishitsubo, H. TOKIWA and S. akai, *Org. Biomol. Chem.*, 2014, DOI: 10.1039/C4OB01627K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/obc

# **RSCPublishing**

### Paper

Cite this: DOI: 10.1039/xoxxooooox

Received ooth January 2012, Accepted ooth January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/obc

# Trifluoromethanesulfonyloxy-Group-Directed Regioselective (3+2) Cycloadditions of Benzynes for the Synthesis of Functionalized Benzo-Fused Heterocycles

Takashi Ikawa,\*<sup>a</sup> Hideki Kaneko,<sup>a</sup> Shigeaki Masuda,<sup>a</sup> Erika Ishitsubo,<sup>b</sup> Hiroaki Tokiwa,<sup>b</sup> Shuji Akai\*<sup>a</sup>

Highly regioselective (3+2) cycloadditions of (trifluoromethanesulfonyloxy)benzynes [(triflyloxy)benzynes] with 1,3-dipoles followed by cross-coupling reactions provided multisubstituted benzo-fused heterocycles. The triflyloxy group at the 3-position of benzynes, and even that at the remote 4-position, greatly affected the regiocontrol of the cycloaddition. These groups also served to install other substituents at their *ipso*-positions.

#### Introduction

The (3+2) cycloaddition reactions of benzynes<sup>1</sup> have been useful in the construction of benzo-fused heterocyclic skeletons, which are widely found in various biologically active compounds.<sup>2,3</sup> However, the regioselectivities of (3+2) cycloadditions of unsymmetrically-substituted benzynes were often low, and the reactions provided difficult-to-separate regioisomer mixtures.<sup>4</sup> For instance, the reactions of pmethoxyphenyl azide (2a) with either 3-methyl- (3a) or 3fluorobenzyne (3b) afforded a 1:1 mixture of distal- and proximal-benzotriazoles (4a and 4b, respectively) (Scheme 1).<sup>4g</sup> This problem could be solved by using a specific directing group that has a strong effect on controlling the orientation of the cycloaddition reactions, and that can also be subsequently converted into any other desired substituent. An alkoxy group at the 3-position of benzynes has been widely utilized as a distinguished directing group of various benzyne reactions such as nucleophilic addition,<sup>5a</sup> cycloaddition<sup>4,5b</sup> and transition-metal-catalysis.<sup>5c</sup> In fact, the reactions of 3-methoxybenzyne 3c, generated from 1c, with 2a exclusively afforded the distal adduct, distal-4a (Scheme 1).4g However, the alkoxy groups on aryl rings have rarely been converted into other substituents after the reactions of alkoxybenzynes.

**Organic & Biomolecular Chemistry** 

Garg *et al.* reported sulfamoyloxy (OSO<sub>2</sub>NMe<sub>2</sub>)-groupcouplings after the cycloaddition, only a limited number of such



Scheme 1 Regioselectivities in (3+2) Cycloaddition of 3-Substituted Benzynes 3, Generated from 1, with Azide 2a.

directed regioselective reactions of 3,4-pyridyne.<sup>6</sup> Although this directing group could be transformed using nickel-catalyzed transformations are known and the substrate scopes of nickel-catalyzed couplings are much narrower than, for example, those of palladium.<sup>7</sup> Recently, we have reported silyl- or boryl-group-directed regioselective (3+2) cycloaddition reactions of benzynes.<sup>8</sup> These methods should be particularly valuable because the reactions of 3-silyl- and 3-borylbenzynes afforded products with complementary regioselectivities, and the directing groups were subsequently converted into various substituents under palladium-catalyzed and uncatalyzed conditions. The development of other directing groups of benzynes that have powerful regiocontrol and versatile convertibility should enrich diversity in synthetic methods and would be useful for organic and medicinal chemistries.

Herein, we report the use of a trifluoromethanesulfonyloxy (triflyloxy,  $OSO_2CF_3$ , OTf) group, at either the 3-position or 4-position of benzynes, in powerfully directed regioselective (3+2) cycloaddition reactions. Moreover, the OTf-group of the cycloadducts could be directly transformed into other substituents, such as aryl groups, using palladium-catalyzed couplings (Scheme 2).

<sup>&</sup>lt;sup>a</sup>Graduate School of Pharmaceutical Sciences, Osaka University, Yamadaoka, Suita, Osaka 565-0871, Japan. E-mails: ikawa@phs.osaka-u.ac.jp; akai@phs.osaka-u.ac.jp, Tel: +81-6-6879-8212; +81-6-6879-8210.

<sup>&</sup>lt;sup>b</sup>Department of Chemistry, Research for Smart Molecules, Rikkyo University, Nishi-Ikebukuro, Toshimaku, Tokyo 171-8501, Japan.

<sup>†</sup>Electronic Supplementary Information (ESI) available: Experimental *details*, full spectral data for all new compounds, distortion analysis, and Cartesian coordinate. see DOI: 10.1039/b000000x/

**Biomolecular Chemistry Accepted M** 



**Scheme 2** Proposed OTf-Group-Directed Regioselective Cycloaddition Followed by Subsequent Transformation.

#### **Results and discussion**

Considering the strong electron-withdrawing nature of a triflyloxy group, we expected that it may act as a more powerful directing group of benzyne cycloadditions than any alkoxy groups. Moreover, it is well known that the triflyloxy groups on aromatic compounds are readily convertible into other various functional groups.<sup>9</sup> However, there have been no reports on the use of the triflyloxy group as a directing group for benzyne reactions.

To examine the feasibility of our idea, a new benzyne precursor 8 (Table 1) bearing two triflyloxy and one tertbutyldimethylsilyl groups on a benzene ring was synthesized from resorcinol in 5 steps (see the Supplementary Information), while 1,3-bis(triflyloxy)-2-(trimethylsilyl)benzene could not be synthesized due the instability of 2to (trimethylsilyl)resorcinol.<sup>10</sup> Under mild conditions using CsF in MeCN at room temperature, novel benzyne 3d was successfully generated, which was confirmed by the subsequent (3+2)cycloaddition reaction with aryl azide 2a to form 1-(pmethoxyphenyl)-4-(triflyloxy)benzotriazole distal-4d (48% yield). To our delight, the regioselectivity of this reaction was exclusively distal (*distal:proximal* = >98:2; Entry 1). Under the same reaction conditions, benzyne 3d reacted with benzyl, pnitrobenzyl, and cyclohexyl azide 2b, 2c, and 2d, to afford distal-adducts 4e, 4f, and 4g, respectively (Entries 2-4), in superior yield to that of 4d. In all cases, the proximal-4 adducts were not observed by <sup>1</sup>H NMR analysis of crude reaction mixtures.

Generated from <b>8</b> , with Azides <b>2</b> . <sup><i>a</i></sup>							
OTf	Si( <i>t</i> -Bu)Me <sub>2</sub> CsF RN <sub>3</sub> 2 OTf MeCN rt, 3 h	OTf 3 2 1 + 3d	NH NH R 2 R distal-4 (L	N N R D-4) (P-4)	imal-4		
Entry	R	2	D-4 : P-4 <sup>b</sup>	4	Yield		
					(%) <sup>c</sup>		
1	C <sub>6</sub> H <sub>4</sub> -4-OMe	2a	>98:2	4d	48		
2	Bn	2b	>98:2	<b>4e</b>	74		
3	$CH_2C_6H_4$ -4- $NO_2$	2c	>98:2	<b>4f</b>	70		
4	cyclohexyl	2d	>98 : 2	4g	63		

Table 1 Regioselective (3+2) Cycloaddition of 3-(Triflyloxy)benzyne 3d,

<sup>*a*</sup>Conditions: **8** (1.0 equiv), **2** (3.0 equiv), CsF (3.0 equiv), MeCN, rt. <sup>*b*</sup>Determined by <sup>1</sup>H NMR analysis of a crude product. <sup>*c*</sup>Isolated yield of *distal*-**4**. The reactions of **3d** generated from **8** with nitrile oxides (**9a** and **9b**)<sup>11</sup> were also achieved to provide proximal adducts (*proximal*-**10a** and -**10b**) exclusively (Scheme 3). Notably, the selectivities of the reactions using **3d** are much higher than that using 3-methoxybenzyne (*proximal:distal* = 80:20).<sup>4h,12</sup> These results suggest that the selectivities are derived not only from the electron-negativity of the oxygen atom but also from the strong electron-withdrawing nature of the sulforyl group.



Next, we attempted the remote regiocontrol of (3+2)cycloadditions using substituents at the 4-position of benzyne, which has been long known as a much more difficult task than that using substituents at the adjacent 3-position.<sup>13</sup> Harrity et al. reported the directing effects of Br, CO<sub>2</sub>Me, and CN groups attached at the 4-position of benzynes on the (3+2) cycloaddition with benzyl azide, which resulted in the production of two regioisomers in up to 2:1 ratio although the transformations of the directing groups have never been tried.<sup>13f</sup> Not surprisingly, the (3+2) cycloaddition of 4-methoxybenzyne 3f, generated from precursor 11a, with benzyl azide 2b afforded a 52:48 mixture of two regioisomers (distal-12a:proximal-12a) (Table 2, Entry 1). On the other hand, a methanesulfonyloxy group in 3g had a greater effect on the regiocontrol of the cycloaddition to afford a 72:28 mixture of distal-12b:proximal-12b (Entry 2). The regioselectivity became slightly better when using a triflyloxy group as the directing

**Table 2** Regioselectivities in the (3+2) Cycloaddition of 4-Substituted Benzynes **3**, Generated from **11**, with Azide **2b**.<sup> $\sigma$ </sup>

x	SiMe OTf 11	CsF <sup>3</sup> BnN <sub>3</sub> 2t MeCN rt, 3 h		4 2 1 3 2 N N Bn 2b -	X distal-	$ \overset{N}{\longrightarrow} \overset{N}{\underset{N}{\underset{N}{\overset{N}{\underset{N}{\underset{N}{\overset{N}{\underset{N}{\underset{N}{\overset{N}{\underset{N}}{\underset{N}{N$
Entry	Х	11	3	<i>D</i> -12 : <i>P</i> -12 <sup>b</sup>	12	Yield
						$(\%)^{c}$
1	OMe	11a	3f	52:48	12a	80
2	OMs	11b	3g	72:28	12b	65
3	OTf	11c	3e	77:23	12c	47
$4^d$	OTf	11c	3e	75:25	12c	60

<sup>*a*</sup>Conditions: **11c** (1.0 equiv), 12 equiv of **2** (12 equiv), CsF (1.0 equiv), MeCN, rt. <sup>*b*</sup>Isolated product ratio. <sup>*c*</sup>Total isolated yield of *proximal*-**12** and *distal*-**12**.

Published on 22 October 2014. Downloaded by UNIVERSITY OF ALABAMA AT BIRMINGHAM on 26/10/2014 06:25:48.

Published on 22 October 2014. Downloaded by UNIVERSITY OF ALABAMA AT BIRMINGHAM on 26/10/2014 06:25:48.

Journal Name

group (*distal*-12c:*proximal*-12c = 77:23; Entry 3). The lower 47% yield of 12c was improved to 60% by using 12 equiv of azide 2b while maintaining similar regioselectivity (Entry 4). Delightfully, the triflyloxy group, which exhibited the highest regioselectivity as a directing group, would be the most promising for the subsequent transformations (*vide infra*, see Scheme 4).<sup>9</sup>

With the optimized benzyne precursor **11c** and reaction conditions in hand, azide **2a**, along with **2c** and **2d**, were applied to the (3+2) cycloaddition reactions (Table 3). Aryl azide **2a** (Entry 1), primary alkyl azide **2c** (Entry 2), and secondary alkyl azide **2d** (Entry 3) successfully reacted with the *in situ* generated 4-(triflyloxy)benzyne **3e** to preferentially produce distal products (*distal*-**12**:*proximal*-**12** = 75–76:25–24). To the best of our knowledge, these are the best results among the reported regioselectivities of 4-substituted benzyne (3+2) cycloadditions.

Table 3 Regioselective (3+2) Cycloaddition of 4-(Triflyloxy)benzyne 3e, Generated from 11c, with Azides 2.<sup>a</sup>

TfO	SiMe <sub>3</sub> CsF RN <sub>3</sub> 2 OTf MeCN rt, 3 h	- TfO_4 3	$\begin{bmatrix} 2 & N \\ + & N^+ \\ 1 & -N \\ R & R \\ 2 \end{bmatrix}$	distal-12 TfO	$ \begin{array}{c} & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & $
	_			proxima	12 (P-12)
Entry	R	2	<i>D</i> -12 : <i>P</i> -12 <sup><i>b</i></sup>	12	Yield $(\%)^c$
1	C <sub>6</sub> H <sub>4</sub> -4-OMe	2a	76 : 24	12d	61
2	CH <sub>2</sub> C <sub>6</sub> H <sub>4</sub> -4-NO <sub>2</sub>	2c	75 : 25	12e	53
3	cyclohexyl	2d	76 : 24	12f	60

<sup>a</sup>Conditions: **11c** (1.0 equiv), 12 equiv of **2** (12 equiv), CsF (1.0 equiv), MeCN, rt. <sup>b</sup>Isolated product ratio. <sup>c</sup>Total isolated yield of *proximal*-**12** and *distal*-**12**.

Aryl trifluoromethanesulfonates have widely been used as good substrates for palladium-catalyzed coupling reactions.<sup>7</sup> The major cycloaddition products, *distal*-4e, *distal*-12c, and *proximal*-10a, were applied to the Suzuki coupling reactions with boronic acids 13a and 13b (Scheme 4). Reactions in the presence of a palladium catalyst, ligand, boronic acid, and base smoothly afforded biaryls 14–16 in 75–88% isolated yields even under unoptimized reaction conditions. These results suggest that the combination of the regioselective (3+2) cycloadditions of 3- or 4-(triflyloxy)benzynes and 1,3-dipoles, with subsequent transformations of the triflyloxy group provides a useful method for the synthesis of multisubstituted benzo-fused heterocyclic compounds.

The origin of the regioselectivity in the (3+2) cycloadditions was analyzed using density functional theory (DFT) calculation<sup>14</sup> and a natural bond orbital (NBO) method.<sup>15</sup> Firstly, the structures of **3c–f** were optimized by DFT [B3LYP/6-31G(d)], and then the electron densities of their reacting  $\pi$ -orbitals were calculated by NBO 6.0.<sup>16</sup> Thereby, the



Scheme 4 Transformations of Cycloaddition Products, *distal*-4e, *distal*-12c, and *proximal*-10a.

selectivities of both 3- and 4-substituted benzynes could be rationally explained using the electron densities of the NBO (Figure 1).<sup>15–20</sup> The electron density in the  $\pi$ -orbital at the C1 of 3c (0.826) and that of 3d (0.781) were lower than those at the C2 (1.010 and 1.031, respectively). A similar tendency was observed for 3f and 3e; however, the difference between the electron densities of 3f (C2–C1 = 0.0005) and that of 3e (C2– C1 = 0.046) were found to be much smaller than those of **3c** and 3d (C2-C1 = 0.184 and 0.250, respectively). These calculated differences were parallel with the magnitudes of regioselectivities. For example, the (3+2) cycloadditions of 3c and 3d with benzyl azide 2b exhibited complete distal regioselectivities (Scheme 1 and Table 2, Entry 2), while that of 3e afforded moderate distal selectivity. Moreover, very poor selectivity was observed using 3f. These results indicated that the inductive effect of the oxygen atom at the 3-position was greater than that at the 4-position, thus inducing a larger difference in electron density between the C1 and C2 orbitals, which resulted in higher regioselectivities. An additional sulfonyl group at the 4-position could amplify the electronwithdrawing inductive effect and achieve directed benzyne (3+2) cycloaddition reactions.



**Fig. 1** Natural Bond Orbital (NBO) Analysis of Substituted Benzynes **3**c–f. Optimized structures of benzynes **3**c–f were calculated by DFT [B3LYP6/31G(d)]. Electron densities of the reacting *p*-orbitals were determined by NBO analysis.

#### Conclusions

In conclusion, the regiocontrolled (3+2) cycloadditions of benzynes with azides and nitrile oxides were demonstrated by the use of a triflyloxy group at either the 3- or 4-position of benzynes. In particular, this method provides the highest reported selectivity for 4-substituted-group-directed benzyne cycloaddition reactions. The sequential combination of regioselective cycloaddition and palladium-catalyzed coupling of the aryl triflates could be implemented in the synthesis of polysubstituted heterocycles. The applications of this chemistry to a wide range of benzyne reactions and detailed mechanistic investigations are now underway in our laboratory.

#### Experimental

#### General Procedure for Regioselective Cycloadditions of Triflyloxybenzynes 3 with dipoles (Tables 1–3, and Scheme 3)

CsF (1.5 equiv) was flame-dried under reduced pressure in a flask equipped with a three-way stopcock, and back-filled with Ar. Azide 2 or nitrile oxide 9 (7.0 equiv) with a stir bar was loaded into the flask and evacuated and backfilled with Ar (This process was repeated three times). MeCN (One-fifth of its total volume) was added into the flask via a syringe. A solution of precursor 8, or 11 (1.0 equiv) in anhydrous MeCN (one-fifth of its total volume) was added to the flask through a cannula and washed with MeCN (three-fifth of its total volume). The mixture was stirred at rt for the period shown in Tables 2-4 and Scheme 3. H<sub>2</sub>O and EtOAc were added to the reaction mixture, and the aqueous phase was extracted thrice with EtOAc. The combined organic phase was washed with a saturated aqueous NaCl solution (brine). The organic phase was dried over anhydrous MgSO<sub>4</sub>, and the solvent was removed under reduced pressure. The residue was subjected to <sup>1</sup>H NMR analysis for calculating the ratio of the two regioisomers (distal- and proximal-4, 10, or 12). The crude product was purified by flash column chromatography on silica gel (hexane, a mixture of hexane and EtOAc, or CH<sub>2</sub>Cl<sub>2</sub>) to afford distal- and proximal-4, 10, or 12.

#### General Procedure for Pd-Catalyzed Couplings of Cycloaddition Products *distal*- and *proximal*-4, 10, or 12

An oven dried Schlenk tube was charged with *distal*- and *proximal*-4, 10, or 12 (1.0 equiv), phenylboronic acid 13 (1.5 equiv), Pd source (0.10 equiv), ligand (0.20 equiv, if applicable), base (2.0 equiv), evacuated and back-filled with Ar. An anhydrous solvent (0.56 mL) was added via syringes, and the reaction mixture was stirred at 100 °C for several hours and filtered through a pad of silica gel cake using EtOAc. The eluent was concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc) to provide the biaryl compound 14–16.

#### Synthesis of benzyne precursors 8, 11b and 11c

**2-Bromoresorcinol (17):**<sup>22,23</sup> To a solution of resorcinol (11 g, 0.10 mol) in CHCl<sub>3</sub> (63 mL, 0.50 M) was added Br<sub>2</sub> (15 mL, 0.30 mol) at 0 °C. After stirring for 10 h at rt, the mixture was concentrated in vacuo. The residue was recrystallized from CHCl<sub>3</sub> to give 2,4,6-tribromoresorcinol (27 g, 77%). To a solution of 2,4,6-tribromoresorcinol (17 g, 50 mmol) in H<sub>2</sub>O/MeOH (0.35 L, H<sub>2</sub>O/MeOH = 6:1, 0.50 M) were added

NaOH (15 mL, 0.30 mol) and Na<sub>2</sub>SO<sub>3</sub> at rt. After stirring for 10 h at rt, the mixture was concentrated in vacuo. The residue was recrystallized from CHCl<sub>3</sub> to give the titled compound **17** (7.3 g, 78%) as a colourless solid. Mp: 101–102 °C (Lit. 101–102 °C).<sup>23 1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$ : 5.39 (2 OH, s), 6.60 (2 H, d, J = 8.5 Hz), 7.11 (1 H, t, J = 8.5 Hz).

1,3-Bis(tert-butyldimethylsilyloxy)-2-bromobenzene (18): To a solution of 17 (6.0 g, 32 mmol) in DMF (63 mL, 0.50 M) were added imidazol (6.5 g, 95 mmol) and TBSCl (14 g, 95 mmol) at 0 °C. After stirring for 1 h at rt, the reaction was stopped by adding a saturated aqueous solution of NH<sub>4</sub>Cl and the mixture was extracted with hexane. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc = 20:1) to provide the titled compound 18 (13 g, quant) as a colourless solid. Mp: 40-42 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ: 0.23 (12 H, s), 1.04 (18 H, s), 6,51 (2 H, d, *J* = 8.5 Hz), 6.99 (1 H, t, *J* = 8.5 Hz). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ: -4.22, 18.4, 25.8, 109.3, 113.0, 127.3, 154.1. IR (neat): 1252, 1464 cm<sup>-1</sup>. HRMS (MALDI) Calcd for  $C_{18}H_{34}O_2Si_2Br [M+H]^+$ : 417.1275, found 417.1257.

#### 2-(tert-Butyldimethylsilyl)-3-[(tert-

butyldimethylsilyl)oxy|phenol (19): To a solution of 18 (10 g, 24 mmol) in THF (0.12 L, 0.20 M) was added 1.6 M n-BuLi in hexane (18 mL, 29 mmol) slowly at -78 °C. After stirring for 40 min, the reaction was stopped by adding a saturated aqueous solution of NH<sub>4</sub>Cl and the mixture was extracted with hexane. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc = 15:1) to provide the titled compound **19** (8.1) g, 85%) as a colourless solid. Mp: 57-60 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ: 0.31 (6 H, s), 0.37 (6 H, s), 0.93 (9 H, s), 1.00 (9 H, s), 4.87 (OH, s), 6.30 (1 H, d, J = 8.0 Hz), 6.40 (1 H, d, J = 8.0 Hz), 7.05 (1 H, t, J = 8.0 Hz). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ: -2.95, -1.37, 18.6, 19.5, 26.9, 27.0, 107.8, 110.5, 112.0, 130.6, 162.3, 162.3. IR (neat): 1254, 1437, 3512 cm<sup>-1</sup> HRMS (MALDI) Calcd for  $C_{18}H_{35}O_2Si_2$  [M+H]<sup>+</sup>:339.2170, found 339.2170.

2-(*tert*-Butyldimethylsilyl)benzene-1,3-diol (20): То а solution of 19 (2.0 g, 5.0 mmol) in THF (50 mL, 0.10 M) was added TBAF (5.0 mL, 5.0 mmol) slowly at 0 °C. After stirring for 1 h, the reaction was stopped by adding a saturated aqueous solution of NH<sub>4</sub>Cl and the mixture was extracted with EtOAc. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo. The residue was recrystallized from CHCl<sub>3</sub> to provide the titled compound 20 (0.86 mg, 77%) as a colourless solid. Mp: 128–131 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ: 0.41 (6 H, s), 0.94 (9 H, s), 4.90 (2 OH, s), 6.29 (2 H, d, J = 8.0 Hz), 7.07 (1 H, t, J = 8.0 Hz). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ: -2.09, 18.4, 26.8, 107.7, 108.0, 131.4, 162.1. IR (neat): 1263, 1327, 1445, 3518 cm<sup>-1</sup>. HRMS (MALDI) Calcd for  $C_{12}H_{21}O_2Si [M+H]^+$ : 225.1305, found 225.1298.

#### 2-(tert-Butyldimethylsilyl)-1,3-

**bis(trifluoromethanesulfonyloxy)benzene (8):** To a solution of **20** (0.20 g, 0.89 mmol) in  $CH_2Cl_2$  (4.5 mL, 0.20 M) were added DIPEA (0.47 mL, 2.7 mmol) and  $Tf_2O$  (0.45 mL, 2.7 mmol) at 0 °C. After stirring for 1 h at rt, the reaction was

Journal Name

stopped by adding NaHCO<sub>3</sub> aq. and the mixture was extracted with EtOAc. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc = 20:1) to provide the titled compound **8** (0.37 g, 86%) as a yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ : 0.53 (6 H, s), 0.98 (9 H, s), 7.49 (2 H, d, J = 9.0 Hz), 7.56 (1 H, t, J = 9.0 Hz). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ : -2.00, 18.6, 26.5, 117.9, 118.6 (q, J = 318 Hz), 122.4, 132.3, 156.0. <sup>19</sup>F NMR (280 MHz, CDCl<sub>3</sub>)  $\delta$ : -73.8. IR (neat): 1215, 1424 cm<sup>-1</sup>. Anal. Calcd for C<sub>14</sub>H<sub>18</sub>F<sub>6</sub>O<sub>6</sub>S<sub>2</sub>Si: C, 34.42; H, 3.71. Found: C, 34.54; H, 3.72.

**2-Bromohydroquinone** (21):<sup>24,25</sup> To a solution of hydroquinone (6.4 g, 58 mmol) in CHCl<sub>3</sub> (0.29 L, 0.20 M) was added Br<sub>2</sub> (3.0 mL, 58 mmol) at 0 °C. After stirring for 1 h at rt, the mixture was concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc = 3:1) to provide the titled compound **21** (6.0 g, 55%) as a colourless solid. Mp: 111–114 °C (Lit. 112 °C).<sup>25 1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$ : 5.12 (2 OH, brs), 6.73 (1 H, dd, *J* = 3.0, 9.0 Hz), 6.90 (1 H, d, *J* = 9.0 Hz), 6.99 (1 H, d, *J* = 3.0 Hz).

**1,4-Bis(trimethylsilyloxy)-2-bromobenzene (22):** To a solution of **21** (2.5 g, 13 mmol) in THF (65 mL, 0.20 M) were added  $Et_3N$  (5.4 mL, 39 mmol) and TMSCl (4.9 mL, 39 mmol). After stirring for 1 h at rt, the mixture was concentrated in vacuo. The residue was filtered through Celite pad (washed with hexane) and concentrated in vacuo as a colourless oil (4.3 g, quant). This compound **22** was used for next reaction without purification due to the instability on silica gel column chromatography.

2-(Trimethylsilyl)hydroquinone (23):<sup>26</sup> To a solution of 22 (4.3 g, 13 mmol) in THF (65 mL, 0.20 M) was added 2.3 M n-BuLi in hexane (11 mL, 26 mmol) slowly at -78 °C. After stirring for 1 h at rt, the reaction was stopped by adding a saturated aqueous solution of NH4Cl and the mixture was extracted with EtOAc. The combined organic extracts were washed with brine, dried over MgSO4 and concentrated in The residue was purified by flash column vacuo. chromatography on silica gel (hexane/EtOAc = 5:1) to provide the titled compound **23** (2.1 g, 89%) as a colourless solid. Mp: 126–127 °C (Lit. 126–127 °C).<sup>26 1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ: 0.30 (9 H, s), 4.49 (OH, s), 4.54 (OH, s), 6.57 (1 H, d, J = 8.5 Hz), 6.70 (1 H, dd, J = 3.5, 8.5 Hz), 6.82 (1 H, d, J = 3.5 Hz). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ: -1.08, 115.5, 116.2, 117.0, 121.4, 149.0, 154.2. IR (neat): 1362, 3349 cm<sup>-1</sup>. HRMS (MALDI) Calcd for  $C_9H_{14}O_2Si [M+H]^+$ : 182.0758, found 182.0759.

#### 1-(Methanesulfonyloxy)-3-(trimethylsilyl)-4-

(trifluoromethanesulfonyloxy)benzene (11b): To a solution of 23 (0.90 g, 4.9 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (25 mL, 0.20 M) were added pyridine (2.6 mL, 32 mmol) and MsCl (1.9 mL, 25 mmol) at 0 °C. After stirring for 2 h at rt, the reaction was stopped by adding H<sub>2</sub>O and the mixture was extracted with Et<sub>2</sub>O. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (toluene/EtOAc = 6:1) to provide the mixture of 11b and 1,4bis(methansulfonyloxy)-3-(trimethylsilyl)benzene (1.3 g). To a solution of the mixture in CH<sub>2</sub>Cl<sub>2</sub>(14 mL) were added pyridine (1.9 mL, 24 mmol) and Tf<sub>2</sub>O (3.1 mL, 18 mmol) at 0 °C. After stirring for 3 h at rt, the reaction was stopped by adding H<sub>2</sub>O and the mixture was extracted with Et<sub>2</sub>O. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc = 5:1) to provide the titled compound **11b** [1.0 g, 52% (2 steps)] as a yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ : 0.38 (9 H, s), 3.18 (3 H, s), 7.36 (1 H, dd, *J* = 3.0, 8.5 Hz), 7.21 (1 H, d, *J* = 8.5 Hz), 7.40 (1 H, d, *J* = 3.0 Hz). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ : -1.21, 37.6, 118.3 (q, *J* = 317 Hz), 121.2, 124.5, 129.3, 135.9, 147.5, 152.7. <sup>19</sup>F NMR (470 MHz, CDCl<sub>3</sub>)  $\delta$ : -73.8. IR (neat): 1373, 1420 cm<sup>-1</sup>. HRMS (APCI) Calcd for C<sub>11</sub>H<sub>16</sub>F<sub>3</sub>O<sub>6</sub>S<sub>2</sub>Si [M+H]<sup>+</sup>: 393.01097, found 393.01356.

#### 1,4-Bis(trifluoromethanesulfonyloxy)-3-

(trimethylsilyl)benzene (11c): To a solution of 23 (1.0 g, 5.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (28 mL, 0.20 M) were added pyridine (2.0 mL, 25 mmol) and Tf<sub>2</sub>O (2.8 mL, 17 mmol) at 0 °C. After stirring for 19 h at rt, the reaction was stopped by adding a saturated aqueous solution of NH<sub>4</sub>Cl and the mixture was extracted with Et<sub>2</sub>O. The combined organic extracts were washed with brine, dried over MgSO4 and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (hexane/EtOAc = 15:1) to provide the titled compound **11c** (2.3 g, 95%) as a yellow oil. <sup>1</sup>H NMR (300 MHz,  $\hat{CDCl}_3$ )  $\delta$ : 0.39 (9 H, s), 7.34 (1 H, dd, J = 3.0, 9.0Hz), 7.38 (1 H, d, J = 3.0 Hz), 7.44 (1 H, d, J = 9.0 Hz). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$ : -1.21, 118.4 (q, J = 318 Hz), 118.7 (q, J = 319 Hz), 121.5, 123.9, 128.6, 136.7, 147.9, 153.3. <sup>19</sup>F NMR (280 MHz, CDCl<sub>3</sub>)  $\delta$ : -73.7, -72.6. IR (neat): 1427 cm<sup>-1</sup>. HRMS (APCI) Calcd for  $C_{12}H_{13}F_6O_6S_2Si [M+H]^+$ : 446.98270, found 446.98508.

#### Acknowledgements

This work was supported by the JSPS KAKENHI (Grant Numbers 23790017 and 25460018), Platform for Drug Discovery, Informatics, and Structural Life Science from the MEXT and Hoansha foundation.

#### **Notes and references**

- For selected reviews on benzynes, see: (a) A. M. Dyke, A. J. Hester and G. C. Lloyd-Jones, *Synthesis*, 2006, 24, 4093–4112; (b) R. Sanz, *Org. Prep. Proced. Int., 2008*, 40, 215–291; (c) H. Yoshida, J. Ohshita and A. Kunai, *Bull. Chem. Soc. Jpn.*, 2010, 83, 199–219; (d) P. M. Tadross and B. M. Stoltz, *Chem. Rev.*, 2012, 112, 3550–3577; (e) C. Wu and F. Shi, *Asian J. Org. Chem.*, 2013, 2, 116–125; (f) R. W. Hoffmann and K. Suzuki, *Angew. Chem. Int. Ed.*, 2013, 52, 2655–2656; (g) C. Holden and M. F. Greaney, *Angew. Chem. Int. Ed.*, 2014, 53, 5746–5749.
- 2 (a) A. R. Katrizky and A. F. Pozharskii, *Handbook of Heterocyclic Chemistry*, 2nd ed., Pergamon: New York, 2000; (b) T. Eicher and S. Hauptmann, *The Chemistry of Heterocycles*, Wiley-VCH: Weinheim, 2003.
- 3 (a) G. Semple, P. J. Skinner, M. C. Cherrier, P. J. Webb, C. R. Sage, S. Y. Tamura, R. Chen, J. G. Richman and D. T. Connolly, *J. Med. Chem.*, 2006, **49**, 1227–1230; (b) S. Hayashi, A. Hirao, A. Imai, H. Nakamura, Y. Murata, K. Ohashi and E. Nakata, *J. Med. Chem.*, 2009, **52**, 610–625; (c) Z. Rezaei, S. Khabnadideh, K. Pakshir, Z.

Page 6 of 7

Published on 22 October 2014. Downloaded by UNIVERSITY OF ALABAMA AT BIRMINGHAM on 26/10/2014 06:25:48.

Hossaini, F. Amiri and E. Assadpour, *Eur. J. Med. Chem.*, 2009, 44, 3064–3067; (d) D. Patel, M. Jain, S. R. Shah, R. Bahekar, P. Jadav, B. Darji, Y. Siriki, D. Bandyopadhyay, A. Joharapurkar, S. Kshirsagar, H. Patel, M. Shaikh, K. V. V. M. Sairam and P. Patel, *ChemMedChem*, 2011, 6, 1011–1016; (e) A. Dubey, S. K. Srivastava and S. D. Srivastava, *Bioorg. Med. Chem. Lett.*, 2011, 21, 569–573.

- 4 (a) T. Matsumoto, T. Sohma, S. Hatazaki and K. Suzuki, Synlett, 1993, 843-846; (b) T. Jin and Y. Yamamoto, Angew. Chem. Int. Ed., 2007, 46, 3323-3325; (c) F. Shi, J. P. Waldo, Y. Chen and R. C. Larock, Org. Lett., 2008, 10, 2409-2412; (d) Z. Liu, F. Shi, P. D. G. Martinez, C. Raminelli and R. C. Larock, J. Org. Chem., 2008, 73, 219-226; (e) M. Dai, Z. Wang and S. J. Danishefsky, Tetrahedron Lett., 2008, 49, 6613-6616; (f) L. Campbell-Verduyn, P. H. Elsinga, L. Mirfeizi, R. A. Dierckx and B. L. Feringa, Org. Biomol. Chem., 2008, 6, 3461-3463; (g) F. Zhang and J. E. Moses, Org. Lett., 2009, 11, 1587-1590; (h) J. A. Crossley and D. L. Browne, Tetrahedron Lett., 2010, 51, 2271-2273; (i) C. Spiteri, C. Mason, F. Zhang, D. J. Ritson, P. Sharma, S. Keeling and J. E. Moses, Org. Biomol. Chem., 2010, 8, 2537-2542; (j) C. Spiteri, S. Keeling and J. E. Moses, Org. Lett., 2010, 12, 3368-3371; (k) A. V. Dubrovskiy and R. C. Larock, Org. Lett., 2010, 12, 1180-1183; (I) P. Li, J. Zhao, C. Wu, R. C. Larock and F. Shi, Org. Lett., 2011, 13, 3340-3343; (m) Y. Fang, C. Wu, R. C. Larock and F. Shi, J. Org. Chem., 2011, 76, 8840-8851; (n) C. Lu, A. V. Dubrovskiy and R. C. Larock, J. Org. Chem., 2012, 77, 2279-2284; (o) A. V. Dubrovskiy, P. Jain, F. Shi, G. H. Lushington, C. Santini, P. Porubsky and R. C. Larock, ACS Comb. Sci., 2013, 15, 193-201.
- 5 For selected reviews on specific benzyne reactions, see: (a) *Comprehensive Organic Synthesis*, Vol. 4 (Eds.: B. M. Trost, I. Fleming and M. F. Semmelhack), Pergamon, Oxford, 1992, pp. 483– 515; (b) A. V. Dubrovskiy, N. A. Markina and R. C. Larock, *Org. Biomol. Chem.*, 2013, **11**, 191–218; (c) S. A. Worlikar and R. C. Larock, *Curr. Org. Chem.*, 2011, **15**, 3214–3232.
- 6 (a) A. E. Goetz and N. K. Garg, *Nature Chem.*, 2013, 5, 54–60; (b) A. E. Goetz and N. K. Garg, *J. Org. Chem.*, 2014, 79, 846–851.
- 7 (a) S. D. Ramgren, A. L. Silberstein, Y. Yang and N. K. Garg, *Angew. Chem. Int. Ed.*, 2011, **50**, 2171–2173; (b) T. K. Macklin and V. Snieckus, *Org. Lett.*, 2005, **7**, 2519–2522; (c) A. G. Sergeev and J. F. Hartwig, *Science*, 2011, **332**, 439–443; (d) T. Mesganaw, N. F. Fine Nathel and N. K. Garg, *Org. Lett.*, 2012, **14**, 2918–2921.
- 8 (a) T. Ikawa, H. Tokiwa and S. Akai, *J. Synth. Org. Chem. Jpn.*, 2012, **70**, 1123–1133; (b) T. Ikawa, A. Takagi, M. Goto, Y. Aoyama, Y. Ishikawa, Y. Itoh, S. Fujii, H. Tokiwa and S. Akai, *J. Org. Chem.*, 2013, **78**, 2965–2983.
- 9 For selected reviews on cross-coupling of aryl triflates, see: (a) K. Ritter, *Synthesis*, 1993, 735–762; (b) I. Baraznenok, V. Nenajdenko and E. Balenkova, *Tetrahedron*, 2000, **56**, 3077–3119; (c) Z. Gilson and R. C. Larock, *Chem. Rev.*, 2006, **106**, 4644–4680.
- 10 For a related benzyne precursor, see: Y. Himeshima, T. Sonoda and H. Kobayashi, *Chem. Lett.*, 1983, 8, 1211–1212.
- (a) J. W. Bode, Y. Hachisu, T. Matsuura and K. Suzuki, *Tetrahedron Lett.*, 2003, 44, 3555–3558; (b) J. W. Bode, Y. Hachisu, T. Matsuura and K. Suzuki, *Org. Lett.*, 2003, 5, 391–394; (c) T. Matsuura, J. W. Bode, Y. Hachisu and K. Suzuki, *Synlett*, 2003, 1746–1748.

- 12 The reported selectivity<sup>4h</sup> was also reproduced under the same reaction conditions using CsF in MeCN at rt (see the Supplementary Information).
- 13 For selected examples of 4-substituted-group-directed benzyne reaction, see: (a) H. Yoshida, Y. Honda, E. Shirakawa and T. Hiyama, *Chem. Commun.*, 2001, 1880–1881; (b) H. Yoshida, H. Fukushima, J. Ohshita and A. Kunai, *J. Am. Chem. Soc.*, 2006, **128**, 11040–11041; (c) Z. Liu and R. C. Larock, *J. Org. Chem.*, 2006, **71**, 3198–3209; (d) C. Raminelli, Z. Liu and R. C. Larock, *J. Org. Chem.*, 2006, **71**, 4689–4691; (e) H. Yoshida, T. Morishita, H. Fukushima, J. Ohshita and A. Kunai, *Org. Lett.*, 2007, **9**, 3367–3370; (f) J. D. Kirkham, P. M. Delaney, G. J. Ellames, E. C. Row and J. P. A. Harrity, *Chem. Commun.*, 2010, **46**, 5154–5156; (g) E. Yoshioka and H. Miyabe, *Tetrahedron*, 2012, **68**, 179–189; (h) H. Yoshida, R. Yoshida and K. Takaki, *Angew. Chem. Int. Ed.*, 2013, **52**, 8629–8632.
- 14 All calculations were performed using the Gaussian 09 program package (revision D.01): M. J. Frisch *et al.* Gaussian, Inc.: Wallingford CT, 2009. For details, see the Supplementary Information.
- 15 For a natural bond orbital (NBO) method, see: (a) K. B. Wiberg, *Tetrahedron*, 1968, 24, 1083–1096; (b) J. P. Foster and F. Weinhold, *J. Am. Chem. Soc.*, 1980, 102, 7211–7218; (c) A. E. Reed, R. B. Weinstock and F. Weinhold, *J. Chem. Phys.*, 1985, 83, 735–746.
- 16 For NBO6, see: E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales and F. Weinhold, Theoretical Chemistry Institute, University of Wisconsin, Madison, 2013.
- 17 The natural "localized" molecular orbital (NLMO) based on the natural bond orbital (NBO) interpretations could give chemists actual chemical and even quantitative insights without methodology/basis set-dependency. For a review on NBO analysis, see: A. E. Reed, L. A. Curtiss, and F. Weinhold, *Chem. Rev.*, 1988, **88**, 899–926.
- 18 For recent reports on the NBO analysis of benzynes, see: (a) A. Takagi, T. Ikawa, Y. Kurita, K. Saito, K. Azechi, M. Egi, Y. Itoh, H. Tokiwa, Y. Kita and S. Akai, *Tetrahedron*, 2013, **69**, 4338–4352; (b) A. Takagi, T. Ikawa, K. Saito, S. Masuda, T. Ito and S. Akai, *Org. Biomol. Chem.*, 2013, **11**, 8145–8150.
- 19 Electron density  $\rho_{C_A}^i$  of the *i*th (reacting  $\pi$ ) NBO of carbon atom C<sub>A</sub> (A = 1 or 2) was evaluated as follows,

$$\rho^{i}_{C_{\lambda}} = n_{i} \times d_{C_{\lambda}} \qquad (1$$

where  $n_i$  is occupancy of the *i*th NBO and  $d_{C_A}$  is percentage contribution from each carbon atom C<sub>A</sub> for the *i*th NBO (Fig. 1).

- 20 Distortion analysis<sup>21</sup> could also rationalize the regioselectivities of (3+2) cycloaddition of 3-substituted benzynes **3c** and **3d**; however, this method could not be applied to the reactions of 4-substituted benzynes **3e** and **3f** (see the Supplementary Information).
- 21 Recently, Houk and Garg *et al.* have proposed that distortions of arynes are useful to predict and understand the regioselectivities of aryne reactions, see: (a) P. H.-Y. Cheong, R. S. Paton, S. M. Bronner, G-Y. J. Im, N. K. Garg and K. N. Houk, *J. Am. Chem. Soc.*, 2010, 132, 1267–1269; (b) G-Y. Im, S. M. J. Bronner, A. E. Goetz, R. S. Paton, P. H.-Y. Cheong, K. N. Houk and N. K. Garg, *J. Am. Chem. Soc.*, 2010, 132, 17933–17944; (c) S. M. Bronner, J. L. Mackey, K.

## Page 7 of 7

Published on 22 October 2014. Downloaded by UNIVERSITY OF ALABAMA AT BIRMINGHAM on 26/10/2014 06:25:48.

Journal Name

N. Houk and N. K. Garg, J. Am. Chem. Soc., 2012, 134, 13966-13969.

- 22 E. Kiehlmann and R. W. Lauener, Can. J. Chem., 1989, 67, 335–344.
- 23 U. Lüning, M. Abbass and F. Fahrenkrug, *Eur. J. Org. Chem.*, 2002, 3294–3303.
- 24 G. Viault, D. Grée, S. Das, J. S. Yadav and R. Grée, *Eur. J. Org. Chem.*, 2011, 1233–1241.
- 25 D. A. M. Egbe, C. Bader, E. Klemn, L. Ding, F. E. Karasz, U. W. Grummt and E. Birckner, *Macromolecules*, 2003, 36, 9303–9312.
- 26 G. Cooper, B. Williams and C. Lape, J. Org. Chem., 1961, 26, 4171– 4173.