

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. M. Levesque, R. G. Kinney and B. A. Arndtsen, *Chem. Sci.*, 2020, DOI: 10.1039/D0SC00085J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

A Palladium-Catalyzed C-H Functionalization Route to Ketones via the Oxidative Coupling of Arenes with Carbon Monoxide

Taleah M. Levesque, R. Garrison Kinney and Bruce A. Arndtsen*

Received 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

We describe the development of a new palladium-catalyzed method to generate ketones via the oxidative coupling of two arenes and CO. This transformation is catalyzed by simple palladium salts, and is postulated to proceed via the conversion of arenes into high energy aryl triflate electrophiles. Exploiting the latter can also allow the synthesis of unsymmetrical ketones from two different arenes.


Introduction

Metal-catalyzed arene C-H bond functionalization has emerged as a powerful technique for the efficient and atom-economical build-up of organic products from feedstock chemicals.¹ One important class of these reactions is to form carbon-carbon bonds from the oxidative coupling of two separate C-H bond containing substrates.² In contrast to classical synthetic methods employing pre-functionalized substrates (e.g. cross-coupling reactions), this strategy offers a platform to assemble biaryls and related structures directly from arenes with high step efficiency and minimal waste (Fig. 1a).³ A potentially useful variant would be to perform these in concert with reactive units such as carbon monoxide. Carbon monoxide is a broadly available C1 building block, and its coupling with organic fragments leads to the formation of valuable carbonyl-containing products.⁴ Moreover, unlike inert aryl-aryl bonds, carbonyl derivatives such as ketones number among the most synthetically versatile functionalities in organic chemistry. The palladium-catalyzed carbonylative functionalization of aryl or even alkyl C-H bonds is well established, and dates back to early work by Fujiwara and others.⁵ However, these reactions commonly lead to carboxylic acids or their derivatives (e.g. esters, amides), and are rarely applicable to the assembly of ketones.⁶ This limitation has been attributed in part to the favored reaction of Pd-acyl intermediates with the carboxylate ligands that are often required for the C-H palladation step, therefore leading instead to the formation of anhydrides.⁷ As shown by Lei, this can be avoided by performing oxidative C-H functionalization in an intramolecular fashion (Fig. 1b).⁸ Nevertheless, the ability to assemble aryl ketones via the simple coupling of its two most fundamental building blocks, arenes and carbon monoxide, has to date presented an unmet challenge.^{9,10}


a. Pd-Catalyzed Oxidative (Hetero)Arene Coupling



b. Intramolecular Carbonylative Arene Coupling



c. Pd-Catalyzed Arene C-H Functionalization via Aryl Triflates



d. This work




Figure 1 Approaches to Intra- and Intermolecular Oxidative C-H Functionalization and Application to Ketone Synthesis.

We have recently described an alternative approach to carbonylative C-H functionalization, wherein palladium catalysts can mediate the *in situ* formation of reactive aryl electrophiles such as acyl iodides and even acyl triflates.¹¹ The latter have the ability to react with unactivated arenes and allowed the development of a carbonylative method to functionalize arenes to form ketones (Fig. 1c). In considering

Department of Chemistry, McGill University, 801 Sherbrooke Street West, Montreal, QC H3A 0B8, Canada. E-mail: bruce.arndtsen@mcgill.ca


*Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x



this chemistry, we questioned if this approach could be taken even further and offer a pathway to perform carbonylative arene/arene coupling with carbon monoxide. In principle, this might be accomplished by devising a method to replace aryl iodides with an oxidative carbonylative platform to convert arenes themselves to high energy acyl triflate electrophiles (Fig. 1d). We describe herein our studies towards such a system. This has led to the discovery of the first catalytic route to assemble ketones directly from two arenes and carbon monoxide. Mechanistic studies suggest this reaction proceeds via a series of steps that ultimately convert arenes into high energy aroyl triflate electrophiles. Exploitation of the latter has opened a novel route to generate heterocoupled ketones from two separate arenes.

Results and Discussion

Our research on this topic grew out of a study on the mechanism of the Pd(0)-catalyzed carbonylative formation of aroyl triflate electrophiles from aryl iodides.^{11d} This reaction can be performed using PdCl₂ and related Pd(II) catalyst precursors (Figure 1c), but it was unclear how these Pd(II) sources are reduced to Pd(0) to mediate the reaction. Carbon monoxide has been established to reduce Pd(II) to Pd(0).^{12,13} However, this often occurs in the presence of nucleophiles (e.g. water) that can trap the oxidized carbon monoxide fragment, and the reaction conditions did not include an obvious nucleophile, or even a dative ligand (e.g. phosphines) that might be oxidized by Pd(II). To probe this, the stoichiometric reaction of PdCl₂ with carbon monoxide and AgOTf was examined in benzene. To our surprise, this leads to the formation of an aromatic product, benzophenone, in up to 71% yield (Scheme 1).



Scheme 1 Generation of Benzophenone via Stoichiometric, PdCl₂-mediated Arene/CO Coupling.

The ability of metal salts such as PdCl₂ to mediate the oxidative carbonylation of arenes to ketones is to our knowledge unknown.¹³ While there are several pathways by which this stoichiometric reaction may occur (*vide infra*), the formal by-product of this reaction is Pd(0), suggesting that a suitable oxidant may transform this stoichiometric chemistry into a catalytic route to form aryl ketones. As shown in Table 1, the addition of various metal salt oxidants that are commonly employed in related oxidative coupling chemistry to the reaction of benzene and CO with 10 mol% PdCl₂ catalyst leads to minimal reaction (entries 1-3), and also inhibits the stoichiometric formation of benzophenone. However, we were pleased to find that simple iodine is an effective oxidant and allows the catalytic formation of benzophenone in good yield (79%, entry 4). Further experimentation showed that the

reaction does not require the use of arene as solvent and proceeds in high yield with a near stoichiometric amount of arene in 1,2-dichloroethane (92%, entry 5) and at temperatures as low as 60 °C with 1 atm CO (entries 6-10). The reaction here was performed with *t*-butylbenzene to aid *in situ* characterization, but is equally efficient with benzene (entry 11). Simple [Pd(allyl)Cl]₂ is the most effective catalytic precursor under these conditions,¹⁴ and the reaction can be performed with as little as 0.1 mol% [Pd(allyl)Cl]₂ at elevated temperatures (entry 12).

Table 1 Development of Conditions for Carbonylative Homocoupling of Arenes.

Entry	Pd Cat.	Ligand	Temp.	Solvent	[Ox]	Yield (%)
1 ^[a]	PdCl ₂	-	100 °C	C ₆ H ₆	Ce(OTf) ₄	0
2 ^[a]	PdCl ₂	-	100 °C	C ₆ H ₆	Na ₂ S ₂ O ₈	2
3 ^[a]	PdCl ₂	-	100 °C	C ₆ H ₆	CuCl ₂	4
4 ^[a]	PdCl ₂	-	100 °C	C ₆ H ₆	I ₂	79
5 ^[a]	PdCl ₂	-	100 °C	1,2-DCE	I ₂	92
6	PdCl ₂	-	60 °C	1,2-DCE	I ₂	23
7	Pd(OAc) ₂	-	60 °C	1,2-DCE	I ₂	60
8	[Pd(allyl)Cl] ₂	PPh ₃ ^[b]	60 °C	1,2-DCE	I ₂	78
9	[Pd(allyl)Cl] ₂	P ^t Bu ₃ ^[b]	60 °C	1,2-DCE	I ₂	57
10	[Pd(allyl)Cl] ₂	-	60 °C	1,2-DCE	I ₂	87
11	[Pd(allyl)Cl] ₂	-	60 °C	1,2-DCE	I ₂	93 (94) ^[c]
12	[Pd(allyl)Cl] ₂ ^[d]	-	100 °C	1,2-DCE	I ₂	72


Conditions: Arene (0.53 mmol), silver triflate (176 mg, 0.68 mmol), Ox (0.25 mmol), 5 mol% Pd, 1 atm CO, 22 h, R = ^tBu. [a] 10 mol% Pd, 4 atm CO, R = H (entries 1-4), ^tBu (entry 5). [b] 10 mol%. [c] 0.75 mmol arene, R = ^tBu (R = H in brackets). [d] 0.1 mol% [Pd(allyl)Cl]₂.

With a palladium-catalyzed method to assemble ketones from arenes and carbon monoxide in hand, we next turned to probing its generality. As shown in Table 2, various electron neutral and electron rich arenes can be employed in this reaction and form ketones in high yields. These include alkyl-substituted arenes (**1b**, **1c**, **1f**) as well as more electron rich anisole derivatives (**1i**, **1j**). In each of the mono-substituted arenes, we observed 4,4'-substituted ketones as the major isolated product, consistent with an electrophilic mechanism for functionalization. Electron deficient substrates, such as mono- and di-halogenated arenes, react under more pressing conditions to form ketones in neat arene (**1d**, **1e**, **1k**). More sterically hindered 1,3- and 1,4-di-substituted arenes can also be converted to ketones (**1f-h**, **1k**, **1l**). In these cases, functionalization proceeds again at the most electron rich site on the arene. This oxidative carbonylation can also be applied to the synthesis of ketones from heteroarenes, with examples including thiophenes (**1n**, **1o**), benzothiophene (**1m**), and *N*-tosyl pyrrole (**1p**). The addition of 2,6-^tbutylpyridine base is



required for these latter examples to inhibit substrate decomposition. Overall, this provides what is to our knowledge the first general catalytic method to assemble aryl ketones in one step from their two most fundamental building blocks, arenes and carbon monoxide, and without the required use of the synthetic, expensive, and/or unstable (hetero)aryl iodides (and organometallic reagents) commonly employed in carbonylative ketone synthesis.

Table 2 Palladium-Catalyzed Oxidative Arene/CO/Arene Coupling to Form Symmetrical Ketones.




Conditions: Arene (1.50 mmol), AgOTf (350 mg, 1.36 mmol), I₂ (126 mg, 0.50 mmol), [Pd(allyl)Cl]₂ (5 mg, 0.013 mmol), 1,2-dichloroethane (2 mL), 4 atm CO, 22 h, 60 °C (see SI for details). Isolated Yield (ratio of 4':2' isomers). [a] At 150 °C in arene solvent. [b] At 100 °C. [c] With 2,6-di-tert-butylpyridine (258 mL, 1.15 mmol) (ratio of 3':2' isomers).


We next turned to examine how this transformation proceeds. As mechanistic possibilities, the stoichiometric reaction of PdCl₂ (Scheme 1) suggests the potential *in situ* formation of phosgene equivalents upon reduction of Pd(II) by CO (Fig. 2a, path A),¹³ or the build-up of electrophilic Pd(II) triflate intermediates that may directly palladate arenes (path B).¹⁵ These could each lead to aryl triflate intermediates for subsequent Friedel-Crafts acylations,^{11d} and are consistent with the regioselectivity in the ketone products formed. The re-oxidation of Pd(0) by I₂ would then make these viable cycles for catalysis. However, iodine is also established to react with arenes in the presence of silver triflate to form aryl iodides.¹⁶ It is therefore possible that the catalytic arene carbonylation

occurs via an alternative Pd(0) cycle to form aryl triflate products (path C). Monitoring the reaction by *in situ* ¹H and ¹³C NMR analysis show that the latter mechanism is likely the case during catalysis, and we observe the rapid, high yield build-up of aryl iodide within 1 h under the catalytic conditions (Fig. 2b, 94% yield). Continued heating converts this aryl iodide to ketone **1a**. Similarly, control experiments show that the AgOTf mediated iodination of the arene is rapid (30 min at r.t.), relative to catalytic ketone formation (23% after 22 h at r.t.; Fig. 2c). Thus, in the presence of I₂ as oxidant, these data suggest that the direct iodination of the arene is the most rapid pathway (path C). Nevertheless, the ability of Pd(II) salts themselves to mediate the direct carbonylative C-H functionalization of arenes also appears to be relevant to catalysis, as it allows the activation of the Pd(II) pre-catalyst (e.g. Table 1). The compatibility between Pd(0) catalysis and I₂ is unusual, since Pd(0) can also undergo rapid oxidation. This presumably reflects the rapid rate of arene iodination relative to the slower carbonylation and build-up of Pd(0), as well as the ability of any *in situ* generated Pd(II) to re-enter the cycle (e.g. via paths A or B). The lack of potentially oxidizable donor ligands in the reaction further enhances this compatibility. Together, this suggests a new avenue to use Pd(0) catalysis in oxidative coupling chemistry with I₂.

a. Possible Mechanistic Pathways



b. In Situ Monitoring of Catalytic Reaction



c. Iodination and Carbonylation Reactions at Room Temperature




Figure 2. Proposed Catalytic Pathway for Ketone Formation.




From a synthetic perspective, the generation of halogenated arene intermediates can have useful implications. Since these intermediates are formed more rapidly than carbonylation, the rate difference can be exploited to perform cross arene/arene coupling to build up more structurally diverse ketones. The latter can present a challenge with many oxidative arene/arene coupling reactions, and typically require significant differences in the electronic, steric, or coordinating ability of the two substrates.² After examination of various reaction conditions, we were pleased to find that the ambient temperature reaction of arene, [Pd(allyl)Cl]₂ catalyst, I₂, and AgOTf, followed by the subsequent addition of a second, more electron rich arene and carbon monoxide leads to the clean


overall heterocoupling of arenes to ketones (Table 3). Both electron neutral and electron rich arenes can be easily cross-coupled to form unsymmetrical ketones in high yield, including sterically encumbered tetra-substituted arenes and an activated benzophenone (**2f**, **2h**, **2i**). Employing more pressing conditions allows for the use of deactivated arenes such as bromo- or dichlorobenzene (**2e**, **2g**), and does so without secondary activation of these carbon-halogen bonds. Thiophene, *N*-substituted pyrrole, and benzothiophene can also be incorporated into this reaction to create mixed (hetero)aryl ketones (**2k**, **2m**, **2n**).

Finally, we have examined the use of this system to assemble more structurally elaborate products. As an example, Crystal Violet (**3**, Scheme 2) is a common dye exploited in textiles. Crystal Violet is typically generated starting from reactive, synthetic acylating agents (e.g. phosgene), in a sequence that proceeds through the initial build-up of ketone **4**.¹⁷ Alternatively, the palladium-catalyzed oxidative carbonylation of *N,N*-dimethylaniline can offer a method to generate **3** from the parent arene and carbon monoxide (Scheme 2). While **3** is generated in moderate yield (41% together with its de-methylated isomer), it represents the product of three concurrent arene C-H bond functionalization steps. This reaction presumably proceeds via the initial generation of **4** that undergoes a spontaneous Friedel-Crafts reaction to liberate H₂O, which can also be trapped by the *in situ* generated aryl electrophiles. As far as we are aware, this is the first synthesis of such an advanced structure directly from arenes and CO.

Table 3 Palladium-Catalyzed Oxidative Arene/CO/Arene Coupling to Form Unsymmetrical Ketones.



Conditions: Arene (1.00 mmol), AgOTf (339 mg, 1.32 mmol), I₂ (126 mg, 0.49 mmol), [Pd(allyl)Cl]₂ (5 mg, 0.013 mmol), 1,2-dichloroethane (2 mL), 4-6 h, r.t., then second arene (1.50 mmol), 4 atm CO, 22 h, 60 °C (see SI for details). Isolated yields (3':2' ratio). [a] Both steps at 100 °C. [b] Step 1 at 100 °C, 22 h; step 2 at 150 °C w/ 10 eq. bromobenzene. [c] With 2,6-di-tert-butylpyridine (1.15 mmol).



Scheme 2 Synthesis of Crystal Violet from Dimethylaniline and Carbon Monoxide

Conclusions

In conclusion, we have described herein the first catalytic route to prepare ketones from arenes and CO. The reaction generates either symmetrical or unsymmetrical aryl ketones in high yields does so from combinations of stable, commercially available, and easily modulated reagents, and without the need for synthetic and often expensive aryl iodide reagents. Mechanistic analysis suggests this transformation proceeds via the palladium-catalyzed, *in situ* generation of potent acylating electrophiles (acyl triflates) from arenes. Considering the broad utility of acylating electrophiles, we anticipate that the ability to access these intermediates from arenes could prove



relevant to the design of various new approaches to carbonylative C-H bond functionalization chemistry.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank NSERC, the Canadian Foundation for Innovation (CFI), the McGill Sustainability Systems Initiative (MSSI) through the Ideas Fund, and the FQRNT supported Centre for Green Chemistry and Catalysis for funding this research.

Notes and references

- For reviews: a) J. F. Hartwig, *J. Am. Chem. Soc.* 2016, **138**, 2-24; b) T. W. Lyons, M. S. Sanford, *Chem. Rev.* 2010, **110**, 1147-1169; c) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem. Int. Ed.* 2009, **48**, 5094-5115; d) T. Cernak, K. D. Dykstra, S. Tyagarajan, P. Vachal, S. W. Kraska, *Chem. Soc. Rev.* 2016, **45**, 546-576; e) J. F. Hartwig, M. A. Larsen, *ACS Cent. Sci.* 2016, **2**, 281-292; f) P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz, L. Ackermann, *Chem. Rev.* 2019, **119**, 2192-2452; g) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* 2009, **42**, 1074-1086; h) P. Wedi, M. van Gemmeren, *Angew. Chem. Int. Ed.* 2018, **57**, 13016-13027.
- a) C. Liu, J. Yuan, M. Gao, S. Tang, W. Li, R. Shi, A. Lei, *Chem. Rev.* 2015, **115**, 12138-12204; b) Y. Yang, J. Lan, J. You, *Chem. Rev.* 2017, **117**, 8787-8863; c) C. S. Yeung, V. M. Dong, *Chem. Rev.* 2011, **111**, 1215-1292; d) Y.-F. Zhang, Z.-J. Shi, *Acc. Chem. Res.* 2019, **52**, 161-169; e) J. A. Ashenhurst, *Chem. Soc. Rev.* 2010, **39**, 540-548; f) C.-J. Li, *Acc. Chem. Res.* 2009, **42**, 335-344.
- For representative examples: a) R. Li, L. Jiang, W. Lu, *Organometallics* 2006, **25**, 5973-5975; b) X. Wang, D. Leow, J.-Q. Yu, *J. Am. Chem. Soc.* 2011, **133**, 13864-13867; c) D. R. Stuart, K. Fagnou, *Science* 2007, **316**, 1172-1175; d) T. W. Lyons, K. L. Hull, M. S. Sanford, *J. Am. Chem. Soc.* 2011, **133**, 4455-4464; e) K. Kawasumi, K. Mochida, T. Kajino, Y. Segawa, K. Itami, *Org. Lett.* 2012, **14**, 418-421; f) X. Zhao, C. S. Yeung, V. M. Dong, *J. Am. Chem. Soc.* 2010, **132**, 5837-5844; g) G. Brasche, J. García-Fortanet, S. L. Buchwald, *Org. Lett.* 2008, **10**, 2207-2210; h) K. Matsumoto, S. Takeda, T. Hirokane, M. Yoshida, *Org. Lett.* 2019, **21**, 7279-7283; i) D. Wang, S. S. Stahl, *J. Am. Chem. Soc.* 2017, **139**, 5704-5707; j) B.-J. Li, S.-L. Tian, Z. Fang, Z.-J. Shi, *Angew. Chem. Int. Ed.* 2008, **47**, 1115-1118; k) Y. Wei, W. Su, *J. Am. Chem. Soc.* 2010, **132**, 16377-16379; l) C.-Y. He, S. Fan, X. Zhang, *J. Am. Chem. Soc.* 2010, **132**, 12850-12852.
- a) M. Beller, X.-F. Wu, *Transition Metal Catalyzed Carbonylations-Carbonylative Activation of C-X Bonds*, Springer, Berlin, 2013; b) Y. Li, Y. Hu, X.-F. Wu, *Chem. Soc. Rev.* 2018, **47**, 172-194; c) A. Brennfürher, H. Neumann, M. Beller, *Angew. Chem. Int. Ed.* 2009, **48**, 4114-4133; d) I. Omae, *Coord. Chem. Rev.* 2011, **255**, 139-160.
- For reviews see: a) C. Jia, T. Kitamura, Y. Fujiwara, *Acc. Chem. Res.* 2001, **34**, 633-639; For recent examples: b) K. Orito, A. Horibata, T. Nakamura, H. Ushito, H. Nagasaki, M. Yuguchi, S. Yamashita, M. Tokuda, *J. Am. Chem. Soc.* 2004, **126**, 14342-14343; c) R. Giri, J.-Q. Yu, *J. Am. Chem. Soc.* 2008, **130**, 14082-14083; d) R. Giri, J. K. Lam, J.-Q. Yu, *J. Am. Chem. Soc.* 2010, **132**, 686-693; e) Z.-H. Guan, M. Chen, Z.-H. Ren, *J. Am. Chem. Soc.* 2012, **134**, 17490-17493; f) S. Luo, F.-X. Luo, X.-S. Zhang, Z.-J. Shi, *Angew. Chem. Int. Ed.* 2013, **52**, 10598-10601; g) D. Willcox, B. G. N. Chappell, K. F. Hogg, J. Calleja, A. P. Smalley, M. J. Guant, *Science* 2016, **354**, 851-857.
- Q. Liu, H. Zhang, A. Lei, *Angew. Chem. Int. Ed.* 2011, **50**, 10788-10799.
- a) M. A. Campo, R. C. Larock, *J. Org. Chem.* 2002, **67**, 5616-5620; b) X.-F. Wu, P. Anbarasan, H. Neumann, M. Beller, *Angew. Chem. Int. Ed.* 2010, **49**, 7316-7319; c) Z. Lian, S. D. Friis, T. Skrydstrup, *Chem. Commun.* 2015, **51**, 1870-1873; d) J. Liu, Z. Wei, H. Jiao, R. Jackstell, M. Beller, *ACS Cent. Sci.* 2018, **4**, 30-38; e) T. A. Cernak, T. H. Lambert, *J. Am. Chem. Soc.* 2009, **131**, 3124-3125; f) F. Zeng, H. Alper, *Org. Lett.* 2013, **15**, 2034-2037.
- a) H. Zhang, R. Shi, P. Gan, C. Liu, A. Ding, Q. Wang, A. Lei, *Angew. Chem. Int. Ed.* 2012, **51**, 5204-5207; b) J. Wen, S. Tang, F. Zhang, R. Shi, A. Lei, *Org. Lett.* 2017, **19**, 94-97.
- a) X.-F. Wu, H. Neumann, M. Beller, *Chem. Soc. Rev.* 2011, **40**, 4986-5009; b) M.-N. Zhao, L. Ran, M. Chen, Z.-H. Ren, Y.-Y. Wang, Z.-H. Guan, *ACS Catal.* 2015, **5**, 1210-1213.
- For an alternative approach from aldehydes: a) S. Tetsuya, I. Tomoaki, M. Masahiro, N. Masakatsu, *Chem. Lett.* 1996, **25**, 823-824; b) S. Ko, B. Kang, S. Chang, *Angew. Chem. Int. Ed.* 2005, **44**, 455-457; c) J. Ruan, O. Saidi, J. A. Iggo, J. Xiao, *J. Am. Chem. Soc.* 2008, **130**, 10510-10511; d) X. Jia, S. Zhang, W. Wang, F. Luo, J. Cheng, *Org. Lett.* 2009, **11**, 3120-3123. e) P. Álvarez-Bercedo, A. Flores-Gaspar, A. Correa, R. Martin, *J. Am. Chem. Soc.* 2010, **132**, 466-467; f) B. Suchand, G. Satyanarayana, *J. Org. Chem.* 2016, **81**, 6409-6423; g) T. Wakaki, T. Togo, D. Yoshidome, Y. Kuninobu, M. Kanai, *ACS Catal.* 2018, **8**, 3123-3128.
- a) J. S. Quesnel, B. A. Arndtsen, *J. Am. Chem. Soc.* 2013, **135**, 16841-16844; b) J. S. Quesnel, L. V. Kayser, A. Fabrikant, B. A. Arndtsen, *Chem. Eur. J.* 2015, **21**, 9550-9555; c) J. Tjuttrins, B. A. Arndtsen, *J. Am. Chem. Soc.* 2015, **137**, 12050-12054; d) R. G. Kinney, J. Tjuttrins, G. M. Torres, N. J. Liu, O. Kulkarni, B. A. Arndtsen, *Nat. Chem.* 2018, **10**, 193-199; e) P.-L. Lagueux-Tremblay, A. Fabrikant, B. A. Arndtsen, *ACS Catal.* 2018, **8**, 5350-5354.
- a) J. Tsuji, *Acc. Chem. Res.* 1969, **2**, 144-152; b) J. Tsuji, *Palladium Reagents and Catalysts – New Perspectives for the 21st Century*, John Wiley & Sons Ltd., West Sussex, 2004; c) C. F. J. Barnard, *Organometallics* 2008, **27**, 5402-5422; d) X.-F. Wu, H. Neumann, A. Spannenberg, T. Schulz, H. Jiao, M. Beller, *J. Am. Chem. Soc.* 2010, **132**, 14596-14602.
- PdCl₂ has been suggested to react with CO to generate phosgene derivatives. a) T. A. Stromnova, M. N. Vargaftik, I. I. Moiseev, *J. Organomet. Chem.* 1983, **252**, 113-120; b) D. B. Dell'Amico, F. Calderazzo, N. Zandona, *Inorg. Chem.* 1984, **23**, 137-140; c) S. Baig, B. Richard, P. Serp, C. Mijoule, K. Hussein, N. Guihéry, J.-C. Barthelat, P. Kalck, *Inorg. Chem.* 2006, **45**, 1935-1944; d) F. Ragaini, H. Larici, M. Rimoldi, A. Caselli, F. Ferretti, P. Macchi, N. Casati, *Organometallics* 2011, **30**, 2385-2393.
- The reaction proceeds well with a near stoichiometric amount of arene (2.1 : 1 arene to I₂ ratio, entry 10), but we observed slightly elevated yields with 1.5 equiv. of arene (3 : 1 ratio of arene to I₂).
- For mechanistic studies with related transition metal acetates: a) D. Lapointe, K. Fagnou, *Chem. Lett.* 2010, **39**, 1118-1126; b) S. I. Gorelsky, *Coord. Chem. Rev.* 2013, **257**, 153-164; c) L. Wang, B. P. Carrow, *ACS Catal.* 2019, **9**, 6821-6836; d) D. Whitaker, J. Burés, I. Larrosa, *J. Am. Chem. Soc.* 2016, **138**, 8384-8387; e) S. Y. Lee, J. F. Hartwig, *J. Am. Chem. Soc.* 2016, **138**, 15278-15284; f) M. D. Lotz, N. M. Camasso, A. J. Canty, M. S. Sanford, *Organometallics* 2017, **36**, 165-171; g) C. Colletto, A. Panigrahi, J. Fernández-Casado, I. Larrosa, *J. Am. Chem. Soc.* 2018, **140**, 9638-9643.



ARTICLE


Journal Name

- 16 a) G. K. Mulholland, Q.-H. Zheng, *Synth. Commun.* 2001, **31**, 3059-3068; b) S. Wan, S. R. Wang, W. Lu, *J. Org. Chem.* 2006, **71**, 4349-4352.
- 17 T. Gessner, U. Mayer, *Triarylmethane and Diarylmethane Dyes: Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, Weinheim, 2000.

View Article Online
DOI: 10.1039/D0SC00085J



Table of Contents Entry



A palladium catalyzed route to prepare aryl ketones from their two fundamental building blocks, two arenes and carbon monoxide, is described.

