

Bioactive Constituents from Chinese Natural Medicines. XV.¹⁾ Inhibitory Effect on Aldose Reductase and Structures of Saussureosides A and B from *Saussurea medusa*

Haihui XIE,^a Tao WANG,^a Hisashi MATSUDA,^a Toshio MORIKAWA,^a Masayuki YOSHIKAWA,^{*,a} and Tadato TANI^b

^a Kyoto Pharmaceutical University; Misasagi, Yamashina-ku, Kyoto 607-8412, Japan; and ^b Department of Kampo-pharmaceutics, Institute of Natural Medicine, Toyama Medical and Pharmaceutical University; 2630 Sugitani, Toyama 930-0194, Japan. Received June 15, 2005; accepted July 15, 2005

The 80% aqueous acetone extract from the whole plant of *Saussurea medusa* MAXIM. was found to inhibit rat lens aldose reductase (IC₅₀=1.4 µg/ml). From this extract, flavonoids, lignans, and quinic acid derivatives were isolated together with two new ionone glycosides, saussureosides A and B. Their absolute stereostructures were elucidated on the basis of chemical and physicochemical evidence including the application of modified Mosher's method. In addition, some isolates were found to show an inhibitory effect on aldose reductase.

Key words *Saussurea medusa*; aldose reductase inhibitor; saussureoside; flavonoid; ionone glycoside

The Compositae plant, *Saussurea medusa* MAXIM., is a rare perennial medicinal herb growing in the northwestern parts of China (e.g., Tibet, Xinjiang, Qinghai, Gansu, Yunnan, and Sichuan provinces). The whole plant or aerial parts of *S. medusa*, popularly known as “Xuelianhua (雪蓮花)”, has been commonly used in both traditional Chinese medicine and Tibetan folk medicine as a remedy for rheumatic arthritis, menoxenia, gynopathy, traumatic bleeding, and headache.^{2–4)} Several lignan, neolignan, flavonoid, terpenoid, and chlorophyll constituents have previously been isolated from *S. medusa*^{5–10)} and their pharmacologic activities such as anti-tumor-promoting,⁵⁾ immunosuppressive⁶⁾ and cell-protecting activities⁷⁾ were also reported. In the course of our studies on bioactive constituents from Chinese natural medicines,^{11–17)} we found that the 80% aqueous acetone extract from the whole plant of *S. medusa* showed a potent inhibitory effect on rat lens aldose reductase (IC₅₀=1.4 µg/ml). In this paper, we report the isolation and structure elucidation of two new megastigmane glycosides, saussureosides A (**1**) and B (**2**), together with 13 flavonoids (**3**–**15**), four lignans (**16**–**19**) and three neolignans (**20**–**22**), three quinic acids (**23**–**25**), and 24 other known constituents.

The 80% aqueous acetone extract from the whole plant of *S. medusa* was partitioned into an EtOAc–H₂O (1:1, v/v) mixture to furnish the EtOAc-soluble fraction and aqueous layer. The aqueous layer was extracted with *n*-butanol (*n*-BuOH) to give *n*-BuOH and H₂O-soluble fractions. The *n*-BuOH-soluble fraction, which was the most active fraction (IC₅₀=0.8 µg/ml), was subjected to normal-phase and reversed-phase silica gel column chromatography and repeated HPLC to give saussureosides A (**1**, 0.0007%) and B (**2**, 0.0038%), apigenin 7-*O*-β-D-glucopyranoside¹⁸⁾ (**4**, 0.0016%), apigenin 4'-*O*-β-D-glucopyranoside¹⁹⁾ (**5**, 0.0089%), apigenin 7-*O*-rutinoside²⁰⁾ (**6**, 0.0054%), luteolin 7-*O*-β-D-glucopyranoside²⁰⁾ (**8**, 0.034%), luteolin 4'-*O*-β-D-glucopyranoside¹⁹⁾ (**9**, 0.042%), luteolin 7-*O*-rutinoside²⁰⁾ (**10**, 0.023%), chrysoeriol 7-*O*-β-D-glucopyranoside²¹⁾ (**11**, 0.0034%), chrysoeriol 7-*O*-rutinoside²²⁾ (**12**, 0.0030%), quercetin 3-*O*-β-D-glucopyranoside²³⁾ (**14**, 0.079%), isorhamnetin 3-*O*-rutinoside²⁴⁾ (**15**, 0.0035%), (+)-pinoreosin

O-β-D-glucopyranoside²⁵⁾ (**17**, 0.0026%), (+)-syringaresinol *O*-β-D-glucopyranoside²⁶⁾ (**19**, 0.0011%), arctiin²⁷⁾ (**22**, 0.062%), 3-*O*-caffeoylquinic acid methyl ester^{28,29)} (**23**, 0.0011%), 4-*O*-caffeoylquinic acid methyl ester²⁸⁾ (**24**, 0.0012%), 5-*O*-caffeoylquinic acid methyl ester³⁰⁾ (**25**, 0.019%), syringin³¹⁾ (0.0063%), junipediol A³²⁾ (0.0011%), phenethyl *O*-β-D-glucopyranoside³³⁾ (0.0007%), benzyl *O*-β-D-glucopyranoside³⁴⁾ (0.0028%), *p*-anisyl *O*-β-D-glucopyranoside³⁵⁾ (0.0006%), (*Z*)-3-hexenyl *O*-β-D-glucopyranoside³⁶⁾ (0.0010%), L-phenylalanine³⁷⁾ (0.0018%), thymidine³⁷⁾ (0.0026%), and sitosterol 3β-*O*-β-D-glucopyranoside^{29,38)} (0.0046%). The EtOAc-soluble fraction was also subjected to normal-phase and reversed-phase silica gel column chromatography and repeated HPLC to give apigenin²³⁾ (**3**, 0.0021%), luteolin²³⁾ (**7**, 0.0069%), quercetin²³⁾ (**13**, 0.0060%), (+)-pinoreosin³⁹⁾ (**16**, 0.0004%), (+)-syringaresinol⁴⁰⁾ (**18**, 0.0004%), (–)-secoisolaricresinol⁴¹⁾ (**20**, 0.0007%), (–)-arctigenin²⁷⁾ (**21**, 0.038%), **22** (0.054%), 3-methoxy-4-hydroxyphenylethanol⁴²⁾ (0.0021%), (–)-2-oxoisodauc-5-en-12-al⁴³⁾ (0.0002%), (+)-glutinin⁴⁴⁾ (0.0006%), α-amyrin⁴⁵⁾ (0.0086%), lupeol⁴⁵⁾ (0.0074%), clionasterol⁴⁶⁾ (0.0026%), azelaic acid³⁷⁾ (0.0024%), linolenic acid³⁷⁾ (0.027%), linoleic acid³⁷⁾ (0.065%), oleic acid³⁷⁾ (0.0093%), hexadecanoic acid³⁷⁾ (0.075%), octadecanoic acid³⁷⁾ (0.0092%), eicosanoic acid³⁷⁾ (0.0028%), docosanoic acid³⁷⁾ (0.0025%), tricosanoic acid³⁷⁾ (0.0007%), and eicosanol³⁷⁾ (0.0011%).

Structure of Saussureosides A (1**) and B (**2**)** Saussureoside A (**1**) was isolated as a white powder with negative optical rotation ([α]_D²⁴ –45.0°). In the UV spectrum of **1**, an absorption maximum was observed at 229 (log ε 4.07) nm, while the IR spectrum of **1** showed absorption bands at 1684, 1671, and 1260 cm^{–1} assignable to α,β-unsaturated carbonyl, olefin, and epoxide functions and strong absorption bands at 3410 and 1078 cm^{–1} suggestive of a glycoside moiety. In the positive-ion FAB-MS of **1**, a quasimolecular ion peak was observed at *m/z* 425 (M+Na)⁺. The molecular formula C₁₉H₃₀O₉ of **1** was determined from the quasimolecular ion peak observed in the FAB-MS and by high-resolution FAB-MS measurement. The ¹H-NMR (pyridine-*d*₅, Table 2) and

* To whom correspondence should be addressed. e-mail: shoyaku@mb.kyoto-phu.ac.jp




Chart 1

Table 1. Inhibitory Effects of 80% Aqueous Acetone Extract and EtOAc-, *n*-BuOH-, H₂O-Soluble Fractions from *Saussurea medusa* on Rat Lens Aldose Reductase

	IC ₅₀ (μ g/ml)
80% Aqueous acetone extract	1.4
EtOAc-soluble fraction	1.4
<i>n</i> -BuOH-soluble fraction	0.8
H ₂ O-soluble fraction	2.2

¹³C-NMR (Table 3) spectra⁴⁷⁾ of **1** showed signals assignable to four methyls [δ 0.98, 1.02, 1.67, 2.24 (all s, 12, 13, 11, 10-H₃)], a methylene [δ 1.49 (dd, $J=3.4, 12.5$ Hz), 1.94 (dd, $J=12.2, 12.5$ Hz), 2-H₂], two methines bearing an oxygen function [δ 4.25 (ddd, $J=3.4, 3.4, 12.2$ Hz, 3-H), 4.40 (d, $J=3.4$ Hz, 4-H)], and a *trans*-olefin pair [δ 6.47, 7.03 (both d, $J=15.9$ Hz, 8, 7-H)] together with a β -glucopyranosyl part [δ 5.00 (d, $J=6.4$ Hz, 1'-H)]. Acid hydrolysis of **1** with 1 M hydrochloric acid (HCl) liberated D-glucose, which was identified by HPLC analysis using an optical rotation detector.^{1,13–15,17,48)} The aglycon **1a** was obtained by enzymatic hydrolysis of **1** with β -glucosidase. As shown in Fig. 1, the

¹H-¹H correlation spectroscopy (¹H-¹H COSY) experiment on **1** indicated the presence of partial structures written in bold lines, and in the heteronuclear multiple-bond correlations (HMBC) experiment, long-range correlations were observed between the following protons and carbons (2-H₂, 12-H₃, 13-H₃ and 1-C; 4-H, 11-H₃ and 5-C; 4-H, 7-H, 12-H₃, 13-H₃ and 6-C; 7-H, 8-H, 10-H₃ and 9-C; 1'-H and 4-C). Furthermore, comparison of the ¹³C-NMR data of **1** with those of **1a** revealed a glycosylation shift around the 4-position in **1**. Thus, the connectivity of the β -D-glucopyranosyl moiety in **1** was clarified to be the 4-position of **1a**. The relative stereostructure of the aglycon part (**1a**) in **1** was characterized by nuclear Overhauser enhancement spectroscopy (NOESY) experiment, which showed NOE correlations between the following proton pairs (2 α -H and 3-H; 2 β -H and 13-H₃; 3-H and 4-H; 11-H₃ and 4-H, 7-H₂) as shown in Fig. 1. Finally, the absolute configurations of **1** and **1a** were characterized by application of the modified Mosher's method.⁴⁹⁾ Namely, treatment of **1a** with (*R*)- or (*S*)-2-methoxy-2-trifluoromethylphenylacetic acid [(*R*)- or (*S*)-MTPA] in the presence of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC·HCl) and 4-dimethylaminopyridine (4-DMAP) selectively yielded the 3-mono-MTPA esters (**1b**, **c**), respectively,



Fig. 1. H-H COSY, HMBC, and NOE Correlations of **1** and **2**

Fig. 2

whose ^1H -NMR data showed an acylation shift at the 3-position in **1a**. As shown in Fig. 2, the signals due to protons attached to the 4 and 11-positions in the 3-mono-(S)-MTPA ester (**1c**) were observed at lower fields compared with those of the 3-mono-(R)-MTPA ester (**1b**) [$\Delta\delta$: positive], while the signals due to protons on the 2 and 13-positions in **1c** were observed at higher fields compared with those of **1b** [$\Delta\delta$: negative]. Consequently, the absolute configuration at the 3-position of **1a** was determined to be S configuration and the absolute stereostructures of **1** and **1a** were elucidated as shown.

Saussureoside B (**2**) was obtained as a white powder with negative optical rotation ($[\alpha]_D^{26} -39.5^\circ$). The molecular formula of **2** was determined from the positive-ion FAB-MS and by high-resolution FAB-MS analysis to be $\text{C}_{19}\text{H}_{30}\text{O}_8$. In the UV spectrum of **2**, absorption maxima were observed at 227 (log ϵ 3.89) and 279 (log ϵ 3.78) nm, while the IR spectrum of **2** showed absorption bands due to hydroxyl, conjugated enone, and ether functions (3432, 1671, 1655, 1076 cm^{-1}). The ^1H -NMR (Table 2, pyridine- d_5) and ^{13}C -NMR (Table 3) spectra⁴⁷⁾ of **2** showed the presence of the following func-

Table 2. ^1H -NMR (500 MHz, Pyridine- d_5) Data of Saussureosides A (**1**) and B (**2**)

H-	1	2
2 α	1.49 dd (3.4, 12.5)	1.70 dd (3.0, 12.5)
2 β	1.94 dd (12.2, 12.5)	2.06 dd (12.5, 12.5)
3	4.25 ddd (3.4, 3.4, 12.2)	4.06 ddd (3.0, 3.4, 12.5)
4	4.40 d (3.4)	4.30 d (3.4)
7	7.03 d (15.9)	7.22 d (16.5)
8	6.47 d (15.9)	6.11 d (16.5)
10	2.24 s	2.27 s
11	1.67 s	2.11 s
12	0.98 s	1.05 s
13	1.02 s	1.02 s
1'	5.00 d (6.4)	5.00 d (6.4)
2'	4.11 m	4.09 m
3'	4.26 m	4.24 m
4'	4.25 m	4.23 m
5'	4.06 m	4.05 m
6'	4.36 dd (5.2, 11.4)	4.35 dd (5.8, 11.9)
	4.58 dd (2.1, 11.4)	4.58 dd (2.5, 11.9)

δ in ppm and J (in parentheses) in Hz.

Table 3. ^{13}C -NMR (125 MHz) Data of Saussureosides A (**1**) and B (**2**), **1a**, and **2a**

C-	1 ^{a)}	1a ^{b)}	2 ^{a)}	2a ^{b)}
1	34.7	34.5	37.1	37.0
2	42.0	39.4	43.0	42.4
3	64.9	65.9	66.1	66.8
4	84.7	71.6	84.3	71.9
5	69.0	68.8	131.5	134.3
6	70.5	70.1	139.7	138.7
7	142.7	142.2	141.9	142.5
8	133.5	132.9	133.9	133.7
9	197.0	197.4	197.0	197.7
10	27.6	28.1	27.4	27.4
11	16.9	16.8	19.9	20.2
12	25.0	24.6	27.1	27.6
13	29.0	29.2	29.9	30.0
1'	106.6		106.9	
2'	75.1		75.0	
3'	78.5		78.3	
4'	71.5		71.4	
5'	78.7		78.7	
6'	62.4		62.4	

Measured in a) pyridine- d_5 and b) CDCl_3 .

tions: four methyls [δ 1.02, 1.05, 2.11, 2.27 (all s, 13, 12, 11, 10- H_3)], a methylene [δ 1.70 (dd, $J=3.0$, 12.5 Hz), 2.06 (dd, $J=12.5$, 12.5 Hz), 2- H_2], two methines bearing an oxygen function [δ 4.06 (ddd, $J=3.0$, 3.4, 12.5 Hz, 3-H), 4.30 (d, $J=3.4$ Hz, 4-H)], and a *trans*-olefin pair [δ 6.11, 7.22 (both d, $J=16.5$ Hz, 8, 7-H)] together with a β -glucopyranosyl part [δ 5.00 (d, $J=6.4$ Hz, 1'-H)]. The acid hydrolysis of **2** with 1 M HCl liberated D-glucose, which was identified by HPLC analysis using an optical rotation detector.^{1,13-15,17,48)} The aglycon **2a** was obtained by enzymatic hydrolysis of **2** with β -glucosidase. The structures of **2** and **2a** were confirmed by ^1H - ^1H COSY and HMBC experiments. That is, the ^1H - ^1H COSY experiment on **2** indicated the presence of the partial structures written in bold lines, and in the HMBC experiment, long-range correlations were observed between the following protons and carbons (2-H, 12- H_3 , 13- H_3 and 1-C; 4-H, 11- H_3 and 5-C; 4-H, 7-H, 12- H_3 ,

Table 4. Inhibitory Effects of Constituents from *Saussurea medusa* on Rat Lens Aldose Reductase

	IC ₅₀ (μM)
Saussuroside A (1)	>100 (3%) ^{a)}
Saussuroside B (2)	>100 (13%)
Apigenin (3)	2.2 ⁵²⁾
Apigenin 7- <i>O</i> -β-D-glucopyranoside (4)	4.4
Apigenin 4'- <i>O</i> -β-D-glucopyranoside (5)	3.2
Apigenin 7- <i>O</i> -rutinoside (6)	4.7
Luteolin (7)	0.45 ⁵²⁾
Luteolin 7- <i>O</i> -β-D-glucopyranoside (8)	0.99 ⁵²⁾
Luteolin 4'- <i>O</i> -β-D-glucopyranoside (9)	4.8
Luteolin 7- <i>O</i> -rutinoside (10)	0.92
Chrysoeriol 7- <i>O</i> -β-D-glucopyranoside (11)	26
Chrysoeriol 7- <i>O</i> -rutinoside (12)	14
Quercetin (13)	2.2 ⁵²⁾
Quercetin 3- <i>O</i> -β-D-glucopyranoside (14)	4.5 ⁵²⁾
Isorhamnetin 3- <i>O</i> -rutinoside (15)	19 ⁵⁶⁾
(+)-Pinorelinol (16)	>100 (34%)
(+)-Pinorelinol <i>O</i> -β-D-glucopyranoside (17)	>100 (37%)
(+)-Syringaresinol (18)	>100 (13%)
(+)-Syringaresinol <i>O</i> -β-D-glucopyranoside (19)	>100 (38%)
(-)-Secoisolariciresinol (20)	>100 (26%)
(-)-Arctigenin (21)	>100 (16%)
Arctiin (22)	20
3-Caffeoylquinic acid methyl ester (23)	13
4-Caffeoylquinic acid methyl ester (24)	16
5-Caffeoylquinic acid methyl ester (25)	1.3
Epalrestat	0.072

a) Values in parentheses indicate inhibition (%) at 100 μM.

13-H₃ and 6-C; 7-H, 8-H, 10-H₃ and 9-C; 1'-H and 4-C) (Fig. 1).

The relative stereostructure of **2** was determined by a NOESY experiment, in which correlations were observed between the following proton pairs (2α-H and 3-H; 2β-H and 13-H₃; 3-H and 4-H). The absolute configurations of **2** and **2a** were determined by a modified Mosher's method.⁴⁹⁾ As shown in Fig. 2, the 3-mono-MTPA esters (**2b** and **2c**) were derived from **2a** by a method similar to that for **1a**. The protons on the 4 and 11-positions of the (*S*)-MTPA ester (**2c**) resonated at lower fields than those of the (*R*)-MTPA ester (**2b**) [Δδ: positive], while the protons on the 2, 12, and 13-positions of **2c** were observed at higher fields compared to those of **2b** [Δδ: negative]. Consequently, the absolute configurations at the 3-position in **2** and **2a** were elucidated to be *S*. On the basis of this evidence, the absolute configurations of **2** and **2a** were determined as shown.^{50,51)}

Inhibitory Effect on Rat Lens Aldose Reductase As a key enzyme in the polyol pathway, aldose reductase has been reported to catalyze the reduction of glucose to sorbitol. Sorbitol does not readily diffuse across cell membranes, and the intracellular accumulation of sorbitol has been implicated in the chronic complications of diabetes such as cataract. Previously, we reported that various flavonoid and terpenoid constituents from several natural medicines and medicinal foods inhibited aldose reductase inhibitory activity.^{52–57)} Since the aqueous acetone extract from the whole plants of *S. medusa* was found to show aldose reductase inhibitory activity, the principal isolated constituents were examined (Table 4).

Among these principal constituents, flavones with the 3',4'-dihydroxyl group [e.g., luteolin 7-*O*-rutinoside (**10**, IC₅₀=0.92 μM)] showed strong activity in agreement with the

previous study.⁵²⁾ In contrast, lignan constituents (**16–22**) showed weak activity. As to the activity of caffeoylquinic acid methyl esters, 5-*O*-caffeoylquinic acid methyl ester (**25**, 1.3 μM) showed stronger activity than **23** and **24**, and this is the first report of caffeoylquinic acid methyl esters with inhibitory activity for aldose reductase.

Experimental

The following instruments were used to obtain physical data: specific rotations, Horiba SEPA-300 digital polarimeter (*l*=5 cm); UV spectra, Shimadzu UV-1600 spectrometer; IR spectra, Shimadzu FTIR-8100 spectrometer; EI-MS and high-resolution EI-MS, JEOL JMS-GCMATE mass spectrometer; FAB-MS and high-resolution FAB-MS, JEOL JMS-SX 102A mass spectrometer; ¹H-NMR spectra, JNM-LA500 (500 MHz) spectrometer; ¹³C-NMR spectra, JNM-LA500 (125 MHz) spectrometer with tetramethylsilane as an internal standard; and HPLC detector, Shimadzu RID-6A refractive index detector.

The following experimental conditions were used for chromatography: ordinary-phase silica gel column chromatography, Silica gel BW-200 (Fuji Silysia Chemical, Ltd., 150–350 mesh); reverse-phase silica gel column chromatography, Chromatorex ODS DM1020T (Fuji Silysia Chemical, Ltd., 100–200 mesh); TLC, precoated TLC plates with Silica gel 60F₂₅₄ (Merck, 0.25 mm) (ordinary phase) and Silica gel RP-18 F_{254S} (Merck, 0.25 mm) (reverse phase); reverse-phase HPTLC, precoated TLC plates with Silica gel RP-18 F_{254S} (Merck, 0.25 mm); and detection was achieved by spraying with 1% Ce(SO₄)₂–10% aqueous H₂SO₄ followed by heating.

Plant Material The whole plant of *Saussurea medusa* MAXIM. was purchased from Kunming, Yunnan province, China in September 2001, and the botanical identification was kindly undertaken by Dr. Seichi Yamaji (Toyama Pharmaceutical and Medical University, Japan).^{3,4)} A voucher specimen (2001.09. Yunnan-01) of this plant is on file in our laboratory.

Extraction and Isolation The dried whole plant of *S. medusa* (2.70 kg) was powdered and extracted with 80% aqueous acetone three times at room temperature. Evaporation of the solvent under reduced pressure provided the aqueous acetone extract (300 g, 11.1%). This extract (286 g) was partitioned in an EtOAc–H₂O (1:1, v/v) mixture to furnish an EtOAc-soluble fraction and aqueous layer. The aqueous layer was extracted with *n*-BuOH and removal of the solvents *in vacuo* from the EtOAc-, *n*-BuOH-, and H₂O-soluble fractions yielded 72 g (2.8%), 60 g (2.3%), and 154 g (6.0%) of the residues, respectively.

Normal-phase silica gel column chromatography [2.2 kg, *n*-hexane–EtOAc (50:1→20:1→10:1→5:1→2:1→1:1, v/v)→MeOH] of the EtOAc-soluble fraction (72 g) gave eight fractions [Fr. 1 (0.99 g), Fr. 2 (2.60 g), Fr. 3 (7.62 g), Fr. 4 (0.85 g), Fr. 5 (8.92 g), Fr. 6 (9.03 g), Fr. 7 (24.40 g), and Fr. 8 (16.10 g)]. Fraction 4 (0.85 g) was separated by reversed-phase silica gel column chromatography [25 g, MeOH–H₂O (70:30→85:15→98:2, v/v)→MeOH] to furnish four fractions [Fr. 4-1 (58 mg), Fr. 4-2 (83 mg), Fr. 4-3 (242 mg), Fr. 4-4 (459 mg)]. Fraction 4-3 (240 mg) was separated by HPLC [YMC-Pack ODS-A (YMC Co., Ltd., Kyoto, Japan, 250×20 mm i.d.), MeOH–H₂O (95:5, v/v)] to give (+)-glutinol (13.9 mg, 0.0006%). Fraction 5 (8.92 g) was subjected to reversed-phase silica gel column chromatography [240 g, MeOH–H₂O (75:25→85:15→98:2, v/v)→MeOH] to furnish four fractions [Fr. 5-1 (0.12 g), Fr. 5-2 (4.72 g), Fr. 5-3 (3.43 g), Fr. 5-4 (0.50 g)]. Fraction 5-1 (120 mg) was separated by HPLC [MeOH–H₂O (77:23, v/v)] to give (–)-2-oxoisodauc-5-en-12-al (5.4 mg, 0.0002%). Fraction 5-2 (630 mg) was purified by HPLC [MeOH–H₂O (90:10, v/v)] to give linoleic acid (118 mg, 0.035%), hexadecanoic acid (178 mg, 0.053%), and oleic acid (33.7 mg, 0.0093%). Fraction 5-3 (590 mg) was separated by HPLC [MeOH–H₂O (98:2, v/v)] to give linoleic acid (19.7 mg, 0.0044%), hexadecanoic acid (94.7 mg, 0.022%), octadecanoic acid (40.3 mg, 0.0092%), eicosanoic acid (12.1 mg, 0.0028%), eicosanol (4.8 mg, 0.0011%), docosanoic acid (10.8 mg, 0.0025%), tricosanoic acid (3.0 mg, 0.0007%), lupeol (31.8 mg, 0.0074%), and α-amyrin (37.2 mg, 0.0086%). Fraction 6 (9.03 g) was separated by reversed-phase silica gel column chromatography [280 g, MeOH–H₂O (75:25→85:15→98:2, v/v)→MeOH] to furnish four fractions [Fr. 6-1 (0.62 g), Fr. 6-2 (2.40 g), Fr. 6-3 (4.03 g), Fr. 6-4 (1.90 g)]. Fraction 6-2 (1.08 g) was separated by HPLC [MeOH–H₂O (87:13, v/v)] to give linolenic acid (305 mg, 0.027%) and linoleic acid (277 mg, 0.025%). Fraction 6-3 (1.05 g) was purified by HPLC [MeOH–H₂O (98:2, v/v)] to give clonasterol (16.6 mg, 0.0026%). Fraction 7 (24.40 g) was subjected to reversed-phase silica gel column chromatography [730 g, MeOH–H₂O (40:60→55:45→70:30→85:15→98:2, v/v)→MeOH] to

furnish nine fractions [Fr. 7-1 (1.18 g), Fr. 7-2 (1.48 g), Fr. 7-3 (1.44 g), Fr. 7-4 (3.00 g), Fr. 7-5 (2.58 g), Fr. 7-6 (1.08 g), Fr. 7-7 (3.20 g), Fr. 7-8 (3.34 g), Fr. 7-9 (6.88 g)]. Fraction 7-2 (1.48 g) was separated by HPLC [MeOH-H₂O (40:60, v/v)] to give 3-methoxy-4-hydroxyphenylethanol (51.9 mg, 0.0021%). Fraction 7-4 (2.76 g) was separated by normal-phase silica gel column chromatography [276 g, CHCl₃:MeOH:H₂O (50:3:1)→30:3:1→10:3:1, lower layer→6:4:1, v/v/v]→MeOH] to give seven fractions [Fr. 7-4-1 (99 mg), Fr. 7-4-2 (185 mg), Fr. 7-4-3 (52 mg), Fr. 7-4-4 (513 mg), Fr. 7-4-5 [=arctiin (**22**, 1359 mg, 0.054%)], Fr. 7-4-6 (465 mg), and Fr. 7-4-7 (45 mg)]. Fraction 7-4-1 (99 mg) was separated by HPLC [CH₃CN-H₂O (25:75, v/v)] to give (+)-pinoresinol (**16**, 8.9 mg, 0.0004%) and (+)-syringaresinol (**18**, 10.8 mg, 0.0004%). Fraction 7-4-4 (513 mg) was purified by HPLC [MeOH-H₂O (42:58, v/v)] to furnish azelaic acid (59.3 mg, 0.0024%) and (-)-secoisolaricresinol (**20**, 16.8 mg, 0.0007%). Fraction 7-5 (2.58 g) was separated by normal-phase silica gel column chromatography [250 g, CHCl₃-MeOH-H₂O (10:3:1→7:3:1, lower layer→6:4:1, v/v/v)→MeOH] to give eight fractions [Fr. 7-5-1 [=(-)-arctigenin (**21**, 939 mg, 0.038%)], Fr. 7-5-2 (385 mg), Fr. 7-5-3 (122 mg), Fr. 7-5-4 (108 mg), Fr. 7-5-5 (229 mg), Fr. 7-5-6 (340 mg), Fr. 7-5-7 (149 mg), and Fr. 7-5-8 (168 mg)]. Fraction 7-5-4 (108 mg) was separated by HPLC [MeOH-H₂O (60:40, v/v)] to give luteolin (**7**, 32.0 mg, 0.0013%). Fraction 7-5-6 (340 mg) was purified by HPLC [MeOH-H₂O (55:45, v/v)] to give quercetin (**13**, 150 mg, 0.0060%). Fraction 7-6 (1.08 g) was separated by HPLC [MeOH-H₂O (60:40, v/v)] to give apigenin (**3**, 53.7 mg, 0.0021%), and **7** (140 mg, 0.0056%).

The *n*-BuOH-soluble fraction (37 g) was subjected to normal-phase silica gel column chromatography [1.1 kg, CHCl₃-MeOH-H₂O (15:3:1)→10:3:1→7:3:1, lower layer→6:4:1, v/v/v]→MeOH] to furnish nine fractions [Fr. 1 (0.27 g), Fr. 2 (0.31 g), Fr. 3 (1.74 g), Fr. 4 (1.38 g), Fr. 5 (5.64 g), Fr. 6 (4.18 g), Fr. 7 (8.57 g), Fr. 8 (13.53 g), and Fr. 9 (1.21 g)]. Fraction 3 (1.74 g) was subjected to reversed-phase silica gel column chromatography [55 g, MeOH-H₂O (30:70→50:50, v/v)→MeOH] to provide five fractions [Fr. 3-1 (0.51 g), 3-2 (0.05 g), 3-3 (**22**, 983 mg, 0.062%), 3-4 (0.02 g), 3-5 (0.11 g), and Fr. 3-6 (0.06 g)]. Fr. 3-2 (0.05 g) was purified by HPLC [MeOH-H₂O (40:60, v/v)] to give (+)-syringaresinol *O*-β-D-glucopyranoside (**19**, 16.7 mg, 0.0011%). Fr. 3-5 (0.11 g) was purified by HPLC [MeOH-H₂O (50:50, v/v)] to give sitosterol 3β-*O*-β-D-glucopyranoside (72.5 mg, 0.0046%). Fraction 4 (1.38 g) was subjected to reversed-phase silica gel column chromatography [41 g, MeOH-H₂O (30:70→50:50, v/v)→MeOH] to furnish three fractions [Fr. 4-1 (0.92 g), 4-2 (0.36 g), and 4-3 (0.10 g)]. Fraction 4-1 (0.92 g) was further purified by HPLC [MeOH-H₂O (10:90, 25:75, or 30:70, v/v)] to give saussureosides A (**1**, 12.0 mg, 0.0007%) and B (**2**, 59.8 mg, 0.0038%), (+)-pinoresinol *O*-β-D-glucopyranoside (**17**, 22.9 mg, 0.0026%), junipediol A (17.0 mg, 0.0011%), benzyl *O*-β-D-glucopyranoside (44.4 mg, 0.0028%), *p*-anisyl *O*-β-D-glucopyranoside (9.1 mg, 0.0006%), phenethyl *O*-β-D-glucopyranoside (11.5 mg, 0.0007%), (*Z*)-3-hexenyl *O*-β-D-glucopyranoside (16.0 mg, 0.0010%), and thymidine (41.0 mg, 0.0026%). Fraction 5 (5.64 g) was subjected to reversed-phase silica gel column chromatography [170 g, MeOH-H₂O (20:80→50:50, v/v)→MeOH] to provide seven fractions [Fr. 5-1 (2.31 g), Fr. 5-2 (0.43 g), Fr. 5-3 (0.40 g), Fr. 5-4 (0.39 g), Fr. 5-5 (0.34 g), Fr. 5-6 (1.38 g), and Fr. 5-7 (0.32 g)]. Fraction 5-2 (80 mg) was purified by HPLC [MeOH-H₂O (30:70, v/v)] to give syringin (22.0 mg, 0.0037%). Fraction 5-5 (340 mg) was purified by HPLC [MeOH-H₂O (45:55, v/v)] to furnish quercetin 3-*O*-β-D-glucopyranoside (**14**, 60.2 mg, 0.0038%). Fraction 5-6 (374 mg) was separated by HPLC [MeOH-H₂O (40:60, v/v)] to give apigenin 7-*O*-β-D-glucopyranoside (**4**, 7.0 mg, 0.0016%), apigenin 4'-*O*-β-D-glucopyranoside (**5**, 38.4 mg, 0.0089%), luteolin 7-*O*-β-D-glucopyranoside (**8**, 7.6 mg, 0.0018%), luteolin 4'-*O*-β-D-glucopyranoside (**9**, 120.4 mg, 0.028%), chrysoeriol 7-*O*-β-D-glucopyranoside (**11**, 14.6 mg, 0.0034%), and **14** (69.5 mg, 0.016%). Fraction 6 (4.18 g) was subjected to reversed-phase silica gel column chromatography [140 g, MeOH-H₂O (30:70→50:50→70:30, v/v)→MeOH] to provide four fractions [Fr. 6-1 (1.50 g), Fr. 6-2 (2.30 g), Fr. 6-3 (0.26 g), and Fr. 6-4 (0.08 g)]. Fraction 6-1 (0.50 g) was separated by HPLC [MeOH-H₂O (71:83, v/v)] to give L-phenylalanine (9.7 mg, 0.0018%). Fraction 6-2 (0.50 g) was separated by HPLC [MeOH-H₂O (40:60, v/v)] to give **6** (14.6 mg, 0.0054%), **8** (61.7 mg, 0.023%), **9** (39.0 mg, 0.014%), chrysoeriol 7-*O*-rutoside (**12**, 8.1 mg, 0.0035%), **14** (160 mg, 0.059%), and isorhamnetin 3-*O*-rutoside (**15**, 9.6 mg, 0.0035%). Fraction 7 (8.57 g) was subjected to reversed-phase silica gel column chromatography [260 g, MeOH-H₂O (30:70→50:50→70:30, v/v)→MeOH] to provide five fractions [Fr. 7-1 (3.51 g), Fr. 7-2 (2.96 g), Fr. 7-3 (1.14 g), Fr. 7-4 (0.35 g), and Fr. 7-5 (0.36 g)]. Fraction 7-2 (350 mg) was separated by HPLC [MeOH-H₂O (40:60, v/v)] to afford **7** (18.0 mg, 0.0096%), **8** (18.0 mg, 0.0096%), luteolin 7-*O*-

rutoside (**10**, 42.3 mg, 0.023%), and 5-*O*-caffeoylquinic acid methyl ester (**25**, 27.0 mg, 0.015%). Fraction 8 (15.53 g) was subjected to reversed-phase silica gel column chromatography [470 g, MeOH-H₂O (10:90→25:75→40:60→50:50, v/v)→MeOH] to provide four fractions [Fr. 8-1 (9.16 g), Fr. 8-2 (2.65 g), Fr. 8-3 (1.15 g), and Fr. 8-4 (1.85 g)]. Fraction 8-2 (600 mg) was separated by HPLC [MeOH-H₂O (33:67, v/v)] to furnish 3-*O*-caffeoylquinic acid methyl ester (**23**, 4.0 mg, 0.0011%), 4-*O*-caffeoylquinic acid methyl ester (**24**, 4.2 mg, 0.0012%), and **25** (15.4 mg, 0.0043%).

The known compounds were identified by comparison of their physical data ([α]_D, IR, ¹H-NMR, ¹³C-NMR, MS) with reported values^{18–36,38–45} or commercial samples.³⁷

Saussureoside A (1): A white powder, [α]_D²⁴ -45.0° (*c*=0.80, MeOH). High-resolution positive-ion FAB-MS: Calcd for C₁₉H₃₀O₉Na (*M*+Na)⁺ 425.1788; Found 425.1783. CD [MeOH, nm (Δε)]: 229 (-4.36). UV [MeOH, nm (log ε)]: 229 (4.07). IR (KBr): 3410, 2932, 1684, 1671, 1260, 1078, 756 cm⁻¹. ¹H-NMR (500 MHz, pyridine-*d*₅) δ: given in Table 2. ¹³C-NMR (125 MHz, pyridine-*d*₅) δ_C: given in Table 3. Positive-ion FAB-MS: *m/z* 425 (*M*+Na)⁺.

Saussureoside B (2): A white powder, [α]_D²⁶ -39.5° (*c*=0.98, MeOH). High-resolution positive-ion FAB-MS: Calcd for C₁₉H₃₀O₈Na (*M*+Na)⁺ 409.1838; Found 409.1843. CD [MeOH, nm (Δε)]: 227 (+3.21), 279 (+3.16), 315 (-3.06). UV [MeOH, nm (log ε)]: 227 (3.89), 279 (3.78). IR (KBr): 3432, 2932, 1671, 1655, 1364, 1262, 1076 cm⁻¹. ¹H-NMR (500 MHz, pyridine-*d*₅) δ: given in Table 2. ¹³C-NMR (125 MHz, pyridine-*d*₅) δ_C: given in Table 3. Positive-ion FAB-MS: *m/z* 409 (*M*+Na)⁺.

Acid Hydrolysis of 1 and 2 A solution of **1** and **2** (2.0 mg of each) in 1 mL HCl (0.5 mL) was heated under reflux for 3 h. After cooling, the reaction mixture was poured into ice-water and neutralized with Amberlite IRA-400 (OH⁻ form), and the resin was removed by filtration. Then, the filtrate was extracted with EtOAc. The aqueous layer was subjected to HPLC analysis under the following conditions: HPLC column, Kaseisorb LC NH₂-60-5, 4.6 mm i.d.×250 mm (Tokyo Kasei Co., Ltd., Tokyo, Japan); detection, optical rotation [Shodex OR-2 (Showa Denko Co., Ltd., Tokyo)]; mobile phase, CH₃CN-H₂O (75:25, v/v); flow rate 0.8 mL/min; column temperature, room temperature. Identification of D-glucose present in the aqueous layer was carried out by comparison of its retention time and optical rotation with those of an authentic sample. *t*_R: 12.3 min (positive optical rotation).

Enzymatic Hydrolysis of 1 and 2 with β-Glucosidase A solution of **1** (8.0 mg, 0.020 mmol) in H₂O (1.0 mL) was treated with β-glucosidase (8.0 mg, from Almond, Oriental Yeast Co., Ltd., Japan) and the solution was stirred at 37 °C for 4 d. After EtOH was added to the reaction mixture, the solvent was removed under reduced pressure and the residue was purified by HPLC [MeOH-H₂O (40:60, v/v)] to furnish **1a** (3.4 mg, 71%). Through a similar procedure, **2a** (4.8 mg, 75%) was prepared from **2** (9.4 mg, 0.024 mol).

1a: A white powder, [α]_D¹⁸ -64.6° (*c*=0.10, MeOH). High-resolution EI-MS: Calcd for C₁₃H₂₀O₄ (*M*⁺) 240.1361; Found 240.1354. CD [MeOH, nm (Δε)]: 229 (+3.92). UV [MeOH, nm (log ε)]: 229 (3.92), UV [EtOH, nm (log ε)]: 232 (4.05). IR (KBr): 3432, 2924, 1682, 1662, 1258, 754 cm⁻¹. ¹H-NMR (500 MHz, CDCl₃) δ: 0.99, 1.17, 1.31, 2.28 (each 3H, all s, 12, 13, 11, 10-H₃), [1.30 (1H, dd, *J*=3.4, 12.5 Hz), 1.60 (1H, dd, *J*=12.5, 12.5 Hz), 2-H₃], 3.96 (1H, ddd, *J*=3.4, 3.4, 12.5 Hz, 3-H), 4.02 (1H, d, *J*=3.4 Hz, 4-H), 6.27, 7.01 (1H each, both d, *J*=15.9 Hz, 8, 7-H). ¹³C-NMR (125 MHz, CDCl₃) δ_C: given in Table 3. EI-MS *m/z* (%): 240 (*M*⁺, 2), 179 (3), 151 (3), 123 (100).

2a: A white powder, [α]_D²⁵ -70.7° (*c*=0.25, MeOH). High-resolution EI-MS: Calcd for C₁₃H₂₀O₃ (*M*⁺) 224.1412; Found 224.1415. CD [MeOH, nm (Δε)]: 225 (+3.21), 274 (+3.28), 312 (-2.94). UV [MeOH, nm (log ε)]: 220 (3.87), 281 (3.82). IR (KBr): 3410, 2934, 1674, 1609, 1363, 1258, 1063 cm⁻¹. ¹H-NMR (500 MHz, CDCl₃) δ: 1.05, 1.12, 2.07, 2.32 (each 3H, all s, 13, 12, 11, 10-H₃), [1.78 (1H, dd, *J*=3.4, 12.5 Hz), 2.07 (1H, dd, *J*=12.5, 12.5 Hz), 2-H₃], 4.18 (1H, ddd, *J*=3.4, 3.4, 12.5 Hz, 3-H), 4.30 (1H, d, *J*=3.4 Hz, 4-H), 6.32, 7.39 (1H each, both d, *J*=16.5 Hz, 8, 7-H). ¹³C-NMR (125 MHz, CDCl₃) δ_C: given in Table 3. EI-MS *m/z* (%): 224 (*M*⁺, 2), 206 (16), 191 (100).

Preparation of the (R)-MTPA Esters (1b, 2b) and (S)-MTPA Esters (1c, 2c) from 1a and 2a A solution of **1a** or **2a** (0.7 mg each) in CH₂Cl₂ (1.0 mL) was treated with (*R*)-α-methoxy-α-trifluoromethylphenylacetic acid [(*R*)-MTPA, 3.4 mg] in the presence of 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC·HCl, 2.9 mg) and 4-dimethylaminopyridine (4-DMAP, 1.1 mg), and the mixture was stirred at room temperature for 10 h. After cooling, the reaction mixture was poured into ice-water and the whole reaction mixture was extracted with EtOAc. The EtOAc extract was successively washed with 5% aqueous HCl, saturated aqueous NaHCO₃, and

brine, then dried over MgSO_4 powder and filtered. Removal of the solvent from the filtrate under reduced pressure furnished a residue, which was separated by HPLC [$\text{MeOH-H}_2\text{O}$ (70:30, v/v)] to give **1b** (0.7 mg, 53%) or **2b** (0.7 mg, 51%), respectively. Using a similar procedure, (*S*)-MTPA esters [**1c** (1.0 mg, 53%) or **2c** (1.2 mg, 51%)] were obtained from **1a** or **2a** (1.0 mg each), respectively, using (*S*)-MTPA (4.9 mg), EDC·HCl (4.2 mg), and 4-DMAP (1.6 mg).

1b: $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 1.06, 1.16, 1.26, 2.27 (each 3H, all s, 12, 13, 11, 10- H_3), [1.44 (1H, dd, $J=3.4$, 12.5 Hz), 1.91 (1H, dd, $J=12.5$, 12.5 Hz), 2- H_2], 3.54 (3H, s, $-\text{OCH}_3$), 4.11 (1H, d, $J=3.4$ Hz, 4-H), 5.28 (1H, ddd, $J=3.4$, 3.4, 12.5 Hz, 3-H), 6.27, 6.96 (1H each, both d, $J=16.5$ Hz, 8, 7-H), [7.40—7.42 (3H, m), 7.48—7.50 (2H, m), Ph-H].

1c: $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 1.06, 1.15, 1.30, 2.29 (each 3H, all s, 12, 13, 11, 10- H_3), [1.44 (1H, dd, $J=3.4$, 12.5 Hz), 1.83 (1H, dd, $J=12.5$, 12.5 Hz), 2- H_2], 3.52 (3H, s, $-\text{OCH}_3$), 4.21 (1H, d, $J=3.4$ Hz, 4-H), 5.28 (1H, ddd, $J=3.4$, 3.4, 12.5 Hz, 3-H), 6.29, 6.98 (1H each, both d, $J=16.5$ Hz, 8, 7-H), [7.42—7.44 (3H, m), 7.50—7.51 (2H, m), Ph-H].

2b: $^1\text{H-NMR}$ (500 MHz, pyridine- d_5) δ : 1.05, 1.15, 1.99, 2.33 (each 3H, all s, 13, 12, 11, 10- H_3), [1.91 (1H, dd, $J=3.4$, 12.5 Hz), 2.45 (1H, dd, $J=12.5$, 12.5 Hz), 2- H_2], 3.74 (3H, s, $-\text{OCH}_3$), 4.47 (1H, d, $J=3.4$ Hz, 4-H), 5.50 (1H, ddd, $J=3.4$, 3.4, 12.5 Hz, 3-H), 6.32, 7.35 (1H each, both d, $J=16.5$ Hz, 8, 7-H), [7.42—7.47 (3H, m), 7.94 (2H, d, $J=7.6$ Hz), Ph-H].

2c: $^1\text{H-NMR}$ (500 MHz, pyridine- d_5) δ : 0.96, 1.13, 2.01, 2.33 (each 3H, all s, 13, 12, 11, 10- H_3), [1.75 (1H, dd, $J=3.4$, 12.5 Hz), 2.38 (1H, dd, $J=12.5$, 12.5 Hz), 2- H_2], 3.81 (3H, s, $-\text{OCH}_3$), 4.63 (1H, d, $J=3.4$ Hz, 4-H), 5.52 (1H, ddd, $J=3.4$, 3.4, 12.5 Hz, 3-H), 6.30, 7.32 (1H each, both d, $J=16.5$ Hz, 8, 7-H), [7.40—7.47 (3H, m), 8.00 (2H, d, $J=7.6$ Hz), Ph-H].

Bioassay. Inhibitory Effect on Aldose Reductase Activity Aldose reductase activity was assayed by the method described previously.^{52–57} The supernatant fluid of rat lens homogenate was used as a crude enzyme. The incubation mixture contained 135 mM Na, K-phosphate buffer (pH 7.0), 100 mM Li_2SO_4 , 0.03 mM NADPH, 1 mM DL-glyceraldehyde as a substrate, and 100 μl of enzyme fraction, with or without 25 μl of sample solution, in a total volume of 0.5 ml. The reaction was initiated by the addition of NADPH at 30 °C. After 30 min, the reaction was stopped by the addition of 150 μl of 0.5 M HCl. Then, 0.5 ml 6 M NaOH containing 10 mM imidazole was added, and the solution was heated at 60 °C for 20 min to convert NADP to a fluorescent product. Fluorescence was measured using a fluorophotometer (Luminescence Spectrometer LS50B, Perkin Elmer, U.K.) at an excitation wavelength of 360 nm and an emission wavelength of 460 nm.

Acknowledgments The authors would like to thank Dr. Seiichi Yamaji, Assistant Professor of the Institute of Natural Medicine, Toyama Pharmaceutical and Medical University, Japan for identifying the plant material. Part of this work was supported by the 21st COE program and the Academic Frontier Project of the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References and Notes

- Part XIV: Morikawa T., Sun B., Matsuda H., Wu L. J., Harima S., Yoshikawa M., *Chem. Pharm. Bull.*, **52**, 1194—1199 (2004).
- Shanghai Scientific and Technologic Press, "Dictionary of Traditional Chinese Medicines," Shogakkan, Tokyo, 1985, p. 1482.
- Yang R. P., Yamaji S., Komatsu K., Namba T., *Nat. Med.*, **51**, 134—147 (1997).
- Yang R. P., Komatsu K., Sato T., Namba T., Yamaji S., *J. Jpn. Bot.*, **78**, 315—329 (2003).
- Takasaki M., Konoshima T., Komatsu K., Tokuda H., Nishino H., *Cancer Lett.*, **158**, 53—59 (2000).
- Duan H., Takaishi Y., Momota H., Ohmoto Y., Taki T., *Phytochemistry*, **59**, 85—90 (2002).
- Fan C. Q., Yue J. M., *Bioorg. Med. Chem.*, **11**, 703—708 (2003).
- Jia Z., Fei H., Li Y., Zhu Z., *Lanzhou Daxue Xuebao, Ziran Kexueban*, **20**, 126, 128 (1984).
- Jia Z., Fei H., Li Y., Zhu Z., *Gaodeng Xuexiao Huaxue Xuebao*, **7**, 789—792 (1986).
- Jia Z., Gong N., Du M., *Gaodeng Xuexiao Huaxue Xuebao*, **11**, 202—204 (1986).
- Sun B., Morikawa T., Matsuda H., Tewtrakul S., Wu L. S., Harima S., Yoshikawa M., *J. Nat. Prod.*, **67**, 1464—1469 (2004).
- Matsuda H., Morikawa T., Xie H., Yoshikawa M., *Planta Med.*, **70**, 847—855 (2004).
- Tao J., Morikawa T., Ando S., Matsuda H., Yoshikawa M., *Chem. Pharm. Bull.*, **51**, 654—662 (2003).
- Morikawa T., Tao J., Ando S., Matsuda H., Yoshikawa M., *J. Nat. Prod.*, **66**, 638—645 (2003).
- Tao J., Morikawa T., Toguchida I., Ando S., Matsuda H., Yoshikawa M., *Bioorg. Med. Chem.*, **10**, 4005—4012 (2002).
- Morikawa T., Matsuda H., Toguchida I., Ueda K., Yoshikawa M., *J. Nat. Prod.*, **65**, 1468—1474 (2002).
- Matsuda H., Morikawa T., Tao J., Ueda K., Yoshikawa M., *Chem. Pharm. Bull.*, **50**, 208—215 (2002).
- Kubo M., Sasaki H., Endo T., Taguchi H., Yosioka I., *Chem. Pharm. Bull.*, **34**, 3097—3101 (1986).
- Nawwar M. A. M., Hussein S. A. M., Merfort I., *Phytochemistry*, **37**, 1175—1177 (1994).
- Wang M., Simon J. E., Aviles I. F., He K., Zheng Q. Y., Tadmor Y., *J. Agric. Food Chem.*, **51**, 601—608 (2003).
- Li H. Y., Koike K., Ohmoto T., *Phytochemistry*, **35**, 751—756 (1994).
- Tomas F., Ferreres F., Barberan F. A. T., *J. Nat. Prod.*, **48**, 506—507 (1985).
- Markham K. R., Ternai B., Stanley R., Geiger H., Marby T. J., *Tetrahedron*, **34**, 1389—1397 (1978).
- Fukunaga T., Kajikawa I., Nishiya K., Watanabe Y., Suzuki N., Takeya K., Itokawa H., *Chem. Pharm. Bull.*, **36**, 1185—1189 (1986).
- Chiba M., Hisada S., Nishibe S., Thieme H., *Phytochemistry*, **19**, 335—336 (1980).
- Kobayashi H., Karasawa H., Miyase T., Fukushima S., *Chem. Pharm. Bull.*, **33**, 1452—1457 (1985).
- Rahman M. M. A., Dewick P. M., Jackson D. E., Lucas J. A., *Phytochemistry*, **29**, 1971—1980 (1990).
- Fuchs C., Spiteller G., *J. Mass Spectrometry*, **31**, 602—608 (1996).
- Flamini G., Antognoli E., Morelli I., *Phytochemistry*, **57**, 559—564 (2001).
- Rumbero-Sanchez A., Vazquez P., *Phytochemistry*, **30**, 311—313 (1991).
- Greca M. D., Ferrara M., Fiorentino A., Monaco P., Previtera L., *Phytochemistry*, **49**, 1299—1304 (1998).
- Comte G., Allais D. P., Chulia A. J., Vercauteren J., Pinaud N., *Phytochemistry*, **44**, 1169—1173 (1997).
- Miyase T., Ueno A., Takizawa N., Kobayashi H., Oguchi H., *Chem. Pharm. Bull.*, **36**, 2475—2484 (1988).
- Miyase T., Ueno A., Takizawa N., Kobayashi H., Karasawa H., *Chem. Pharm. Bull.*, **35**, 1109—1117 (1987).
- Kitajima J., Ishikawa T., Tanaka Y., Ono M., Ito Y., Nohara T., *Chem. Pharm. Bull.*, **46**, 1587—1590 (1998).
- Mizutani K., Yuda M., Tanaka O., Saruwatari Y., Fuwa T., Jia M. R., Ling Y. K., Pu X. F., *Chem. Pharm. Bull.*, **36**, 2689—2690 (1988).
- Those known constituents were identified by comparison of their physical data with commercially obtained samples.
- Voutquenne L., Lavaud C., Massiot G., Sevenet T., Hadi H. A., *Phytochemistry*, **50**, 63—69 (1999).
- Deyama T., Ikawa T., Kitagawa S., Nishibe S., *Chem. Pharm. Bull.*, **35**, 1785—1789 (1987).
- Nawwar M. A. M., Buddrus J., Bauer H., *Phytochemistry*, **21**, 1755—1758 (1982).
- Fonseca S. F., Campello J. D. P., Barata L. E. S., Rúveda E. A., *Phytochemistry*, **17**, 499—502 (1978).
- Kodaira H., Ishikawa M., Komoda Y., Nakajima T., *Chem. Pharm. Bull.*, **29**, 2391—2393 (1981).
- Hansson T., Wickberg B., *J. Org. Chem.*, **57**, 5370—5376 (1992).
- Kitajima J., Arai M., Tanaka Y., *Chem. Pharm. Bull.*, **42**, 608—610 (1994).
- Furukawa S., Takagi N., Ikeda T., Ono M., Nafady A. M., Nohara T., Sugimoto H., Doi S., Yamada H., *Chem. Pharm. Bull.*, **50**, 439—440 (2002).
- Reynolds W. F., McLean S., Tay L. L., Yu M., Enriquez R. G., Estwick D. M., Pascoe K. O., *Magn. Res. Chem.*, **35**, 455—462 (1997).
- The ^1H - and ^{13}C -NMR spectra of **1**, **2**, **1a**, and **2a** were assigned with the aid of distortionless enhancement by polarization transfer (DEPT), homo- and hetero-correlation spectroscopy (^1H - ^1H , ^{13}C - ^1H COSY), and HMBC experiments.
- Yokosuka A., Mimaki Y., Sashida Y., *J. Nat. Prod.*, **63**, 1239—1243 (2000).
- Ohtani I., Kusumi T., Kashman Y., Kakisawa H., *J. Am. Chem. Soc.*, **113**, 4092—4096 (1991).
- Although the structure investigation of **2** was reported,⁵¹ the absolute stereostructure was not determined. This is the first report of the struc-

- ture elucidation including the absolute stereochemistry of **2**.
- 51) Shimizu S., Miyase T., Ueno A., Usmanghani K., *Phytochemistry*, **28**, 3399—3402 (1989).
- 52) Matsuda H., Morikawa T., Toguchida I., Yoshikawa M., *Chem. Pharm. Bull.*, **50**, 788—795 (2002).
- 53) Yoshikawa M., Morikawa T., Murakami T., Toguchida I., Harima S., Matsuda H., *Chem. Pharm. Bull.*, **47**, 340—345 (1999).
- 54) Matsuda H., Morikawa T., Toguchida I., Harima S., Yoshikawa M., *Chem. Pharm. Bull.*, **50**, 972—975 (2002).
- 55) Matsuda H., Morikawa T., Ueda H., Yoshikawa M., *Heterocycles*, **55**, 1499—1504 (2001).
- 56) Yoshikawa M., Murakami T., Ishiwada T., Morikawa T., Kagawa M., Higashi Y., Matsuda H., *J. Nat. Prod.*, **65**, 1151—1155 (2002).
- 57) Morikawa T., Kishi A., Pongpiriyadacha Y., Matsuda H., Yoshikawa M., *J. Nat. Prod.*, **66**, 1191—1196 (2003).