Communications

1. cat. CrCl₂ / cat. CrL* Mn, Me₃SiCl R¹CHO R²X + 2. Hvdrolvsis 2 1 3 *t*Bu tBu OН (R,R)-4*t*Bu fBu fRı HO *t*Bu

Highly Enantioselective Catalysts

A Highly Enantioselective Catalyst for the Asymmetric Nozaki–Hiyama–Kishi Reaction of Allylic and Vinylic Halides**

Albrecht Berkessel,* Dirk Menche, Christoph A. Sklorz, Michael Schröder, and Ian Paterson

Dedicated to Professor Emanuel Vogel on the occasion of his 75th birthday

The nucleophilic addition of organochromium reagents to aldehvdes (Scheme 1) is a versatile carbon-carbon bondforming reaction that is compatible with a wide range of functionality in both components.^[1] A large number of applications, particularly in the synthesis of structurally diverse natural products, have resulted. The organochromium reagents are easily prepared in situ by the oxidative addition of chromium(II) species to allyl, propargyl, aryl, and vinyl halides or triflates 2 (requiring catalytic amounts of a Ni^{II} salt for sp^2 carbon centers), and efficiently add to aldehydes 1 to give the corresponding alcohols **3** in good yields (Scheme 1).^[1] While the original procedure for the Nozaki-Hiyama-Kishi (NHK) reaction relies on the use of stoichiometric quantities of chromium(II) chloride, Fürstner et al. have developed a method requiring only catalytic amounts of the active Cr^{II} species; cheap and toxicologically benign manganese is used as the reducing agent, and a trialkylchlorosilane effects the perpetuation of the catalytic cycle.^[2] This important finding triggered numerous efforts to develop a catalytic and

[*] Prof. Dr. A. Berkessel, Dr. D. Menche, Dr. C. A. Sklorz, Dipl.-Chem. M. Schröder
Institut für Organische Chemie
Universität zu Köln
Greinstrasse 4, 50939 Köln (Germany)
Fax: (+49) 221-470-5102
E-mail: berkessel@uni-koeln.de
Prof. Dr. I. Paterson
University Chemical Laboratory
Lensfield Road, Cambridge CB2 1EW (United Kingdom)

[**] This work was supported by the European Community (Research Training Network "Combinatorial Catalysis") and by the Fonds der Chemischen Industrie. We thank Dr. Ira Fiolia for technical support. **Scheme 1.** Ligands for the catalytic asymmetric Nozaki–Hiyama–Kishi reaction; diaminocyclohexane-derived salen **4** and the novel ligand **5**, based on *endo*,*endo*-2,5-diaminonorbornane (DIANANE).

*t*Bú

(S,S)-5

*t*Bu

enantioselective version of this reaction.^[2-5] Nevertheless, only one such example appears to have been reported, namely by the research group of Umani-Ronchi, which uses the salen ligand 4, derived from trans-1,2-diaminocyclohexane (Scheme 1).^[5] Whereas the enantioselectivities observed for allyl chorides were high (for example, up to 84% ee in the addition to benzaldehyde), the use of the ligand 4 does not yet provide a broadly applicable synthetic method. In particular, the substrate spectrum is still fairly narrow; other halides such as allyl bromide and iodide were found to add with only moderate (52% ee) or no selectivity to benzaldehyde. As an asymmetric NHK process for generating enantiomerically enriched allylic alcohols would be extremely valuable, the omission of vinyl halides as substrates for this reaction is significant.^[5a-e] We now report our preliminary results using the novel salen ligand (S,S)-5 in NHK reactions, which surpasses the current limitations of 4 by promoting the catalytic addition of allylic and vinylic halides (and triflates) to aromatic and aliphatic aldehydes with synthetically useful levels of asymmetric induction.

Salen ligands used in asymmetric catalysis are mainly based on commercially available chiral 1,2-diamines and various salicylic aldehyde components. Comparatively little attention has been paid to significantly modifying the diamine backbone, for example, by attenuating the separation of the nitrogen atoms. We imagined that this parameter might play a particularly important role in further improving the catalytic properties and, thus, we devised a novel chiral diamine that exhibits a larger nitrogen-atom separation.^[6,7] We envisaged endo,endo-2,5-diamino-norbornane (9, DIANANE, Scheme 2) to be a promising candidate. This novel building block has the advantages of possessing C_2 symmetry and having a rigid hydrocarbon backbone. In our synthesis of 9, the diketone 6 is converted to a bis(endo diamine) by



Scheme 2. Enantiodivergent synthesis of (S,S)-**5** and (R,R)-**5**. a) HCO₂H, reflux, 84%;^[9] b) Jones reagent, 44%;^[9] c) NH₂OH·HCl, EtOH, reflux, 1 h; d) NiCl₂, NaBH₄, MeOH, -35 °C to -20 °C, 1 h, 70– 75% (over two steps); e) TsCl, NEt₃, CH₂Cl₂, -78 °C to RT; f) crystallization from MeOH, 35% (over two steps); g) preparative HPLC, Chiralpak AD, 50 cm, 5 cm diameter, flow: 80 mLmin⁻¹, eluent: hexane/isopropanol 60:40, retention times: 24–33 min for (R,R)-**8** and 46.5– 62.5 min for (S,S)-**8**; h) Li, NH₃, -33 °C, 80 min, 85%; i) 3,5-di-*tert*butyl salicylic aldehyde, MeOH, RT, 30 min, 76–81%.

treatment of the corresponding dioxime with NaBH₄/NiCl₂ (Scheme 2). Consequently, the pure enantiomers of DIA-NANE 9 (that is, (S,S)-9 and (R,R)-9) can be prepared from the enantiomerically pure diketone 6 and *ent*-6, respectively.

However, instead of relying on the known enantioselective, but multistep approach to 6/ent-6 from norbornadiene (7),^[8] we employed the readily available racemic form of the diketone (*rac*-6),^[9] and resolved the racemic mixture of DIANANE at a later stage. As summarized in Scheme 2, the *N*,*N'*-bistosylates of DIANANE (8 and *ent*-8) were efficiently separated by preparative HPLC on a chiral stationary phase. Absolute configurations of the crystalline sulfonamides 8 and *ent*-8 were determined by X-ray crystallography.^[10] Reductive *N*-detosylation followed by condensation with 3,5-di-*tert*butyl salicylic aldehyde afforded the desired salen ligands (*S*,*S*)-5 and (*R*,*R*)-5, respectively.

An evaluation of the catalytic performance of the (S,S)-5 and (R,R)-5 ligands was performed. Typical reaction conditions^[2,5] comprised 10 mol% of CrCl₃, an excess of Mn and Me₃SiCl, and in situ preparation of the Cr^{II} complex, in the presence of NEt₃ and MeCN as solvent (Table 1). The addition of allyl bromide to benzaldehyde proceeded asymmetrically (46% ee, room temperature). After optimization of the reaction parameters (solvent, concentration, and temperature), the corresponding homoallylic alcohol was obtained with high enantioselelectivity (90% ee) and good yield (72%; Table 1, entry 1). THF proved to be the best solvent among those tested (DMF, DME, CH₃CN), while a catalyst concentration of 0.025 M gave the best results. Optimal enantioselectivity was observed for the allyl bromide/benzaldehyde system at 5°C (90% ee). A variety of different substrates was screened under these conditions. In

....

 Table 1: Catalytic, enantioselective Nozaki–Hiyama–Kishi-type additions in the presence of ligand (S,S)-5.

 CrCl₃, (0.1 equiv), (S,S)-5, (0.1 equiv), NEt₃ (0.2 equiv)^[a]

	R ¹ CHO + R ² X ⁻ (1 equiv) (1.5 equiv)					
			Mn (3 equiv), Me ₃ SiCl (1.5 equiv) THF		R^1 R^2	
Entry	Aldehyde	Halide	Product	ee [%]	Yield [%] ^[b]	Reaction temperature ^[c]
1	O Ph H	Br	HO Ph	90 ^[d,e]	72	5°C
2	O Ph H		HO Ph	31 ^[d,f]	[g]	RT
3	O Ph H	CI	HO Ph	79 ^[d,e]	76	RT
4	O Ph H	CI	HO Ph	54 ^[e,h]	78	RT
5	O Ph H	Br	HO Ph	64 ^[f,h]	[g]	RT
6	РМВО Н	Br	Но РМВО	92 ^[e,h]	69 ^[1]	10°C
7	РМВО Н	$\sim\sim\sim$	PMBO * R	75 ^[h,j]	59 ^[i]	15 °C
8	РМВО Н			61 ^[h,k,l]	54 ^[i]	20°C

[a] For entries 7 and 8, the reaction was run in the presence of 0.02 equiv of NiCl₂. [b] Yield after flash chromatography. [c] Reaction temperature was optimized for entries 1, 2, 5, 6, 7, and 8. [d] Enantiomeric excess was determined by gas chromatography (GC) of the corresponding TMS ethers on a chiral stationary phase (Macherey–Nagel: Lipodex A, 95 °C). [e] Absolute configurations were assigned by comparison of optical rotations with literature data.^[5d, 11, 12] [f] Absolute configuration based on GC/HPLC data (that is, comparison of the products of entries 2 and 1 and of the products of entries 3 and 5 on an analytical scale). [g] Not determined. [h] Enantiomeric excess determined by HPLC on a chiral phase (Daicel: Chiralcel OD-H). [i] Some debenzylation occured as a side reaction. [j] Absolute configuration determined by 1) oxidative cleavage (ozonolysis), 2) reduction to 1,2,4-butanetriol, 3) GC co-injection with a sample of known absolute configuration.^[13] [k] Reaction was performed with (*R*,*R*)-**5**. [l] Assignment of absolute configuration in analogy to entry 7.

Angew. Chem. Int. Ed. 2003, 42, No. 9 © 2003 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim 1433-7851/03/4209-1033 \$ 20.00+.50/0

Communications

contrast to the salen ligand 4, ligand 5 was also able to effect, for the first time, an enantioselective addition of allyl iodide (Table 1, entry 2). Allyl chloride added with similar selectivity (84% ee using 4,^[5] versus 79% ee using DIANANE-based 5; Table 1, entry 3). Of the three allyl halides tested, the bromide thus afforded the best enantioselectivity. The related β methylallyl halides (Table 1, entries 4 and 5) reacted just as smoothly as the allyl halides. However, the observed enantioselectivities for these compounds (54% and 64% ee for the chloride and bromide, respectively) were not as high as for allyl bromide (90% ee). As a model for projected applications in polyketide natural-product synthesis, we tested PMBprotected 3-hydroxypropanal (PMB = para-methoxybenzyl), shown in entry 6 of Table 1.^[12,14,15] Again, after optimizing the reaction temperature, coupling with allyl bromide occurred with high enantioselectivity (92% ee).^[16]

The encouraging results achieved with allylic halides prompted us to examine the use of our catalytic NHK process with vinyl iodides and triflates for the enantioselective synthesis of allylic alcohols. As expected, 2 mol% of Ni^{II} (in the form of NiCl₂) was required for the coupling reaction to occur efficiently.^[2] Under these reaction conditions, the addition of E-1-iodohex-1-ene^[17] to the PMB-protected 3hydroxypropanal test system afforded the corresponding Eallylic alcohol adduct in an unoptimized 59% yield with 75% ee (Table 1, entry 7). Likewise, the vinyl triflate^[18] (entry 8, Table 1) added to this same aldehyde to produce the isomeric allylic alcohol with 61 % ee. Notably, these last two addition reactions represent the first examples of synthetically useful levels of asymmetric induction being realized for catalytic, enantioselective NHK reactions of vinylic halide and triflate substrates.

In summary, we have demonstrated that our DIANANEbased salen ligands, (S,S)-5 and (R,R)-5, promote efficient and highly enantioselective catalytic Nozaki-Hiyama-Kishi reactions. With this novel ligand modification, the addition of various organochromium intermediates to aromatic and aliphatic aldehydes proceeded under catalytic conditions with good levels of stereoinduction (up to 92% ee). Notably, the asymmetric additions of allyl iodide and, in particular, vinyl halides and triflates to aldehydes were achieved with useful levels of enantioselectivity. We are currently exploring the generality and functional-group compatibility of this catalytic carbon-carbon bond-forming process, together with examining its utility with chiral substrates, in the context of projected applications to the stereocontrolled synthesis of more complex structures, such as those that occur in polyketide natural products.

Experimental Section

Typical Procedure: Anhydrous THF (1 mL) was added to CrCl₃ (4.0 mg, 25.0 µmol) and Mn (42 mg, 750 µmol) in a dried Schlenk tube under argon, and the mixture was stirred for 1 h at room temperature. After addition of the ligand (*S*,*S*)-**5** (14.0 mg, 25.0 µmol) and anhydrous NEt₃ (7 µL, 5.0 mg, 50 µmol), the suspension was stirred for another hour at room temperature.^[19] Finally, allyl bromide was added (32 µL, 45.4 mg, 375 µmol). After 1 h, the mixture was cooled to 5 °C, PhCHO (25 µL, 26.2 mg, 250 µmol) and Me₃SiCl (48 µL, 41.2 mg, 380 µmol) were added, and the suspension was

stirred at that temperature. When the aldehyde was completely consumed (as confirmed by thin-layer chromatography (TLC)), saturated aqueous NaHCO₃ was added. After filtration and evaporation, the aqueous phase was extracted with Et₂O. After evaporation of the combined organic phases, the residue was dissolved in THF (1 mL). Aqueous 1M HCl (0.5 mL) was added and the mixture was stirred until desilylation was complete (confirmed by TLC). The solvent was removed, the residue extracted with Et₂O, and the combined organic phases were dried over MgSO₄. Evaporation of the solvent and flash chromatography (hexane/Et₂O 4:1) gave (*R*)-1-phenyl-3-buten-1-ol as a pale-yellow oil (26.7 mg, 180 µmol, 72 % yield, 90 % *ee*).

Received: October 21, 2002 [Z50397]

- For reviews, see: a) A. S. K. Hashmi, J. Prakt. Chem. 1996, 338, 491-495; b) L. A. Wessjohann, G. Scheid, Synthesis 1999, 1—36; c) A. Fürstner, Chem. Rev. 1999, 99, 991-1045; d) A. Fürstner, Chem. Eur. J. 1998, 4, 567-570.
- [2] A. Fürstner, N. Shi, J. Am. Chem. Soc. 1996, 118, 12349-12357.
- [3] a) C. Chen, K. Tagami, Y. Kishi, J. Org. Chem. 1995, 60, 5386–5387; b) Y. Kishi, *Tetrahedron* 2002, 58, 6239–6258; c) H. S. Schrekker, M. W. G. de Bolster, R. V. A. Orru, L. A. Wessjohann, J. Org. Chem. 2002, 67, 1975–1981.
- [4] For asymmetric versions of the NHK reaction using stoichiometric amounts of chiral organochromium agents, see: a) B. Cazes, C. Verniere, J. Goré, *Synth. Commun.* **1983**, *13*, 73–79; b) K. Sugimoto, S. Aoyagi, C. Kibayashi, *J. Org. Chem.* **1997**, *62*, 2322–2323; and ref. [3a].
- [5] a) M. Bandini, P. G. Cozzi, P. Melchiorre, A. Umani-Ronchi, Angew. Chem. 1999, 111, 3558-3561; Angew. Chem. Int. Ed.
 1999, 38, 3357-3359; b) M. Bandini, P. G. Cozzi, A. Umani-Ronchi, Polyhedron 2000, 19, 537-539; c) M. Bandini, P. G. Cozzi, A. Umani-Ronchi, Angew. Chem. 2000, 112, 2417-2420; Angew. Chem. Int. Ed. 2000, 39, 2327-2330; d) M. Bandini, P. G. Cozzi, A. Umani-Ronchi, Tetrahedron 2001, 57, 835-843; e) M. Bandini, P. G. Cozzi, P. Melchiorre, S. Morganti, A. Umani-Ronchi, Org. Lett. 2001, 3, 1153-1155; f) After submission of this manuscript, a report by Kishi et al. on a catalytic enantio-selective process using an oxazoline sulfonamide ligand appeared: H.-W. Choi, K. Nakajima, D. Demeke, F.-A. Kang, H.-S. Jun, Z.-K. Wan, Y. Kishi, Org. Lett. 2002, 4, 4435-4438. However, the enantioselectivities reported are not as high as our best examples.
- [6] For a review of applications of Cr-salen complexes in asymmetric catalysis, see: M. Bandini, P. G. Cozzi, A. Umani-Ronchi, *Chem. Commun.* 2002, 919–927.
- [7] For two examples, see: a) X.-G. Zhou, J.-S. Huang, X.-Q. Yu, Z.-Y. Zhou, C.-M. Che, J. Chem. Soc. Dalton Trans. 2000, 1075–1080; b) D. Seebach, A. K. Beck, A. Heckel, Angew. Chem. 2001, 113, 96–142; Angew. Chem. Int. Ed. 2001, 40, 92–138.
- [8] a) T. Hayashi, Acta Chem. Scand. 1996, 50, 259–266; b) A.
 Weissfloch, R. Azerad, Bioorg. Med. Chem. 1994, 2, 493–500.
- [9] R. T. Hawkins, R. S. Hsu, S. G. Wood, J. Org. Chem. 1978, 43, 4648-4650.
- [10] A full account of the synthesis and configurational assignment of (+)- and (-)-DIANANE (9) will be published elsewhere.
- [11] K. Ishihara, M. Mouri, Q. Gao, T. Maruyama, K. Furuta, H. Yamamoto, J. Am. Chem. Soc. 1993, 115, 11490–11495.
- [12] A. B. Smith III, K. P. Minibiole, P. R. Verhoest, M. Schelhaas, J. Am. Chem. Soc. 2001, 123, 10942–10953.
- [13] V. K. Tandon, A. M. van Leusen, H. Wynberg, J. Org. Chem. 1983, 48, 2767–2769.
- [14] Y. Wu, L. Esser, J. K. De Brabander, Angew. Chem. 2000, 112, 4478-4480; Angew. Chem. Int. Ed. 2000, 39, 4308-4310.

- [15] The synthesis of the aldehyde was performed according to:
 a) M. J. Martinelli, J. Org. Chem. 1990, 55, 5065-5073;
 b) T. Oka, A. Murai, Tetrahedron 1998, 54, 1-20.
- [16] The possibility to recover the valuable ligand by column chromatography after aqueous work-up further adds to the efficiency of our process. Usually, at least 80% of ligand could be recovered in pure form.
- [17] J. K. Stille, J. H. Simpson, J. Am. Chem. Soc. 1987, 109, 2138– 2152.
- [18] K. Takai, K. Sakogawa, Y. Kataoka, K. Oshima, K. Utimoto, Org. Synth. 1995, 72, 180–188.
- [19] For entries 7 and 8, 0.02 equiv of $NiCl_2$ were added prior to the halide.