Cite this: New J. Chem., 2011, 35, 909-914

Silver-free synthesis of nitrate-containing room-temperature ionic liquids†

Nicholas W. Smith, Sai P. Gourisankar,[‡] Jean-Luc Montchamp and Sergei V. Dzyuba*

Received (in Gainesville, FL, USA) 7th November 2010, Accepted 25th January 2011 DOI: 10.1039/c0nj00881h

Nitrate-containing 1-alkyl-3-methylimidazolium ionic liquids, $[C_n-mim]NO_3$, where n = 4, 6, 8, 12, are synthesized in one step by reacting 1-methylimidazole with alkyl nitrates under various conditions without using AgNO₃ and 1-alkyl-3-methylimidazolium halides.

Introduction

Ionic liquids have found numerous applications in various areas of chemistry, physics, biology and engineering.¹ Development of more efficient, facile and greener synthetic approaches to ionic liquids constitutes a significant area of research, which should further increase the utility of ionic liquids.

The majority of ionic liquids feature halogen-containing anions, including $[PF_6]^-$, $[BF_4]^-$, $[CF_3SO_3]^-$ and $[(CF_3SO_2)_2N]^{-}$.² Since some of these anions such as PF_6^- , for example, are prone to hydrolysis, the liberation of HF presents a concern.³ In this light, the synthesis and applications of several halogen-free ionic liquids were reported.⁴

Among room-temperature ionic liquids, nitrate-containing ionic liquids remained a relatively unexplored class of ionic solvents.⁵ Several accounts indicated that nitrate-containing ionic liquids are of interest for potential nuclear waste treatment,⁶ co-solvents and promotors for organic and organometallic reactions⁷ as well as applications in biological⁸ and materials chemistry.⁹

The relatively small number of applications for NO₃-ionic liquids, compared to other types of ionic liquids, is likely due to the fact that the reported syntheses of nitrate-containing ionic liquids rely almost exclusively on the conversion of halide-containing ionic liquids, *i.e.*, chloride or bromide, into the nitrate-containing ionic liquids using AgNO₃. Preparation of NO₃-containing ionic liquids using HNO₃ and an ion-exchange resin was recently reported.¹⁰ Arguably, the significant cost of AgNO₃¹¹ coupled with generation of the AgCl or AgBr by-products complicate large-scale preparations of NO₃-containing ionic liquids using AgNO₃. Thus, the AgNO₃-free synthesis of nitrate-containing ionic liquids using AgNO₃.

important addition to the synthetic repertoire of ionic liquids, potentially enhancing and expanding the applications of these solvents.

We envisioned that a nucleophilic substitution of the ONO₂ group from an alkyl nitrate by an electron pair of the nitrogen of an amine, a reaction that is similar to the synthesis of halide-containing ionic liquids from an amine and alkyl halide,¹² will yield nitrate-containing ionic liquids in one step. This approach would be more economical than the currently available two-step process, *i.e.*, preparation of a halide-containing ionic liquids will also eliminate problems associated with the presence of halides in ionic liquids, which are known to have a drastic impact on the physical and chemical properties of ionic liquids.¹³ To our surprise, a one-step approach to nitrate-containing ionic liquids has not been reported in the literature.

Results and discussion

To test this hypothesis, we used commercially available butyl nitrate and a set of amines, including 1-methylimidazole, pyridine, triethylamine and *N*-methylmorpholine (Scheme 1). This initial reaction was performed under neat conditions at elevated temperatures. Among the tested amines, ionic liquid formation was noted only for 1-methylimidazole and pyridine. In the case of pyridine, however, the conversion was only *ca.* 10%. Neither triethylamine nor *N*-methylmorpholine produced a product: starting materials were recovered in the case of triethylamine,¹⁴ whereas a complex mixture of starting materials and unidentified products was detected in the case of 1-methylmorpholine. Thus, we decided to explore the synthesis of 1-butyl-3-methylimidazolium nitrate, $[C_4-mim]NO_3$, in more detail.

We prepared [C₄-mim]NO₃ ionic liquid using neat conditions (Table 1, entries 1–5). Similarly to halide-containing imidazolium ionic liquids,¹⁵ the efficiency of the synthesis of nitrate-containing ionic liquids was found to be temperature-dependent: the higher the temperature, the higher the conversion. Although temperatures above 120 °C led to even faster

Department of Chemistry, Texas Christian University,

Fort Worth, TX 76129, USA. E-mail: s.dzyuba@tcu.edu;

Fax: +1 817 257 5751; Tel: +1 817 257 6218

[†] Electronic supplementary information (ESI) available: Synthetic procedures for preparation of alkyl nitrates and 1-alkyl-3-methylimidazolium nitrates, additional spectroscopic data. See DOI: 10.1039/ c0nj00881h

[‡] Visiting scholar from Paschal High School, Fort Worth, TX, USA.



Scheme 1 Initial attempts of the one-step synthesis of nitratecontaining ionic liquids.

formation of $[C_4$ -mim]NO₃, the ionic liquid was obtained as a dark brown liquid. It appeared that increasing the amount of the butyl nitrate (Table 1, entries 4–5) produced the ionic liquid in high yield in a short period of time as a dark yellow viscous liquid (Fig. S1, ESI†). It is of interest to point out that

the increased amounts of butyl nitrate up to 2.0 eq. prevented the formation of dark colored impurities. Notably, the excess of butyl nitrate was recovered by decantation from the $[C_4\text{-mim}]NO_3$, which was obtained in high yield.

Consistent with literature accounts,¹⁶ NMR spectroscopy (Fig. S1, ESI[†]) is a convenient way of establishing the purity of ionic liquids suitable for the majority of applications, but it is insensitive to minute amounts of impurities that might be significant for spectroscopic studies. In order to remove the colored impurities (Fig. S1, ESI[†]), the ionic liquid was subjected to several purification protocols according to other reports.¹⁶

We found that passing the solution of the ionic liquid in a molecular solvent (either acetonitrile or ethanol) through a silica gel/charcoal column produced a light brown liquid, albeit in a very low yield.^{16b} Large quantities of eluent were also required to recover the ionic liquid from the column, thus, this procedure was abandoned.

Next, we attempted purification of the $[C_4$ -mim]NO₃ using charcoal/solvent.¹⁶ It appeared that the temperature at which the charcoal treatment was done had a significant impact on the final color of the ionic liquid. Specifically, refluxing the ionic liquid with charcoal in dichloromethane produced a light yellow solution; in refluxing acetonitrile—a yellow solution was obtained; and finally, in refluxing water—a colorless solution was obtained (Fig. S1, ESI†). Notably, repetitive treatment with charcoal appeared to have little impact on

Table 1	Synthesis of [C _n -mim]NO ₃ ionic liquids			R	ionic liquid
				C₄H ₉	[C ₄ -mim]NO ₃
				C ₆ H ₁₃	[C ₆ -mim]NO ₃
		conditions		C ₈ H ₁₇	[C ₈ -mim]NO ₃
				C ₁₂ H ₂₅	[C ₁₂ -mim]NO ₃

Entry	RONO ₂ /eq.	Conditions ^a	Conversion ^{b} (%)	Yield ^{c} (%)
1	$C_4H_0ONO_2(1.1)$	120 °C, 1 h	89	ND^d
2	$C_4 H_9 ONO_2$ (1.1)	120 °C, 3 h	93	ND^d
3	$C_4 H_9 ONO_2 (1.1)$	120 °C, 5 h	98	ND^d
4	$C_4H_9ONO_2$ (2.0)	120 °C, 1 h	93	ND^d
5	$C_4H_9ONO_2$ (2.0)	120 °C, 2 h	>99	90
6	$C_4H_9ONO_2$ (1.1)	MW, 120 °C, 10 min	80	ND^d
7	$C_4H_9ONO_2$ (1.1)	MW, 120 °C, 30 min	85	ND^d
8	$C_4H_9ONO_2$ (2.0)	MW, 120 °C, 10 min	88	ND^d
9	$C_4H_9ONO_2$ (2.0)	MW, 120 °C, 30 min	>99	96
10	$C_4H_9ONO_2$ (2.0)	H ₂ O, 120 °C, 20 h	84	ND^d
11	$C_4H_9ONO_2$ (2.0)	H ₂ O, 120 °C, 40 h	>99	95
12	$C_4H_9ONO_2$ (2.0)	120 °C, 2 h^e	ND^d	99
13	$C_4H_9ONO_2$ (2.0)	H ₂ O, 120 °C, 40 h ^e	ND^d	95
14	$C_6H_{13}ONO_2$ (2.0)	120 °C, 6 h	>99	96
15	$C_6H_{13}ONO_2$ (2.0)	H ₂ O, 120 °C, 20 h	>99	98
16	$C_6H_{13}ONO_2$ (2.0)	MW, 120 °C, 30 min	95	ND^d
17	$C_6H_{13}ONO_2$ (2.0)	MW, 120 °C, 45 min	>99	99
18	$C_8H_{17}ONO_2$ (2.0)	120 °C, 6 h	>99	95
19	$C_8H_{17}ONO_2$ (2.0)	H ₂ O, 120 °C, 20 h	>99	96
20	$C_8H_{17}ONO_2$ (2.0)	MW, 120 °C, 30 min	>99	95
21	$C_{12}H_{25}ONO_2$ (2.0)	120 °C, 6 h	>99	79 ^f
22	$C_{12}H_{25}ONO_2$ (2.0)	MW, 120 °C, 45 min	>99	94 ^f

^{*a*} Unless indicated the reactions were performed on a 12.5 mmol scale (based on 1-methylmidazole) under stirring, exposed to air using the specified set of conditions; H_2O indicates that the reaction was done in the presence of H_2O , MW indicates microwave irradiation. ^{*b*} Determined from ¹H NMR of the crude reaction mixture. ^{*c*} Isolated yields after purification with charcoal/H₂O. ^{*d*} Not determined. ^{*e*} 0.188 mol scale based on 1-methylmidazole. ^{*f*} Isolated yields after washing with Et₂O.

further decolorization of $[C_4$ -mim]NO₃. It should be pointed out that the same color of the $[C_4$ -mim]NO₃ was obtained when this ionic liquid was prepared from a metathesis reaction of the $[C_4$ -mim]Br with AgNO₃. Thus, water/charcoal treatment was chosen as the purification method for all subsequent preparations. Upon removing water under vacuum at 40 °C, the ionic liquid was obtained as a pale yellow viscous liquid. It should also be pointed out, that $[C_4$ -mim]NO₃ can develop a more intense, darker color if exposed to extensive heating.

It has been previously proposed that absorbance and fluorescence spectroscopies could be used to check the purity of ionic liquids.¹⁶ Specifically, transitions above 350 nm in the UV/vis spectrum of an ionic liquid could generally indicate the presence of impurities.^{16c}

The presence of the shoulder at 350 nm indicated that the crude $[C_4$ -mim]NO₃ (Fig. 1) contained such impurities. This transition progressively decreased as the color of the ionic liquid changed from dark yellow to pale yellow (Fig. S1, ESI†). The transition at 307 nm is due to the NO₃-anion: we prepared a 0.7 M solution of KNO₃ in water and obtained the absorbance spectra, which also shows a peak at 307 nm (Fig. S2, ESI†).

Emission spectroscopy has also been used as a tool for determining the purity of ionic liquids. However, unlike absorbance, the use of fluorescence is somewhat controversial as it seems to arise not only due to the impurities,^{16d} but also due to the nature of the ionic liquid.^{16a,b} In the case of [C₄-mim]NO₃, fluorescence spectroscopy appeared to have no correlation with the presence of impurities (Fig. S3, ESI†). The emission of the crude ionic liquid was quite similar to that of the purified ones. Repetitive treatment with charcoal/water did not reduce the emission at *ca.* 430–460 nm. Furthermore, independent preparations of [C₄-mim]NO₃ followed by charcoal treatment exhibited similar emission profiles.

With the purification protocol in hand, we explored the AgNO₃-free synthesis of $[C_n$ -mim]NO₃ ionic liquids in more detail (Table 1). Synthesis of ionic liquids using micro-wave (MW) irradiation is well documented.¹⁷ Not surprisingly, $[C_4$ -mim]NO₃ was efficiently obtained under MW conditions.



As expected, the conversion of 1-methylimidazole increased

We also considered the synthesis of [C4-mim]NO3 in a solvent. Our motivation for using the solvent was primarily to decrease/prevent the formation of colored impurities. After screening several solvents, such as ethylacetate, acetonitrile, ethanol and water, we found that water was the most suitable solvent in terms of reaction efficiency and compatability with our purification procedure. It should be pointed out that this list of solvents is not comprehensive, and other solvents might prove to be more efficient.¹⁸ Considering that the reagent concentration plays significant role in the efficiency of the ionic liquid formation, the amount of water was kept to a minimum (ca. 24% v/v). Under these conditions, a complete conversion of 1-methylimidazole was noted after about 2 days. The purity of the obtained ionic liquid was identical to that obtained under neat conditions. However, although the resulting [C₄-mim]NO₃ had a less intense color, purification with charcoal was still required.

In addition, we carried out scaled-up syntheses of $[C_4$ -mim]NO₃ using the neat and the water-containing conditions (Table 1, entries 12 and 13, respectively). In both cases, after the specified reaction time, the excess of butyl nitrate was decanted and the ionic liquid was subjected to charcoal purification to give [C₄-mim]NO₃ in high yields; the purity of these ionic liquids was identical to that obtained during small scale preparations. The efficiency of the butyl nitrate's recovery depended on the specific conditions. For the reaction done in the presence of water (Table 1, entry 13), we recovered 70% of butyl nitrate. However, under neat reaction conditions (Table 1, entry 12), only 40% of butyl nitrate was recovered. It appeared that butyl nitrate is partially miscible with $[C_4$ -mim]NO₃, which is responsible for an inefficient recovery. Arguably, the presence of water in the reaction mixture (Table 1, entry 13) reduces the miscibility of butyl nitrate and [C₄-mim]NO₃, thus affording more efficient recovery. If more efficient recovery of butyl nitrate is required, the reaction mixture may be washed with Et₂O, prior to subjecting it to the charcoal/water treatment.

We have also tested all of the above conditions, *i.e.*, neat, MW and water, for the synthesis of *N*-butylpyridinium, butyltriethylammonium and *N*-butyl-*N*-methylmorpholinium nitrates. Neither triethylamine nor *N*-methylmorpholine produced the corresponding ionic liquids. Formation of *N*-butylpyridinium nitrate never proceeded above 30-40%yield. Hence, we decided to focus on the synthesis of imidazolium-based nitrate-containing room-temperature ionic liquids. The survey of the literature indicated that only 1-alkyl-3-methylimidazolium nitrates, [C_n-mim]NO₃, where n = 4, 6 and 8 are room-temperature ionic liquids;^{6a,19} other known [C_n-mim]NO₃ ionic liquids, with n = 12, 14, 16, 18 are solids at room temperature.^{9a}



Fig. 1 UV/vis spectra of neat $[C_4$ -mim]NO₃ ionic liquid; 1—crude reaction mixture; 2—1 after charcoal/CH₂Cl₂; 3—1 after charcoal/CH₃CN; 4—1 after charcoal/H₂O (see text for details).



alkyl nitrates are not. Synthesis of alkyl nitrates is well documented, albeit most procedures employ either AgNO₃ or highly corrosive mixture of H₂SO₄ and HNO₃ acids.²⁰ Two procedures reported by Olah et al.²¹ and Gavrila et al.²² had all the features suitable for convenient, facile, scalable and relatively safe syntheses of alkyl nitrates using inexpensive alkyl alcohols as the starting materials (Scheme 2).

BF₃-2H₂O, KNO₃

R-ONO2

Olah et al.²¹ procedure worked well for the synthesis of butyl nitrate, which afforded a pure product in 80% yield without any purification. However, hexyl, decyl and dodecyl nitrates were obtained in significantly lower yields, and were contaminated with a BF₃-complexed alcohol, which in the case of dodecyl alcohol was obtained as the major product (the presence of fluorine was confirmed by ¹⁹F NMR). All attempts to separate the desired nitrates by distillation and extraction proved ineffective. Upon treatment with ethylene diamine followed by acidic work-up, the BF₃-complex was destroyed liberating the corresponding alcohol. Regretfully, separation of the nitrate from the alcohol by distillation or basic extraction also failed. We found that treating the crude mixture of nitrate and alcohol with phenylisocyanate led to the formation of the nitrate and the corresponding carbamate. From this mixture, the alkyl nitrate could be easily isolated by distillation. However, this lengthy procedure might be of little practical use. In order to increase the conversions, we also increased the amounts of KNO₃ and/or BF₃-2H₂O as well as reaction times, yet neither improved the efficiency of the reaction.

It should be pointed out that the presence of the alcohol does not have a measurable effect on the efficiency of the ionic liquid formation, regardless of the conditions, *i.e.*, neat/heat, water/reflux or MW. Considering that the ratio between the alkyl nitrate and the corresponding alcohol could be easily determined from ¹H NMR spectra (the CH₂ resonance at *ca*. 4.4 ppm for the NO₃CH₂R and 3.6 ppm for HOCH₂R), the alcohol-nitrate mixture could, in principle, be directly used for the synthesis of the nitrate-containing ionic liquids. From the ionic liquid, the alkyl alcohol is separated by repetitive washings of the ionic liquid with either ethyl acetate or diethyl ether, and followed by further purification with charcoal in water.

Next, we tested Gavrila et al.22 procedure, which uses a mixture of (CF₃CO)₂O and LiNO₃ as the nitrating reagent. It appeared that by directly adopting this protocol, alkyl nitrates were obtained in moderate yields, albeit some unreacted alcohol remained. In order to achieve a complete conversion

of the starting alcohol, the amounts of both (CF₃CO)₂O and LiNO₃ had to be increased to 2.7 equivalents. Notably, the alkyl trifluoroacetates were obtained as the major by-products. The nitrate and the alkyl trifluoroacetate could not easily be distinguished by ¹H NMR due to resonance overlaps; hence, both ¹³C and ¹⁹F NMR were used for unambiguous identification and purity confirmation. Nonetheless, following this modification, the desired alkyl nitrates were purified by simple distillation from the crude mixture. This procedure was used for the synthesis of the long-chain alkyl nitrates utilized in this work.

Although we have not experienced any safety-related issues during the preparation, purification and handling of alkyl nitrates, care should be exercised when handling alkyl nitrates due to their high flammability.

With the improved procedure in hand, we prepared hexyl, octyl and dodecyl nitrates from the corresponding alcohols. These nitrates were reacted with 1-methylimidazole (Table 1, entries 14-20). The reactions under any given set of conditions appeared to be very efficient and the corresponding nitrate-containing ionic liquids were obtained in high yields. Interestingly, the isolated yields of $[C_n-mim]NO_3$ are comparable to those obtained via a two step procedure that involves AgNO₃. Although, the syntheses of [C₆-mim]NO₃ and [C₈-mim]NO₃ followed similar trends as for [C₄-mim]NO₃ syntheses, several minor deviations were noted: (i) longer reaction times under neat conditions were observed; (ii) a longer reaction time for [C₆-mim]NO₃ formation under MW conditions (Table 1, entries 16 and 17); (iii) the reaction times in water (Table 1, entries 15 and 19) were significantly shorter.

In addition to room-temperature ionic liquid nitrates, we also explored the synthesis of [C12-mim]NO3 as a representative member of solid alkylimidazolium nitrates.^{9a} Although the synthesis of [C12-mim]NO3 appeared to be efficient (Table 1, entries 21 and 22), the purification of this ionic liquid proved to be distinct from other nitrate-containing ionic liquids. Any attempts to utilize charcoal/water treatment failed, due to formation of a soapy composition at the water removal step leading to significant losses of the ionic liquid. Also, dodecylnitrate appeared to be completely miscible with the [C₁₂-mim]NO₃, and no phase separation was noted at the endpoint of the reaction. Therefore, colored impurities and the excess of dodecylnitrate were removed by repetitive washings with Et₂O. The dodecylnitrate was recovered upon removal of Et₂O.

Conclusions

In conclusion, we have developed a facile, AgNO₃-free synthesis of $[C_n-mim]NO_3$ ionic liquids, which does not utilize 1-alkyl-3-methylimidazolium halides as intermediates. Although the reported procedure is limited in scope, the facile nature of this one-step synthesis of [C_n-mim]NO₃ should broaden the range of applications of the nitrate-containing ionic liquids. The green aspect²³ of this one step process should be taken with caution, since the access to many alkyl nitrates is not very efficient at the moment and is currently being explored in our laboratory.

Experimental

Materials and methods

All reagents and solvents were from commercial sources (Sigma-Aldrich, Acros, Alfa Aesar) and were used as received. Butyl nitrate was purchased from Frontier Scientific, Inc. and it was also synthesized according to the reported procedure.²⁴ 1H NMR spectra were recorded on a Varian (300 MHz) spectrometer. The chemical shifts are reported in ppm (δ) downfield from tetramethylsilane in CDCl3 or residual DMSO in DMSO-d6. IR spectra of ionic liquid were acquired on a Midac M series instrument, using NaCl pressed windows either as neat $[C_n-mim]NO_3$ ionic liquid (for n = 4, 6, 8) or deposited from a dichloromethane solution for [C12-mim]NO3. Microwave-based reactions were done using a CEM discovery microwave reactor in manufacturer supplied 10 ml vials. Absorbance measurements were performed on a Agilent 8453 UV-visible instrument with a resolution of 1 nm using 0.1 cm quartz cells for solutions and 0.01 cm cell for neat ionic liquids. Fluorescence measurements were performed using Shimadzu RF-5301PC. Fluorescence measurements were carried out as follows: excitation and emission width slits were 3 mm and 3 mm; intensity-high; using 1 cm quartz cells.

Synthesis of dodecyl nitrate-a representative procedure

 $(CF_3CO)_2O$ (30.0 ml, 0.216 mol) was added to a suspension of LiNO₃ (14.9 g, 0.216 mol) in acetonitrile (400 ml). The mixture was allowed to be stirred at room temperature until LiNO₃ completely dissolved, and the solution was cooled to 0 °C. Na₂CO₃ (22.9 g, 0.216 mol) was added in one portion, followed by the dodecyl alcohol (18.0 ml, 0.080 mol) and the reaction mixture was stirred for 3 h. Subsequently, the reaction mixture was poured into an ice-cold solution of saturated NaHCO₃ (500 ml), and extracted with CH₂Cl₂ (2 × 200 ml). The organic phase was dried (MgSO₄) and volatiles were removed *in vacuo*. The residue was subjected to a vacuum distillation (113–115 b.p.; 1 mmHg) to yield 8.62 g (47% yield) of dodecyl nitrate as a colorless liquid.²⁴

¹H NMR (CDCl₃): δ 4.43 (t, J = 6.9 Hz), 1.71 (p, J = 6.9 Hz, 2H), 1.26 (m, 18H), 0.87 (t, J = 6.9 Hz, 3H); ¹³C NMR (CDCl₃): δ 73.65, 32.14, 29.84 (overlap of 2 peaks; based on signal intensity), 29.73, 29.63, 29.57, 29.34, 26.94, 25.85, 22.91, 14.29.

Hexyl nitrate²¹

Colorless liquid (b.p. 36 °C; 1 mmHg), 30% yield. ¹H NMR (CDCl₃): δ 4.42 (t, J = 6.8 Hz, 2H), 1.70 (p, J = 6.6 Hz, 2H), 1.32 (m, 6H), 0.88 (t, J = 6.6 Hz, 3H); ¹³C NMR (CDCl₃): δ 73.67, 31.47, 26.88, 25.50, 22.62, 14.05.

Octyl nitrate²¹

Colorless liquid (b.p. 86 °C; 4 mmHg), 40% yield. ¹H NMR (CDCl₃): δ 4.43 (t, J = 7.2 Hz, 2H), 1.70 (p, J = 6.6 Hz, 2H), 1.27 (m, 10H), 0.87 (t, J = 6.9 Hz, 3H); ¹³C NMR (CDCl₃): δ 73.72, 31.92, 29.29, 29.27, 26.94, 25.85, 22.83, 14.29.

Synthesis of $[C_4$ -mim $]NO_3$ —a representative procedure under neat conditions

A 25 ml round bottom flask was charged with a stirring bar, 1-methylimidazole (1.0 ml, 12.5 mmol) and butyl nitrate

(2.9 ml, 25.0 mmol) connected to a water condenser and placed into a 120 °C oil bath for 5 h under vigorous stirring. Next, the flask was allowed to cool to room temperature, the excess of butyl nitrate was decanted, and water (24 ml) was added, followed by charcoal (0.73 g). The mixture was refluxed under stirring for 15 min. The reaction mixture was filtered through a double layer of filter paper while hot, the charcoal washed with water (12 ml), and the water was removed *in vacuo* at 40 °C. The residue was redissolved in dichloromethane, filtered to remove any residual charcoal and the volatiles were removed *in vacuo* to give 2.27 g (90% yield) of [C₄-mim]NO₃. ¹H NMR (DMSO-d₆): δ 9.18 (s, 1H), 7.78 (s, 1H), 7.70 (s, 1H), 4.15 (t, J = 7.2 Hz, 2H), 3.83 (s, 3H), 1.74 (p, J = 7.4 Hz, 2H), 1.23 (sept, J = 7.4 Hz, 2H), 0.87 (t, J = 6.9 Hz, 3H); IR (neat): 3150, 2964, 2936, 2875, 1652, 1574, 1560, 1351, 1169 cm^{-1.3}

$[C_6-mim]NO_3^{25}$

¹H NMR (DMSO-d₆): δ 9.14 (s, 1H), 7.77 (t, J = 1.7 Hz, 1H), 7.69 (t, J = 1.7 Hz, 1H), 4.14 (t, J = 7.2 Hz, 2H), 3.83 (s, 3H), 1.75 (m, 2H), 1.23 (m, 6H), 0.84 (t, J = 6.6 Hz, 3H); IR (neat): 3147, 3100, 2961, 2860, 1680, 1652, 1574, 1567, 1560, 1470, 1458, 1349, 1169 cm⁻¹.

[C₈-mim]NO₃²⁵

¹H NMR (DMSO-d₆): δ 9.14 (s, 1H), 7.77 (s, 1H), 7.70 (s, 1H), 4.13 (t, J = 7.2 Hz, 2H), 3.83 (s, 3H), 1.75 (m, 2H), 1.23 (m, 10H), 0.83 (t, J = 6.6 Hz, 3H); IR (neat): 3148, 2958, 2937, 2857, 1680, 1652, 1575, 1560, 1458, 1363, 1168 cm⁻¹.

[C₁₂-mim]NO₃^{9a}

¹H NMR (DMSO-d₆): δ 9.11 (s, 1H), 7.76 (s, 1H), 7.69 (s, 1H), 4.13 (t, J = 7.2 Hz, 2H), 3.83 (s, 3H), 1.75 (m, 2H), 1.23 (m, 18H), 0.87 (t, J = 6.6 Hz, 3H); IR (neat): 2924, 2854, 1682, 1652, 1575, 1560, 1460, 1372, 1169 cm⁻¹.

Synthesis of [C₄-mim]NO₃—a representative procedure under MW irradiation

1-Methylimidazole (1.0 ml, 12.5 mmol), butyl nitrate (2.9 ml, 25.0 mmol) and a stirring bar were charged into a 10 ml vial, sealed and MW irradiated at 120 °C. After 30 min, the vial was allowed to cool to room temperature, the excess of butyl nitrate was decanted, and the residue washed with water (24 ml) into a 100 ml round bottom flask containing a stirring bar. Charcoal (0.73 g) was added and the mixture was refluxed under stirring for 15 min. The reaction mixture was filtered through a double layer of filter paper while hot, the charcoal washed with water (12 ml), and the water was removed *in vacuo* at 40 °C. The residue was redissolved in dichloromethane, filtered to remove any residual charcoal and the volatiles were removed *in vacuo* to give 2.41 g (96% yield) of $[C_4-mim]NO_3$.

Synthesis of $[C_4$ -mim]NO₃—a representative procedure in the presence of water

A 25 ml round bottom flask was charged with a stirring bar, 1-methylimidazole (1.0 ml, 12.5 mmol), butyl nitrate (2.9 ml, 25.0 mmol), water (1.25 ml), connected to a water condenser and placed into a 120 $^{\circ}$ C oil bath for 40 h under vigorous

stirring. Next, the flask was allowed to cool to room temperature, the excess of butyl nitrate was decanted, and water (24 ml) was added, followed by charcoal (0.73 g). The mixture was refluxed under stirring for 15 min. The reaction mixture was filtered through a double layer of filter paper while hot, the charcoal washed with water (12 ml), and the water was removed *in vacuo* at 40 °C. The residue was redissolved in dichloromethane, filtered to remove any residual charcoal and the volatiles were removed *in vacuo* to give 2.4 g (95% yield) of [C₄-mim]NO₃.

Acknowledgements

This work was supported in part by the donors of the ACS-Petroleum Research Fund 47965-G7 (SVD), Andrews Institute of Science Technology and Mathematics (SVD), and the National Science Foundation (CHEM-0953368) (JLM). NWS acknowledges the ACS Division of Medicinal Chemistry for a predoctoral fellowship. SPG thanks the Andrews Institute and TCU-Research Apprentice Program for support. We would like to thank Prof. D. E. Minter (TCU) for access to his facilities.

Notes and references

- For some recent reviews, see: (a) B. Ni and A. D. Headley, *Chem.-Eur. J.*, 2010, 16, 4426–4436; (b) M. D. Green and T. E. Long, *Polym. Rev.*, 2009, 49, 291–314; (c) Z. Samec, J. Langmaier and T. Kakiuchi, *Pure Appl. Chem.*, 2009, 81, 1473–1488; (d) F. Zhou, Y. Liang and W. Liu, *Chem. Soc. Rev.*, 2009, 38, 2590–2599; (e) S. M. Zakeeruddin and M. Graetzel, *Adv. Funct. Mater.*, 2009, 19, 2187–2202; (f) M. Armand, F. Endres, D. R. MacFarlane, H. Ohno and B. Scrosati, *Nat. Mater.*, 2009, 8, 621–629; (g) C. Yao and J. L. Anderson, *J. Chromatogr.*, *A*, 2009, 1216, 1658–1712.
- 2 (a) V. Kempter and B. Kirchner, J. Mol. Struct., 2010, 972, 22–34;
 (b) H. Xue, R. Verma and J. M. Shreeve, J. Fluorine Chem., 2006, 127, 159–176;
 (c) H. Xue and J. M. Shreeve, Eur. J. Inorg. Chem., 2005, 2573–2580;
 (d) T. Kitazume, J. Fluorine Chem., 2000, 105, 265–278.
- (a) M. G. Freire, C. M. S. S. Neves, I. M. Marrucho, J. A. P. Coutinho and A. M. Fernandes, J. Phys. Chem. A, 2010, 114, 3744–3749; (b) J. Ranke, S. Stolte, R. Stoermann, J. Arning and B. Jastorff, Chem. Rev., 2007, 107, 2183–2206; (c) G. S. Fonseca, A. P. Umpierre, P. F. P. Fichtner, S. R. Teixeira and J. Dupont, Chem.-Eur. J., 2003, 9, 3263–3269; (d) R. P. Swatloski, J. D. Holbrey and R. D. Rogers, Green Chem., 2003, 5, 361–363.
- 4 (a) M. Smiglak, N. J. Bridges, M. Dilip and R. D. Rogers, *Chem.-Eur. J.*, 2008, 14, 11314–11319; (b) M. Smiglak, J. D. Holbrey, S. T. Griffin, W. M. Reichert, R. P. Swatloski, A. R. Katritzky, H. Yang, D. Zhang, K. Kirichenko and R. D. Rogers, *Green Chem.*, 2007, 9, 90–98; (c) P. D. Vu, A. J. Boydston and C. W. Bielawski, *Green Chem.*, 2007, 9, 1158–1159; Green Chem. 2008, 10, 1342; (d) E. Kuhlmann, S. Himmler, H. Giebelhaus and P. Wasserscheid, *Green Chem.*, 2007, 9, 233–242; (e) P. Wasserscheid, R. van Hal and A. Bösmann, *Green Chem.*, 2002, 4, 400–404.
- 5 (a) R. Atkin and G. G. Warr, ACS Symp. Ser., 2009, 1030, 317–333; (b) H. R. Mortaheb, M. Mafi, B. Mokhtarani, A. Sharifi, M. Mirzaei, N. Khodapanah and F. Ghaemmaghami, *Chem. Eng. J.*, 2010, 158, 384–392; (c) C. Chiappe, M. Malvaldi, B. Melai, S. Fantini, U. Bardi and S. Caporali, *Green Chem.*, 2010,

12, 77–80; (d) B. Mokhtarani, A. Sharifi, H. R. Mortaheb, M. Mirzaei, M. Mafi and F. Sadeghian, J. Chem. Thermodyn., 2009, 41, 1432–1438.

- 6 (a) M. Fields, G. V. Hutson, K. R. Seddon and C. Gordon, PCT Int. Appl. WO 98061606 A2 19980212, 1998; (b) D. Allen, G. Baston, A. E. Bradley, T. Gorman, A. Haile, I. Hamblett, J. E. Hatter, M. J. F. Healey, B. Hodgson and R. Lewin, et al., Green Chem., 2002, 4, 152–158; (c) A. E. Bradley, C. Hardacre, M. Nieuwenhuyzen, W. R. Pitner, D. Sanders, K. R. Seddon and R. C. Thied, Inorg. Chem., 2004, 43, 2503–2514.
- 7 (a) B. Ballarin, L. Busetto, M. C. Cassani and C. Femoni, *Inorg. Chim. Acta*, 2010, 363, 2055–2064; (b) S. Rostamizadeh, R. Aryan, H. R. Ghaieni and A. M. Amani, *J. Heterocycl. Chem.*, 2009, 46, 74–78; (c) C. Chiappe, E. Leandri and M. Tebano, *Green Chem.*, 2006, 8, 742–745.
- 8 (a) R. Vijayaraghavan, A. Izgorodin, V. Ganesh, M. Surianarayanan and D. R. MacFarlane, *Angew. Chem., Int. Ed.*, 2010, **49**, 1631–1633; (b) N. Byrne and C. A. Angell, *Molecules*, 2010, **15**, 793–803; (c) N. Byrne and C. A. Angell, *Chem. Commun.*, 2009, 1046–1048.
- 9 (a) E. Guillet, D. Imbert, R. Scpelliti and J.-C. G. Bünzli, *Chem. Mater.*, 2004, 16, 4063–4070; (b) O. A. Mazyar, G. K. Jennings and C. McCabe, *Langmuir*, 2009, 25, 5103–10; (c) J.-W. Lee and Y.-T. Yoo, *Sens. Actuators, B*, 2009, B137, 539–546; (d) N. Cai, J. Zhang, D. Zhou, Z. Yi, J. Guo and P. Wang, *J. Phys. Chem. C*, 2009, 113, 4215–4221.
- 10 I. Dinares, C. Garcia de Miguel, A. Ibanez, N. Mesquida and E. Alcalde, *Green Chem.*, 2009, 11, 1507–1510.
- 11 Cost of AgNO₃ (ACS reagent, >99%) from Sigma-Aldrich: #209139-500G \$701.70.
- 12 (a) S. V. Dzyuba, K. D. Kollar and S. S. Sabnis, J. Chem. Educ., 2009, 86, 856–858; (b) A. Stark, D. Ott, D. Kralisch, G. Kreisel and B. Ondruscka, J. Chem. Educ., 2010, 87, 196–201.
- 13 K. R. Seddon, A. Stark and M.-J. Torres, Pure Appl. Chem., 2000, 72, 2275–2287.
- 14 Under the identical conditions reacting Et_3N with BuBr failed to produce $BuEt_3N^+Br^-$ and unchanged starting materials were recovered.
- 15 S. V. Dzyuba and R. A. Bartsch, J. Heterocycl. Chem., 2001, 38, 265–268.
- (a) P. Nockemann, K. Binnemans and K. Driesen, Chem. Phys. Lett., 2005, 415, 131–136; (b) A. Paul, P. K. Mandal and A. Samanta, J. Phys. Chem. B, 2005, 109, 9148–9158; (c) M. J. Earle, C. M. Gordon, N. V. Plechkova, K. R. Seddon and T. Welton, Anal. Chem., 2007, 79, 758–764; (d) A. K. Burrell, R. W. Del Sesto, S. N. Baker, T. M. McCleskey and G. A. Baker, Green Chem., 2007, 9, 449–454; Green Chem. 2007, 9, 809.
- (a) V. V. Namboodiri and R. S. Varma, *Tetrahedron Lett.*, 2002,
 43, 5381–5383; (b) R. S. Varma and V. V. Namboodiri, *Chem. Commun.*, 2001, 643–644.
- 18 J. C. Schleicher and A. M. Scurto, *Green Chem.*, 2009, **11**, 694–703. 19 (a) A. L. Carmichael and K. R. Seddon, *L. Phys. Org. Chem.*, 2000.
- (a) A. J. Carmichael and K. R. Seddon, J. Phys. Org. Chem., 2000,
 13, 591–595; (b) L. Cammarata, S. G. Kazarian, P. A. Salter and T. Welton, Phys. Chem. Chem. Phys., 2001, 3, 5192–5200.
- 20 (a) G. Heigle, J. Nhuyen and Y. Zhou, *Synth. Commun.*, 2004, 34, 2507–2511; (b) L. Lazzarato, B. Rolando, M. L. Lolli, G. C. Tron, R. Fruttero, A. Casco, G. Deleide and H. L. Guether, *J. Med. Chem.*, 2005, 48, 1322–1329.
- 21 G. A. Olah, Q. Wang, X.-y. Li and G. K. S. Prakash, *Synthesis*, 1993, 207–208.
- 22 A. Gavrila, L. Andersen and T. Skrydstrup, *Tetrahedron Lett.*, 2005, 46, 6205–6207.
- 23 (a) R. A. Sheldon, Green Chem., 2007, 9, 1273–1283;
 (b) M. Deetlefs and K. R. Seddon, Green Chem., 2010, 12, 17–30.
- 24 F. L. M. Pattison and G. M. Brown, Can. J. Chem., 1956, 34, 879–884.
- 25 M. Fields, G. V. Hutson, K. R. Seddon and C. Gordon, PCT Int. Appl. WO 98061606 A2 19980212, 1998.