

Published on Web 03/06/2007

Bis(phenyl)dirhodium(III) Caprolactamate: A Dinuclear Paddlewheel Complex with No Metal–Metal Bond

Jason M. Nichols,[‡] Joffrey Wolf,[§] Peter Zavalij,[‡] Bindhu Varughese,[‡] and Michael P. Doyle^{*,‡}

Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742, and Laboratoire de Chimie de Coordination, 205 route de Narbonne, 31077 Toulouse Cedex 4, France

Received January 7, 2007; E-mail: mdoyle3@umd.edu

The preparation and electronic structure of dinuclear metal complexes with a metal-metal bond have been long-standing areas of interest.1 Recent work has provided synthetic access to and/or elucidated the electronic structure of M_2^{6+} complexes (M = transition metal).² Despite the body of work, structurally welldefined Rh2⁶⁺ complexes have remained an elusive target.¹ This manuscript reports the first unambiguous and fully characterized dirhodium(III) paddle-wheel complex, bis(phenyl)dirhodium(III) caprolactamate (1), and its synthesis by a novel oxidation of dirhodium(II) caprolactamate (2) (Scheme 1).³ Isolating 1 allows a comparative analysis of the Rh_2^{n+} complexes (n = 4, 5, and 6) of dirhodium caprolactamate. This analysis demonstrates that 1 is a dinuclear paddlewheel complex without a formal metal-metal bond because of a ligand induced change in its electronic structure.⁴ The formation of 1 under such mild oxidation conditions implies that similar species could play a role in the redox chemistry of dirhodium complexes.5

The aerobic oxidation of 2 with Cu(I)OTf (10 mol %) in the presence of NaBPh₄ (5 equiv) provided 1 in 77% yield after purification (Scheme 1b). The structure of **1** was first indicated by signals corresponding to two equivalent phenyl ligands in addition to the caprolactamate ligands observed in both the ¹H and ¹³C NMR spectra. Aryl ring vibrations at 1583 and 1550 cm⁻¹, and the parent ion corresponding to 1 (M+H) by high-resolution mass spectroscopy (ESI) provided further evidence. The visible spectrum of 1 with λ_{max} (ϵ M⁻¹ cm⁻¹) at 430 (4540) nm was consistent with previous reports of electrochemically generated Rh2⁶⁺ complexes that were not isolated.⁶ X-ray diffraction of a single-crystal grown in CH₂Cl₂ confirmed that 1 was indeed [Rh₂(cap)₄](C₆H₅)₂ (Figure 1). To complete the Rh_2^{n+} series, the Rh_2^{5+} complex, $[Rh_2(cap)_4]$ -(OTf)·2H₂O (3), was prepared by treating 2 with Cu(II)[OTf]₂ in wet ethyl acetate (Scheme 1a).⁷ The visible spectrum of 3 in CH₂Cl₂ was typical of Rh₂⁵⁺ structures with λ_{max} (ϵ M⁻¹ cm⁻¹) at 505 (4502) and 970 (1086) nm.⁶ As an odd electron species, 3 exhibited severe line broadening in the ¹H NMR, but its structure was confirmed crystallographically to have the typical dirhodium core with axial aquo ligands and an outersphere triflate ion.8

In terms of the molecular geometry, **1** does not deviate from the general dirhodium(II) carboxamidate structure. The coordination sphere of **1**, however, is completed by two axial C₆H₅ anionic ligands with notable structural distortions. The Rh–Rh–C angles (155.4°, 156.2°) are significantly distorted from linearity toward the Rh–O bonds, and the Rh–Rh bond of **1** (2.519 Å) is lengthened compared to **2** (2.422 Å). Considering the general $\sigma^2 \pi^4 \delta^2 \pi^{*4} \sigma^{*n}$ electronic structure of dinuclear paddlewheel complexes,⁹ a shorter Rh–Rh bond is expected with increasing bond order upon oxidation from a Rh₂⁴⁺ (n = 2) to a Rh₂⁶⁺ complex (n = 0).¹⁰ Consistent with this electronic structure, the Rh–Rh bond length shortens









Figure 1. ORTEP drawing of one of two crystallographically independent molecules of 1, $C_{36}H_{50}N_4O_4Rh_2$, $R_1 = 3.00\%$. Selected bondlengths (Å) and angles (deg): Rh2–Rh2' (2.519); Rh2–C60 (2.000), Rh2'–O40' (2.085); Rh2'–O50 (2.083); Rh2–N40 (2.011); Rh2–N50 (2.014); Rh2–Rh2'–C60 (155.4); Rh2'–Rh2–N40 (96.4); Rh2'–Rh2–N50 (94.0); Rh2–Rh2'–O40 (77.4); Rh2–Rh2'–O50 (79.8). Ellipsoids are shown at a 30% probability level. Solvent and hydrogens are omitted for clarity.

modestly as oxidation occurs from **2** (n = 2) to **3** (n = 1), 2.422 to 2.384 Å, respectively. However, the Rh–Rh bond length in **1** (n = 0) is approximately 0.1 Å *longer* than the corresponding bond length in **2**. The lengthening of the bond is an indication of a change in the electronic structure of the dirhodium core.^{6,11}

The formation of two new Rh–C bonds can be expected to change the hybridization of the Rh-centered orbitals from 2 (dsp²) to 1 (d²sp³).¹² An alternative electronic configuration of 1 would then be $\pi^4 \delta^2 \pi^{*4} \delta^{*2}$ for a d²sp³ dimer resulting in the formal cleavage of the Rh–Rh bond.¹³ X-ray photoelectron spectroscopy (XPS) of compounds 1–3 indicates that a fundamental change in the metal-centered orbitals indeed occurs when 2 is oxidized to 1. The electron binding energy (*E*_b) for the rhodium 3d-orbitals (Rh 3d_{5/2}) increased from 308.08 to 309.09 eV upon oxidation from 2 to 3. The ~1 eV increase in *E*_b from Rh₂⁴⁺ to Rh₂⁵⁺ is consistent with other dirhodium(II) complexes.¹⁴ However, upon oxidation from Rh₂⁵⁺ to Rh₂⁶⁺ (**3** to 1), *E*_b(Rh 3d_{5/2}) remained virtually unchanged

[‡] University of Maryland. [§] Laboratoire de Chimie de Coordination.





at 309.11 eV. The binding energy of 1 suggests that the complex does not retain the $\sigma^2 \pi^4 \delta^2 \pi^{*4} \delta^{*n}$ electronic structure with n = 0. However, $E_{\rm b}({\rm Rh} \ 3d_{5/2})$ is consistent with monomeric d²sp³ hybridized Rh(III) complexes and a $\pi^4 \delta^2 \pi^{*4} \delta^{*2}$ electron configuration.¹⁵ The presence of ¹H and ¹³C NMR signals, the lack of a measurable magnetic susceptibility for 1, and the disappearance of the $\delta - \delta^*$ transition at 970 nm in the visible spectrum of ${\bf 1}$ indicate that the δ^* orbital is occupied and **1** is diamagnetic, consistent with the configuration assignment.

In addition to the structural features of **1**, its formation from **2** was surprising. How does the oxidation of 2 occur in the presence of Cu(II) and NaBPh₄? The instability of NaBPh₄ toward metal cations has been well documented, including aryl transfer processes,16 but the oxidation of a metal center by a tetraarylborate salt is unlikely.¹⁷ The more likely oxidant is a Cu(II) salt catalytically regenerated in the presence of oxygen. The preparation of 3 from 2 with Cu(II)[OTf]₂ demonstrated clearly that the first oxidation was possible. Submitting 3 to the original reaction conditions gave 1 in 73% yield after purification (Scheme 1c). This indicated that the key transformation occurs from the Rh25+ redox state.

A series of experiments helped determine the role of copper salts and NaBPh₄ in the reaction. Complex **3** was diluted in ethyl acetate, washed repeatedly with water, and concentrated to dryness to minimize the amount of residual copper salts from its preparation. The oxidation of 3 to 1 was observed over 60 min via spectral changes at the λ_{max} of **1** (430 nm). Under the reaction conditions, there was no reaction between 3 and NaBPh₄ (5 equiv) or 3 and excess Cu(II)[OTf]₂. Very little conversion of **3** to **1** occurred under nitrogen while the aerobic reaction in the presence of a catalytic amount of Cu(I)OTf (10 mol %) rapidly generated 1 in almost complete conversion. A simple model can be proposed for the oxidation of 2 to 1 on the basis of these data (Scheme 2). A Cu(II)/Cu(I) couple with oxygen as a terminal oxidant oxidizes 2 to 3. An aryl-transfer from NaBPh₄ to 3 gives a transient 3-Ph species whose oxidation potential is low enough¹⁸ for Cu(II)/Cu(I) couple to oxidize it to 3-Ph⁺.¹⁹ Phenyl transfer from an additional equivalent of NaBPh₄ completes the process and generates **1**.

The isolation of **1** provides the first unequivocal example of a dirhodium(III) complex. A copper catalyzed aerobic oxidation with NaBPh₄ as a phenyl transfer agent provides entry into this novel Rh₂⁶⁺ species. Structural and spectroscopic data indicate a cleavage of the Rh-Rh bond resulting from a change in the electronic structure of the complex. Preliminary data suggest that this synthetic approach can be employed with other dirhodium(II) carboxamidates, and work is underway to explore the structural scope and applications of dirhodium(III) complexes.

Acknowledgment. We are grateful to the National Science Foundation for their generous support. We also thank the Fond Social Européen for funding Joffrey Wolf and Bristol-Meyers-Squibb for funding Jason Nichols with an ACS Organic Division Fellowship.

Supporting Information Available: Experimental procedures, CIF files, XPS data, and characterizations. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (1) Cotton, F. A., Murillo, C. A., Walton, R. A., Eds. Multiple Bonds Between
- Octton, F. A., Murillo, C. A., Walton, R. A., Eds. Multiple Bonds Between Metal Atoms, 3rd ed.; Springer Science: New York, 2005.
 (a) Shi, Y.-H.; Chen, W.-Z.; John, K. D.; Da Re, R. E.; Cohn, J. L.; Xu, G.-L.; Eglin, J. L.; Sattelberger, A. P.; Hare, C. R.; Ren, T. Inorg. Chem. 2005, 44, 5719. (b) Xu, G. L.; Crutchley, R. J.; DeRosa, M. C.; Pan, Q. J.; Zhang, H. X.; Wang, X.; Ren, T. J. Am. Chem. Soc. 2005, 127, 13354.
 (c) Barral, M. C.; Gallo, T.; Herrero, S.; Jimenez-Aparicio, R.; Torres, M. R.; Urbanos, F. A. Inorg. Chem. 2006, 45, 3639. (d) Cotton, F. A.; Murillo, C. A.; Villagran, D.; Yu, R. J. Am. Chem. Soc. 2006, 128, 3281.
 (e) Mendiratta, A.; Cummins, C. C.; Cotton, F. A.; Ibragimov, S. A.; Murillo, C. A.; Villagran, D. Inorg. Chem. 2006, 45, 3438.
 (b) A referee has pointed out that the bent dinitrosyl complexes of dirhodium
- (3) A referee has pointed out that the bent dinitrosyl complexes of dirhodium tetracarboxylates could be formally dirhodium(III) complexes with longer Rh-Rh bondlengths than the parent compounds, as with 1, but unambiguous electronic assignment was not made and these complexes are nitric oxide rather than nitroxyl donors. Hilderbrand, S. A.; Lim, M. H.; Lippard, S. J. J. Am. Chem. Soc. 2004, 126, 4972.
- (4) (a) In contrast to the many $d^8 d^8$ dimers that lack a metal bond with the (4) (a) In contrast to the many d^{--d⁻} d⁻ dimers that lack a metal bond with the standard electronic configuration. Cotton, F. A.; Gu, J.; Murillo, C. A.; Timmons, D. J. J. Am. Chem. Soc. **1998**, *120*, 13280. (b) Berry, J. F.; Bothe, E.; Cotton, F. A.; Ibragimov, S. A.; Murillo, C. A.; Villagran, D.; Wang, X. Inorg. Chem. **2006**, *45*, 4396.
 (5) (a) Catino, A. J.; Forslund, R. E.; Doyle, M. P. J. Am. Chem. Soc. **2004**, *126*, 13622. (b) Espino, C. G.; Fiori, K. W.; Kim, M.; DuBois, J. J. Am. Chem. Soc. **2004**, *126*, 156278. (a) Chiefordide, H. T.; DuBors, K. P. Ace.
- Chem. Soc. 2004, 126, 15378. (c) Chifotides, H. T.; Dunbar, K. R. Acc. Chem. Res. 2005, 38, 146. (d) Catino, A. J.; Nichols, J. M.; Nettles, B. J.; Doyle, M. P. J. Am. Chem. Soc. 2006, 128, 5648.
 (6) Kadish, K. M.; Phan, T. D.; Giribabu, L.; Van Caemelbecke, E.; Bear, J.
- . Inorg. Chem. 2003, 42, 8663.
- (7) The redox potential of the Cu^{II} - Cu^{I} couple (1.2 eV vs SCE in CH₃CN) suggested its ability to oxidize 2. Sumalekshmy, S.; Gopidas, K. R. Chem. Phys. Lett. 2005, 413, 294.
- (8) See Supporting Information for the XRD data of 2 and 3.
- (9) Berry, J. F.; Cotton, F. A.; Ibragimov, S. A.; Murillo, C. A.; Wang, X. Inorg. Chem. 2005, 44, 6129.
- (10) Chifotides, H. T.; Dunbar, K. R. In Multiple Bonds between Metal Atoms, 3rd ed.; Springer Scientific: New York, 2005; p 465.
- (11) Axial ligand donor strength has been shown to have little effect on the Rh-Rh bond length for complexes with standard electronic configurations. Cotton, F. A.; Hillard, E. Â.; Murillo, C. A. J. Am. Chem. Soc. 2002, 124, 5658.
- (12) An analogous phenomenon was observed for Ru26+ complexes with a similar lengthening of the metal bond. Bear, J., L.; Chen, W. Z.; Han, B.; Huang, S.; Wang, L. L.; Thuriere, A.; Van Caemelbecke, E.; Kadish Karl, M.; Ren, T. *Inorg. Chem.* **2003**, *42*, 6230.
- (13) The full extent of Rh-Rh bond cleavage is unclear at this time as the Rh atoms are still within the covalent radius for Rh (136 pm) by 10 pm. Pauling, L. Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem. 1978, 34, 746.
- (14) Dennis, A. M.; Howard, R. A.; Kadish, K. M.; Bear, J. L.; Brace, J.; Winograd, N. Inorg. Chim. Acta **1980**, 44, L139. (15) Gassman, P. G.; Macomber, D. W.; Willging, S. M. J. Am. Chem. Soc.
- 1985, 107, 2380
- (16) (a) Strauss, S. H. Chem. Rev. 1993, 93, 927. (b) Fagnou, K.; Lautens, M. Chem. Rev. 2003, 103, 169.
- (17) In most cases NaBAr4 is a reductant. Murphy, S.; Schuster, G. B. J. Phys. Chem. 1995, 99, 511.
- Axial ligands can dramatically effect oxidation potentials. Bear, J. L.; Han, B.; Huang, S.; Kadish, K. M. *Inorg. Chem.* **1996**, *35*, 3012. (18)
- The 5⁺/6⁺ redox couple for Rh₂(cap)₄ is estimated to be \sim 1300 mV versus SCE in CH3CN. Doyle, M. P.; Ren, T. In Progress in Inorganic Chemistry; Karlin, K. Ed.; Wiley: New York, 2001; Vol. 49, p 113.

JA070122S