

N–H Insertion reactions of rhodium carbenoids. Part 2.¹ Preparation of *N*-substituted amino(phosphoryl)acetates (*N*-substituted phosphorylglycine esters)²


Leigh Ferris,^a David Haigh^b and Christopher J. Moody^{*†,a}

^a Department of Chemistry, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK

^b SmithKline Beecham Pharmaceuticals, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ, UK


Rhodium(II) acetate-catalysed reaction of ethyl 2-diazo-2-diethoxyphosphorylacetate **2** with carbamates, amides, ureas or anilines gives a range of *N*-substituted 2-amino-2-diethoxyphosphorylacetates **3–18** by N–H insertion reaction of the intermediate rhodium carbenoid.

N-Acylaminophosphonates (phosphorylglycines) **1** are useful



intermediates in synthesis. Originally prepared and used by the Merck group in the synthesis of cephalothin,^{3,4} they have subsequently been employed in the preparation of other cephalosporins and analogues.^{5–9} However, it is in the preparation of dehydro amino acids by the Wadsworth–Emmons reaction that phosphorylglycine derivatives have found the widest application.^{10–28}

Phosphorylglycines were originally prepared by methoxycarbonylation of the anion of the Schiff base of diethyl (aminomethyl)phosphonate,³ but are more commonly obtained by Michaelis–Arbusov reaction of trialkyl phosphites with α -halo- or α -alkoxy-glycine esters^{8,10,11,29–33} or with aziridines.³⁴ Other preparative methods include the electrophilic amination of phosphorylacetates,^{35,36} the oximation of phosphorylacetates followed by reduction,^{37,38} the addition of diethyl phosphite anion to the Schiff base of ethyl glyoxalate³⁹ and the Me₃SiBr-promoted conversion of phosphoranones into phosphonates.²⁷ These methods are summarised in Scheme 1.



† Present address: Department of Chemistry, University of Exeter, Stocker Road, Exeter EX4 4QD, UK

Two slightly conflicting reports also describe the preparation of amino(phosphoryl)acetates from phosphorylacetate by reaction with a sulfonyl azide followed by reduction. Thus ethyl diethoxyphosphorylacetate is reported to react with trifluoromethanesulfonyl azide to give ethyl 2-azido-2-diethoxyphosphorylacetate, EtO₂CCHN₃PO(OEt)₂, hydrogenation (Pd–C) of which gave the corresponding α -amino compound.³⁹ On the other hand, the closely related reaction of *tert*-butyl diethoxyphosphorylacetate, Bu^tO₂CCH₂PO(OEt)₂, with toluene-*p*-sulfonyl azide was reported to give the 2-diazo-2-diethoxyphosphorylacetate;³⁸ again hydrogenation over Pd–C gave the 2-amino-2-diethoxyphosphorylacetate.

In view of the continuing interest in phosphorylglycines and in aminophosphonates in general,⁴⁰ we decided to investigate the N–H insertion reactions of rhodium carbenoids derived from readily available ethyl 2-diazo-2-diethoxyphosphorylacetate as a simple route to a range of *N*-acyl- and *N*-aryl-amino(phosphoryl)acetates. We now report our results in detail.²

Results and discussion

Insertion reactions of metallocarbenoids are widely used in synthesis, and a comprehensive discussion of the N–H insertion reaction is included in the preceding paper.¹ However, very few of these reactions involve diazophosphonates, and therefore a study of the N–H insertion reactions of ethyl 2-diazo-2-diethoxyphosphorylacetate **2** was initiated. Compound **2** was prepared from ethyl diethoxyphosphorylacetate in 60–63% yield by diazo transfer reaction using toluene-*p*-sulfonyl azide.^{41,42} Although both the diazo transfer reagent and the diazo compound are potentially explosive (**CAUTION**), with due regard for the hazards, we carried out the preparation on the 0.2 mol scale without incident. As an alternative to toluene-*p*-sulfonyl azide, the commercially available diazo transfer reagent, azidotris(diethylamino)phosphonium bromide,⁴³ can be used. This gives an improved yield of 93% (on a 8 mmol scale).⁴⁴

The rhodium(II) acetate-catalysed decomposition of **2** was investigated in the presence of a wide range of H–H compounds (Scheme 2). The reactions were carried out in boiling




Table 1 Synthesis of compounds 3–18


Entry	R ¹	R ²	Product	Yield (%)
1	Bu ^t O ₂ C	H	3	75
2	BnO ₂ C	H	4	78
3	Ac	H	5	79
4	EtCO	H	6	56
5	Bu ^t CO	H	7	76
6	MeNHCO	H	8	40
7	Pr	Bu ^t O ₂ C	9	46
8	Ph	H	10	72
9	4-MeC ₆ H ₄	H	11	73
10	4-MeOC ₆ H ₄	H	12	70
11	4-NO ₂ C ₆ H ₄	H	13	81
12	4-ClC ₆ H ₄	H	14	71
13	2-MeOC ₆ H ₄	H	15	74
14	2-BrC ₆ H ₄	H	16	70
15	2,6-Me ₂ C ₆ H ₃	H	17	71
16	2-Cl-4-CF ₃ C ₆ H ₃	H	18	61

toluene; diazophosphonates are generally more stable than simple diazo esters,⁴⁵ and no reaction occurred at room temperature. Likewise, there was no reaction at the elevated temperature in the absence of the rhodium catalyst.

Reaction with *tert*-butyl or benzyl carbamate gave the corresponding N–H insertion products 3 and 4 in good yield (Table 1). In the case of the *Z* protected phosphorylglycine, the reaction was carried out on a 2 g scale without reduction in yield. Other *N*-acyl compounds also react: simple amides such as acetamide, propionamide and isovaleramide gave the corresponding *N*-acylamino phosphorylacetates 5–7, and *N*-methylurea gave the carbamoylamino derivative 8 in modest yield (Table 1), with no product resulting from insertion into the MeN–H bond being observed. Attempts to effect N–H insertion reactions on phthalimide, pyrrolidin-2-one or azetidin-2-one using 2 were unsuccessful. The lactams pyrrolidin-2-one and piperidin-2-one are known to undergo N–H insertion reactions with the carbenoid-derived by copper-catalysed decomposition of methyl diazoacetate, although the yields are poor (2 and 18% respectively).⁴⁶ Likewise, there are reports of intermolecular insertions into the N–H bond of β -lactams (azetidin-2-ones) in the rhodium(II)-catalysed reactions of more reactive diazo esters;^{47–50} the intramolecular insertion into β -lactam N–H bonds is much better known and is widely used as a route to bicyclic β -lactams.⁴⁰

Rhodium(II) acetate-catalysed decomposition of 2 in the presence of simple alkylamines did not result in the formation of the corresponding N–H insertion product, presumably as a result of catalyst poisoning. However, the overall product of insertion into the N–H bond of a primary alkylamine can readily be obtained by use of the corresponding *N*-Boc derivative, the *N*-*tert*-butoxycarbonyl group being removed during the reaction (Table 1, Entry 7).

Anilines, on the other hand, react readily to give the corresponding *N*-aryl- α -phosphorylglycines in good yield (Table 1, Entries 8–16). *N*-Aryl- α -phosphorylglycines are a poorly described group of compounds,⁵¹ and hitherto, no general method of preparation has been developed. A range of anilines was used, and the reaction is apparently insensitive to the basicity of the aniline NH₂ group, 4-nitroaniline reacting just as readily as 4-methoxyaniline. In the case of 2-chloro-4-trifluoromethylaniline, the N–H insertion product 18 was accompanied by the amide 19, formed by reaction of the initial carboxylate with the excess aniline.

19 Ar = 2-Cl-4-CF₃C₆H₃

In summary, the rhodium(II)-catalysed N–H insertion reaction of compound 2 represents a simple route (two steps from ethyl diethoxyphosphorylacetate) to a wide range of *N*-substituted amino(phosphoryl)acetates.

Experimental

For general experimental points, see the preceding paper.

General procedure for N–H insertion reactions

A stirred solution of ethyl 2-diazo-2-diethoxyphosphorylacetate 2 (250 mg, 1 mmol) and the NH compound (5 mmol) in dry toluene (5 ml) was treated with rhodium(II) acetate (9 mg, 2 mol%). The mixture was heated at reflux overnight, the solvent evaporated and the residue chromatographed on silica gel (light petroleum–ether) to give the product.

Ethyl 2-*tert*-butoxycarbonylamino-2-diethoxyphosphorylacetate 3. Colourless crystalline solid (75%), mp 61–62 °C (from ether–light petroleum) (lit.,¹⁰ 64.5–65.5 °C) (Found: C, 46.0; H, 7.9; N, 4.2. Calc. for C₁₃H₂₆NO₇P, C, 46.0; H, 7.7; N, 4.1%); $\nu_{\max}(\text{CHCl}_3)/\text{cm}^{-1}$ 1754, 1732, 1250, 1140 and 1030; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 5.35(1 H, br d, exch. CF₃CO₂D, NH), 4.80 [1 H, dd, *J* 9.07 (exch. CF₃CO₂D) and 22.55, CHP], 4.34–4.15 (6 H, m, OCH₂), 1.45 (9 H, s, Bu^t) and 1.35–1.29 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 167.2, 154.9, 80.7, 63.7, 62.3, 52.2, 28.2, 16.3 (d, *J* 6.7) and 14.1; $\delta_{\text{P}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 15.29; *m/z* (FAB) 362 (MNa⁺, 100%) and 340 (MH⁺, 12).

Ethyl 2-benzyloxycarbonylamino-2-diethoxyphosphorylacetate 4. Colourless crystalline solid (78%), mp 50–51 °C (from ether–light petroleum) (lit.,¹⁰ 47–48 °C) (Found: C, 51.5; H, 6.5; N, 3.75. Calc. for C₁₆H₂₄NO₇P, C, 51.55; H, 6.6; N, 3.5%); $\nu_{\max}(\text{CHCl}_3)/\text{cm}^{-1}$ 2940, 1710, 1250 and 1120; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 7.37–7.35 (5 H, m, ArH), 5.67 (1 H, br s, NH), 5.25 (2 H, d, *J* 3.4, PhCH₂), 4.97, [1 H, dd, *J* 9.07 (exch. CF₃CO₂D) and 22.27, CHP], 4.28–4.21 (6 H, m, OCH₂) and 1.35–1.29 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 166.9, 155.6, 135.9, 128.9, 128.7, 128.6, 67.5, 63.8, 63.7, 62.5, 52.7 (d, *J* 146.4), 16.3 (d, *J* 6.8) and 14.1; $\delta_{\text{P}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 14.74; *m/z* (FAB) 374 (M + NH₄⁺, 59%), 192(43) and 91(100).

Ethyl 2-acetylamino-2-diethoxyphosphorylacetate 5. Colourless crystalline solid (79%), mp 88–89 °C (from ether–light petroleum) (lit.,³⁰ 84–85 °C) (Found: C, 42.8; H, 7.4; N, 5.0. Calc. for C₁₀H₂₀NO₆P, C, 42.7; H, 7.2; N, 5.0%); $\nu_{\max}(\text{CHCl}_3)/\text{cm}^{-1}$ 1720, 1680, 1250 and 1100; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 6.27 (1 H, br s, NH, exch. D₂O), 5.17 (1 H, dd, *J* 8.8 and 21.9, CHP), 4.31–4.10 (6 H, m, OCH₂), 2.08 (3 H, s, MeCO) and 1.33 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 169.6, 166.9, 63.7, 63.8, 62.4, 50.8 (d, *J* 146.6), 22.9, 16.3 (d, *J* 6.8) and 14.1; $\delta_{\text{P}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 14.96; *m/z* (CI) 299 (M + NH₄⁺, 100%) and 282 (MH⁺, 67).

Ethyl 2-diethoxyphosphoryl-2-propionylaminoacetate 6. Colourless crystalline solid (56%), mp 39–40 °C (Found: M⁺, 295.1188. C₁₁H₂₂NO₆P requires M, 295.1185); $\nu_{\max}(\text{CHCl}_3)/\text{cm}^{-1}$ 1743, 1633 and 1250; $\delta_{\text{H}}(250 \text{ MHz}; \text{CDCl}_3)$ 6.30 (1 H, br d, *J* 8, NH, exch. D₂O), 5.17 (1 H, dd, *J* 8.8 and 22.1, CHP), 4.18 (6 H, m, OCH₂), 2.30 (2 H, q, *J* 7.5, CH₂CONH), 1.33 (9 H, m, Me) and 1.18 (3 H, t, *J* 7.5, Me); $\delta_{\text{C}}(62.9 \text{ MHz}; \text{CDCl}_3)$ 173.6, 167.3, 64.1, 62.8, 51.0 (d, *J* 145.4), 29.7, 16.6 (d, *J* 6.8), 14.4 and 9.7; $\delta_{\text{P}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 15.28; *m/z* (EI) 295 (M⁺, 5%), 250 (10), 222 (11), 166 (100), 138 (80), 102 (40), 82 (20), 65 (22), 57 (80) and 44 (8).

Ethyl 2-diethoxyphosphoryl-2-(3-methylbutanoylamino)acetate 7. Pale yellow crystalline solid (76%), mp 57–58 °C (Found: C, 48.4; H, 8.3; N, 4.3. C₁₃H₂₆NO₆P requires C, 48.3; H, 8.1; N, 4.3%) (Found: M⁺, 323.1499. C₁₃H₂₆NO₆P requires M, 323.1497); $\nu_{\max}(\text{CHCl}_3)/\text{cm}^{-1}$ 1710, 1677, 1221 and 1025; $\delta_{\text{H}}(360 \text{ MHz}; \text{CDCl}_3)$ 6.17 (1 H, br d, *J* 7.6, exch. D₂O, NH), 5.16 (1 H, dd, *J* 7.6 and 22.0, CHP), 4.20–4.16 (2 H, m, OCH₂), 4.10–4.06 (4 H, m, OCH₂), 2.09–2.05 (3 H, m, CH and CH₂), 1.25 (9 H, t, *J* 7.6, Me) and 0.90 (6 H, d *J* 7.6, Me₂CH); $\delta_{\text{C}}(67.8$

MHz; CDCl_3) 172.0, 166.6, 63.4 (d, J 67.8), 63.3 (d, J 67.8), 62.0, 50.2 (d, J 162.7), 45.1, 25.8, 22.2, 16.1 (d, J 6.8) and 13.8; δ_{p} (101.3 MHz; CDCl_3) 15.16; m/z (EI) 324 (MH^+ , 54%), 281 (66), 235 (42), 166 (100), 138 (72), 102 (38), 85 (65), 69 (22), 57 (99), 41 (83) and 29 (76).

Ethyl 2-diethoxyphosphoryl-2-[*N*-methylcarbamoyl]aminoacetate 8. Colourless crystalline solid (40%), mp 50–51 °C (from ether–light petroleum) (lit.,³⁷ 49–50 °C) (Found: M^+ , 296.1137. Calc. for $\text{C}_{10}\text{H}_{21}\text{N}_2\text{O}_6\text{P}$, M , 296.1137); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 3350, 2930, 1720, 1675, 1250 and 1030; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 6.04 (1 H, br s, exch. D_2O , NH), 5.21 (1 H, br s, exch. D_2O , NHMe), 5.09 (1 H, dd, J 9.3 and 22.54, CHP), 4.30–4.11 (6 H, m, OCH_2), 2.78 (3 H, d, J 4.4, MeNH) and 1.33 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 167.9, 158.2, 64.5, 63.7, 62.1, 51.4 (d, J 143.5), 26.9, 16.3 (d, J 6.2) and 14.0; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 16.64; m/z (CI) 314 ($\text{M} + \text{NH}_4^+$, 10%) and 297 (MH^+ , 52).

Ethyl 2-diethoxyphosphoryl-2-propylaminoacetate 9. Colourless oil (46%) (Found: M^+ , 281.1047. $\text{C}_{11}\text{H}_{24}\text{NO}_5\text{P}$ requires M , 281.1392); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1740 and 1257; $\delta_{\text{H}}(250 \text{ MHz}; \text{CDCl}_3)$ 5.38 (1 H, br s, exch. D_2O , NH), 5.36 (1 H, d, J 17.5, CHP), 4.28–4.11 (6 H, m, OCH_2), 3.12 (2 H, 2 overlapping t, CH_2N), 1.47 (2 H, m, $\text{CH}_2\text{CH}_2\text{N}$), 1.28 (9 H, m, 3 × Me) and 0.86 (3 H, t, J 7.5, Me), $\delta_{\text{C}}(62.9 \text{ MHz}; \text{CDCl}_3)$ 166.1, 68.6 (d, J 160.4), 63.7, 62.0, 42.0, 22.8, 16.2 (d, J 6.7), 13.9 and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 12.56; m/z (EI) 281 (M^+ , 2%), 280 (12), 240 (14), 197 (22), 167 (95), 155 (35), 138 (38), 111 (85), 69 (100) and 43 (23).

Ethyl 2-diethoxyphosphoryl-2-phenylaminoacetate 10. Colourless crystalline solid (72%), mp 56 °C (from ether–light petroleum) (lit.,⁵¹ 55–56 °C) (Found: M^+ , 315.1232. Calc. for $\text{C}_{14}\text{H}_{22}\text{NO}_5\text{P}$, M , 315.1236); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1720, 1240 and 1120; $\delta_{\text{H}}(400 \text{ MHz}; \text{CDCl}_3)$ 7.18 (2 H, t, J 8.4, ArH), 6.79 (1 H, t, J 7.2, ArH), 6.77 (2 H, d, J 8, ArH), 4.54 (1 H, s, exch. D_2O , NH), 4.51 (1 H, d, J 24, CHP), 4.26–4.17 (6 H, m, OCH_2) and 1.36–1.24 (9 H, m, Me); $\delta_{\text{C}}(100.6 \text{ MHz}; \text{CDCl}_3)$ 168.4, 146.2, 129.3, 120.0, 114.1, 64.3, 63.6, 62.2, 56.5 (d, J 147.6, CHP), 22.9, 16.4 and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 16.14; m/z (FAB) 316 (MH^+ , 29%), 315 (M^+ , 31), 242 (12), 178 (100), 139 (6), 104 (36) and 77 (5).

Ethyl 2-diethoxyphosphoryl-2-(4-methylphenylamino)acetate 11. Beige crystalline solid (73%), mp 76–77 °C (from ether–light petroleum) (Found: C , 54.8; H, 7.4; N, 3.8. $\text{C}_{15}\text{H}_{24}\text{NO}_5\text{P}$ requires C , 54.7; H, 7.35; N, 4.25%); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1725, 1255 and 1150; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 6.99 (2 H, d, J 8.30, ArH), 6.60 (2 H, d, J 8.30, ArH), 4.72 (1 H, br s, exch. D_2O , NH), 4.46 (1 H, d, J 23.2, CHP), 4.30–4.12 (6 H, m, OCH_2), 2.24 (3 H, s, ArMe) 1.39–1.32 (6 H, m, Me) and 1.27 (3 H, t, J 7.14, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 168.6, 143.8, 129.8, 128.7, 114.2, 64.0 (d, J 7.2), 63.5, 62.1, 56.8 (d, J 149.5), 20.4, 16.4 (d, J 6.7) and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 16.21; m/z (FAB) 330 (MH^+ , 35%), 329 (M^+ , 51), 256 (13), 192 (100) and 118 (47).

Ethyl 2-diethoxyphosphoryl-2-(4-methoxyphenylamino)acetate 12. Colourless crystalline solid (70%), mp 55–56 °C (from ether–light petroleum) (Found: M^+ , 345.1344. $\text{C}_{15}\text{H}_{24}\text{NPO}_6$ requires M , 345.1341); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1718, 1250 and 1130; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 6.86 (2 H, d, J 9.1, ArH), 6.74 (2 H, d, J 9.1, ArH), 4.55 (1 H, d, J 22.3, CHP), 4.37–4.32 (6 H, m, OCH_2), 3.82 (3 H, s, OMe) and 1.46–1.27 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 168.6, 153.5, 140.2, 115.7, 114.8, 64.0 (d, J 6.4, CH_2OP), 63.5 (d, J 7.3), 62.1, 57.5 (d, J 149.5), 55.5, 16.4 (d, J 6.8) and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 16.12; m/z (FAB) 346 (MH^+ , 49%), 315 (12), 272 (6), 225 (8), 208 (87), 134 (74), 93 (100), 75 (33) and 57 (26).

Ethyl 2-diethoxyphosphoryl-2-(4-nitrophenylamino)acetate 13. Pale yellow solid (81%), mp 96–97 °C (from ether–light petroleum) (Found: C , 46.8; H, 5.9; N, 7.6. $\text{C}_{14}\text{H}_{21}\text{N}_2\text{O}_7\text{P}$ requires C , 46.6; H, 5.9; N, 7.8%); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 2960, 1720, 1590, 1490, 1330, 1260 and 1120; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 8.22 (2 H, d, J 9.34, ArH), 6.75 (2 H, d, J 9.34, ArH), 5.38 (1 H, dd, J 8.25,

exch. D_2O , NH), 4.68 (1 H, dd, J 21.75 and 8.25, CHP), 4.35–4.24 (6 H, m, OCH_2) and 1.41 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 167.2, 151.2, 139.7, 126.2, 112.4, 64.2 (d, J 7), 63.9 (d, J 7), 62.8, 55.4 (d, J 147.4), 16.4 (d, J 6.8) and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 14.77; m/z (FAB) 361 (M^+ , 35%), 345 (47), 287 (20), 271 (15), 259 (10), 223 (82), 207 (52), 195 (13), 178 (23), 149 (70), 139 (100), 121 (27), 111 (42), 103 (35), 93 (30), 83 (42), 77 (13) and 65 (33).

Ethyl 2-(4-chlorophenylamino)-2-diethoxyphosphorylacetate 14. Colourless crystalline solid (71%), mp 80–81 °C (from ether–light petroleum) (Found: C , 48.0; H, 6.0; N, 4.1. $\text{C}_{14}\text{H}_{21}\text{ClNO}_5\text{P}$ requires C , 48.1; H, 6.05; N, 4.0%); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1715, 1245, 1130 and 1090; $\delta_{\text{H}}(400 \text{ MHz}; \text{CDCl}_3)$ 7.14 (2 H, d, J 8.79, ArH), 6.61 (2 H, d, J 8.79, ArH), 4.57 (1 H, br s, exch. D_2O , NH), 4.44 (1 H, d, J 22.27, CHP), 4.22–4.17 (6 H, m, OCH_2) and 1.36–1.24 (9 H, m, Me); $\delta_{\text{C}}(100.6 \text{ MHz}; \text{CDCl}_3)$ 168.1, 144.8, 129.2, 124.2, 115.2, 64.4, 64.1, 62.3, 56.6 (d, J 147.7), 16.4 (d, J 6.8) and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 15.78; m/z (FAB) 372 (MNa^+ , 41%), 350 (MH^+ , 27), 325 (12), 293 (8), 267 (58), 245 (35), 212 (38), 189 (24) and 167 (34).

Ethyl 2-diethoxyphosphoryl-2-(2-methoxyphenylamino)acetate 15. Colourless oil (74%) (Found: M^+ , 345.1341. $\text{C}_{15}\text{H}_{24}\text{NO}_6\text{P}$ requires M , 345.1341); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1720, 1250 and 1135; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 6.86–6.72 (3 H, m, ArH), 6.56 (1 H, d, J 7.8, ArH), 5.15 (1 H, d, J 9.0, exch. D_2O , NH), 4.55 (1 H, dd, J 11.2 and 22.6, CHP), 4.30–4.15 (6 H, m, OCH_2), 3.86 (3 H, s, OMe), 1.33 (6 H, dt, J 11.2 and 7.15, MeCH_2OP) and 1.27 (3 H, t, J 7.15, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 166.5, 147.6, 136.2, 121.0, 118.0, 110.9, 110.0, 63.9 (d, J 6.85), 63.3 (d, J 6.85), 62.1, 56.2, (d, J 147.2), 55.6, 16.4 (d, J 6.8) and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 16.18; m/z (CI) 346 (MH^+ , 27%), 345 (M^+ , 39), 272 (12), 208 (100 and 134 (42).

Ethyl 2-(2-bromophenylamino)-2-diethoxyphosphorylacetate 16. Pale yellow oil (70%) (Found: M^+ , 393.0357. $\text{C}_{14}\text{H}_{21}^{79}\text{BrNO}_5\text{P}$ requires M , 393.0341); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1720, 1250 and 1134; $\delta_{\text{H}}(400 \text{ MHz}; \text{CDCl}_3)$ 7.42 (1 H, d, J 9.25, ArH), 7.13 (1 H, d, J 8.75, ArH), 6.65–6.56 (2 H, m, ArH), 5.22 (1 H, br t, J 8.5, NH, exch. D_2O), 4.50 (1 H, dd, J 22.5 and 8.5, CHP), 4.25–4.16 (6 H, m, OCH_2), 1.34–1.21 (9 H, m, Me); $\delta_{\text{C}}(100.6 \text{ MHz}; \text{CDCl}_3)$ 167.7, 143.1, 132.6, 128.4, 119.7, 112.2, 110.7, 64.0 (d, J 6.1, CH_2OP), 63.7 (d, J 7.0), 62.3, 56.2 (d, J 156.3), 16.3 (d, J 6.8) and 14.0; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 15.19; m/z (EI) 395/393 (M^+ , 30%), 256/258 (90) and 182/184 (100).

Ethyl 2-diethoxyphosphoryl-2-(2,6-dimethylphenylamino)acetate 17. Colourless oil (71%) (Found: M^+ , 343.1546. $\text{C}_{16}\text{H}_{26}\text{NO}_5\text{P}$ requires M , 343.1549); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 1720, 1260 and 1130; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 6.95 (2 H, d, J 7.24, ArH), 6.80 (1 H, dd, J 7.98 and 8.55, ArH), 4.44 (1 H, d, J 23.37, CHP), 4.26–4.10 (6 H, m, OCH_2), 2.36 (6 H, s, ArMe), 1.34 (6 H, dt, J 7.15 and 6.88, Me) and 1.21 (3 H, t, J 7.15, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 169.1, 143.3, 129.1, 128.7, 122.3, 63.7 (d, J 7.3), 63.3 (d, J 7.3), 61.9, 58.8 (d, J 144.3), 18.6, 16.4, (d, J 6.2) and 14.0; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 16.87; m/z (CI) 344 (MH^+ , 100%) and 206 (4).

Ethyl 2-(2-chloro-4-trifluoromethylphenylamino)-2-diethoxyphosphorylacetate 18. Colourless oil (61%) (Found: M^+ , 417.0724. $\text{C}_{15}\text{H}_{20}^{35}\text{ClF}_3\text{NO}_5\text{P}$ requires M , 417.0720); $\nu_{\text{max}}(\text{CHCl}_3)/\text{cm}^{-1}$ 3370, 1743, 1293, 1241 and 1020; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 7.54 (1 H, s, ArH), 7.38 (1 H, d, J 8.5, ArH), 6.69 (1 H, d, J 8.5, ArH), 5.54 (1 H, br d, J 8.5), 4.55 (1 H, dd, J 8.2 and 21.2, CHP), 4.40–4.12 (6 H, m, OCH_2) and 1.39–1.29 (9 H, m, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 167.3, 144.8, 126.7, 125.9, 125.1, 121.9, 121.3, 120.8, 120.0, 111.3, 64.2 (d, J 7.3), 64.0 (d, J 7.3), 62.7, 55.8 (d, J 148.4), 16.4 (d, J 5.2) and 14.1; $\delta_{\text{p}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 14.68; m/z (FAB) 418 (MH^+ , 47%), 417 (M^+ , 45), 398 (22), 344 (28), 280 (100), 206 (85) and 139 (55). Compound 19 was also formed during the reaction.

Diethyl 1,2-bis(2-chloro-4-trifluoromethylphenylamino)-2-oxoethylphosphonate 19. Colourless oil (25%) (Found: M^+ , 566.0362. $\text{C}_{20}\text{H}_{19}^{35}\text{Cl}_2\text{F}_6\text{N}_2\text{O}_4\text{P}$ requires M , 566.0364);

$\nu_{\max}(\text{CHCl}_3)/\text{cm}^{-1}$ 1680, 1281 and 1026; $\delta_{\text{H}}(270 \text{ MHz}; \text{CDCl}_3)$ 9.22 (1 H, br s, NHCO), 8.55 (1 H, d, J 8.5, ArH), 7.62 (2 H, d, J 8.8, ArH) 7.55 (1 H, d, J 8.8, ArH), 7.42 (1 H, d, J 8.5, ArH), 6.73 (1 H, d, J 8.5, ArH), 5.79 (1 H, dd, J 5.2, J 9.9, CHNH), 4.51 (1 H, dd, J 5.2 and 21.7, CHP), 4.27 (4 H, dt, J 7.2 and 8.2, OCH₂) and 1.37 (6 H, q, J 7.2, Me); $\delta_{\text{C}}(67.8 \text{ MHz}; \text{CDCl}_3)$ 164.4, 144.8, 136.9, 129.7, 126.8, 126.4, 125.3, 125.0, 124.5, 123.3, 122.1, 121.2, 120.6, 111.9, 64.7 (d, J 7.3), 64.5 (d, J 7.3), 57.8 (d, J 144.4) and 16.4 (d, J 5.6); $\delta_{\text{P}}(101.3 \text{ MHz}; \text{CDCl}_3)$ 15.55; m/z (FAB) 589 (MNa⁺, 67%), 567 (MH⁺, 31), 429 (25), 345 (57), 206 (100) and 139 (73).

Acknowledgements

We thank Loughborough University and SmithKline Beecham Pharmaceuticals for their support of this research and the EPSRC for funds towards the purchase of a 400 MHz NMR spectrometer. We also thank the EPSRC Mass Spectrometry Centre at Swansea for mass spectra.

References

- 1 For Part 1, see *J. Chem. Soc., Perkin Trans. 1*, 1996, 2879, preceding paper.
- 2 Preliminary communication, L. Ferris, D. Haigh and C. J. Moody, *Synlett*, 1995, 921.
- 3 R. W. Ratcliffe and B. G. Christensen, *Tetrahedron Lett.*, 1973, 4645.
- 4 R. W. Ratcliffe and B. G. Christensen, *Tetrahedron Lett.*, 1973, 4649.
- 5 L. S. Hegedus, L. M. Schultze, J. Toro and C. Yijun, *Tetrahedron*, 1985, **41**, 5833.
- 6 P. Herdewijn, P. J. Claes and H. Vanderhaeghe, *J. Med. Chem.*, 1986, **29**, 661.
- 7 T. Ogasa, H. Saito, Y. Hashimoto, K. Sato and T. Hirata, *Chem. Pharm. Bull.*, 1989, **37**, 315.
- 8 Y. Narukawa, K. N. Juneau, D. Snustad, D. B. Miller and L. S. Hegedus, *J. Org. Chem.*, 1992, **57**, 5453.
- 9 B. T. Lotz and M. J. Miller, *J. Org. Chem.*, 1993, **58**, 618.
- 10 U. Schmidt, A. Lieberknecht, U. Schanbacher, T. Beuttler and J. Wild, *Angew. Chem., Int. Ed. Engl.*, 1982, **21**, 776.
- 11 U. Schmidt, A. Lieberknecht and J. Wild, *Synthesis*, 1984, 53.
- 12 C. Shin, T. Obara, S. Segami and Y. Yonezawa, *Tetrahedron Lett.*, 1987, **28**, 3827.
- 13 C. Shin, Y. Yonezawa, T. Obara and H. Nishio, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 885.
- 14 P. G. Ciattini, E. Morera and G. Ortari, *Synthesis*, 1988, 140.
- 15 D. Kim, Y. Li, B. A. Horenstein and K. Nakanishi, *Tetrahedron Lett.*, 1990, **31**, 7119.
- 16 C. Shin, N. Takahashi and Y. Yonezawa, *Chem. Pharm. Bull.*, 1990, **38**, 2020.
- 17 E. J. Moran and R. W. Armstrong, *Tetrahedron Lett.*, 1991, **32**, 3807.
- 18 U. Schmidt, A. Kleefeldt and R. Mangold, *J. Chem. Soc., Chem. Commun.*, 1992, 1687.
- 19 U. Schmidt, H. Griesser, V. Leitenberger, A. Lieberknecht, R. Mangold, R. Meyer and B. Riedl, *Synthesis*, 1992, 487.

- 20 R. W. Armstrong, J. E. Tellew and E. J. Moran, *J. Org. Chem.*, 1992, **57**, 2208.
- 21 U. Schmidt and B. Riedl, *Synthesis*, 1993, 815.
- 22 R. S. Coleman and A. J. Carpenter, *J. Org. Chem.*, 1993, **58**, 4452.
- 23 J. E. Baldwin, K. D. Merritt and C. J. Schofield, *Tetrahedron Lett.*, 1993, **34**, 3919.
- 24 H.-J. Kreuzfeld, C. Döbler, H. W. Krause and C. Facklam, *Tetrahedron: Asymmetry*, 1993, **4**, 2047.
- 25 K. Sato, T. Miyata, I. Tanai and Y. Yonezawa, *Chem. Lett.*, 1994, 129.
- 26 R. C. Holcomb, S. Schow, S. Ayril-Kaloustian and D. Powell, *Tetrahedron Lett.*, 1994, **35**, 7005.
- 27 M. Seki, K. Kondo and T. Iwasaki, *J. Chem. Soc., Perkin Trans. 1*, 1996, 3; M. Seki and K. Matsumoto, *Synthesis*, 1996, 580.
- 28 R. W. Armstrong, J. E. Tellew and T. R. Hoye, *Tetrahedron Lett.*, 1996, **37**, 447.
- 29 H. Gross and J. Freiberg, *Angew. Chem., Int. Ed. Eng.*, 1965, **4**, 975.
- 30 R. Kober and W. Steglich, *Liebigs Ann. Chem.*, 1983, 599.
- 31 B. Ku and D. Y. Oh, *Tetrahedron Lett.*, 1988, **29**, 4465.
- 32 M. Dumas, L. Vo-Quang and E. L. Goffic, *Synth. Commun.*, 1990, **22**, 3395.
- 33 R. Shankar and A. I. Scott, *Tetrahedron Lett.*, 1993, **34**, 231.
- 34 M. Vaultier, M. S. Ouali and R. Carrié, *Bull. Soc. Chim. Fr.*, 1979, 343.
- 35 D. I. C. Scopes, A. F. Kluge and J. A. Edwards, *J. Org. Chem.*, 1977, **42**, 376.
- 36 E. W. Colvin, G. W. Kirby and A. C. Wilson, *Tetrahedron Lett.*, 1982, **23**, 3835.
- 37 P. S. Khokhlov, B. A. Kashemirov, A. D. Mikityuk and Y. A. Strepikheev, *Zh. Obshch. Khim.*, 1983, **53**, 2146.
- 38 C. Shiraki, H. Saito, K. Takahashi, C. Urakawa and T. Hirata, *Synthesis*, 1988, 399.
- 39 G. H. Hakimelahi and G. Just, *Synth. Commun.*, 1980, **10**, 429.
- 40 For a comprehensive list of references, see *J. Chem. Soc., Perkin Trans. 1*, 1996, 2879, preceding paper.
- 41 M. Regitz, W. Anshütz and A. Liedhegener, *Chem. Ber.*, 1968, **101**, 3734.
- 42 C. J. Moody, E.-R. H. B. Sie and J. J. Kulagowski, *Tetrahedron*, 1992, **48**, 3991.
- 43 M. McGuinness and H. Shechter, *Tetrahedron Lett.*, 1990, **31**, 4987.
- 44 D. J. Miller, Loughborough University, unpublished result.
- 45 G. G. Cox, D. J. Miller, C. J. Moody, E.-R. H. B. Sie and J. J. Kulagowski, *Tetrahedron*, 1994, **50**, 3195.
- 46 G. Bartels, R.-P. Hinze and D. Wullbrandt, *Liebigs Ann. Chem.*, 1980, 168.
- 47 G. Brooks, T. T. Howarth and E. Hunt, *J. Chem. Soc., Chem. Commun.*, 1981, 642.
- 48 P. G. Mattingly and M. J. Miller, *J. Org. Chem.*, 1981, **46**, 1557.
- 49 T. Kametani, N. Kanaya, T. Mochizuki and T. Honda, *Heterocycles*, 1982, **19**, 1023.
- 50 M. Hrytsak and T. Durst, *Heterocycles*, 1987, **26**, 2393.
- 51 D. Haigh, *Tetrahedron*, 1994, **50**, 3177.

Paper 6/04713K

Received 5th July 1996

Accepted 13th August 1996