

Experimental Section

1-Benzyl-1*H*-triazole-4,5-dicarboxaldehyde (I) was prepared according to the procedure reported by Henkel and Weygand (5). The cyclic ketones were prepared according to the methods of Blomquist (6, 7). Melting points were determined with a Thomas-Hoover Unimelt instrument and are uncorrected. NMR spectra were recorded on a Varian A-60 spectrometer using tetramethylsilane as an internal reference and shifts (δ) are reported in ppm.

1-Benzyl-5,7-dodecano-6 (2*H*)-cycloheptatriazolone (III). A solution of 2.15 g (0.01 mol) of 1-benzyl-1*H*-triazole-4,5-dicarboxaldehyde and 2.24 g (0.01 mol) of cyclopentadecanone in 0.3 g of KOH and 50 mL of methanol was heated under reflux. After the mixture was cooled, the product was collected and then recrystallized from ethanol.

4-Hydroxy-1-benzyl-4,5-dihydro-5,7-polymethyleno-6-(2*H*)-cycloheptatriazolones (IVa-c). In the same manner 2.15 g (0.01 mol) of 1-benzyl-1*H*-triazole-4,5-dicarboxaldehyde and 0.01 mol of the corresponding cyclic ketones in 0.4 g of KOH and 60 mL of methanol was heated under reflux. The

solvent was removed under pressure and the resulting product was recrystallized from chloroform-petroleum ether.

The melting points, reflux time, yields, and NMR data for compounds III and IVa-c are listed in Table I.

Registry No. I, 103532-75-8; IIa, 1502-06-3; IIb, 878-13-7; IIc, 830-13-7; IId, 502-72-7; III, 105309-39-5; IVa, 105309-40-8; IVb, 105309-41-9; IVc, 105309-42-0.

Literature Cited

- (1) Abu-Orabi, S. T.; Harmon, R. E. *J. Chem. Eng. Data* **1986**, *31*, 379.
- (2) Abu-Orabi, S. T.; Harmon, R. E. *J. Chem. Eng. Data* **1986**, *31*, 504.
- (3) Kloster-Jensen, V. E.; Tarköy, N.; Eschenmoser, A.; Heilbronner, E. *Helv. Chim. Acta* **1956**, *39*, 786.
- (4) Harmon, R. E.; Suder, R.; Gupta, S. K. *Chem. Commun.* **1969**, 1170.
- (5) Henkel, H.; Weygand, F. *Chem. Ber.* **1943**, *76*, 812.
- (6) Blomquist, A. T.; Spencer, R. D. *J. Am. Chem. Soc.* **1948**, *70*, 30.
- (7) Blomquist, A. T.; Prager, J.; Wollinsky, J. J. *Am. Chem. Soc.* **1955**, *77*, 1804.

Received for review May 13, 1986. Revised July 25, 1986. Accepted August 25, 1986. I thank Yarmouk University for financial support of this work.

Synthesis of Some New 2,4,6-Triarylpyridines Using Phenacylidenedimethylsulfuranes

Komal C. Gupta,* Pankaj K. Pathak, Brijesh K. Saxena, Nirupma Srivastava, and Kalpna Pandey

Department of Chemistry, D.V. (P.G.) College, Orai 285 001, U.P., India

Phenacylidenedimethylsulfurane and *p*-chlorophenacylidenedimethylsulfurane were reacted with α,β -unsaturated ketones in the presence of ammonium acetate in glacial acetic acid to give 2,4,6-triarylpyridines in 45–70% yields. The structures of the pyridines were confirmed by IR and NMR spectra.

In continuation of our earlier researches (1–6) on the synthetic potentialities of ylides, we now report herein the aza ring closure reaction of two phenacylidenedimethylsulfuranes with a wide variety of α,β -unsaturated ketones with a view to test the domain of applicability of these ylides.

Experimental Section


Phenacyldimethylsulfonium bromide (1a) and *p*-chlorophen-

Table I. Physical and Spectral Data of Compounds^a

compd	R ₁	R ₂	R ₃	yield, %	recryst solvent	mp, °C	IR (KBr), ^b cm ⁻¹				NMR ^c data (CDCl ₃) δ , ppm
							ν (Ar-H)	ν (C=C)	ν (C=N)	ϕ (C-H)	
5a	H	H	4-FC ₆ H ₄	45	a	150–52	3150	1605	1500	995	5.95 (s, 2H, -O ₂ CH ₂); 6.40–8.00 (m, 14 H, Ar-H)
5b	H	3,4-O ₂ CH ₂	4-FC ₆ H ₄	60	b	160–62	3010	1610	1508	995	
5c	H	3,4-di-OCH ₃	4-FC ₆ H ₄	60	a	120–22	3000	1608	1500	992	3.70–7.75 (d ($J = 5$ Hz), 6 H, diOCH ₃); 6.58–7.95 (m, 14 H, ArH)
5d	4-Cl	H	4-FC ₆ H ₄	55	b	168–70	3060	1600	1505	1005	3.70 (s, 3 H, OCH ₃); 6.40–8.00 (m, 15 H, ArH)
5e	4-ClC ₆ H ₄	4-CH ₃ OC ₆ H ₄	C ₆ H ₅	50	c	95–96	3030	1510	1520	986	
5f	4-ClC ₆ H ₄	3-NO ₂ -C ₆ H ₄	C ₆ H ₅	65	d	90–92	3060	1595	1005	1005	3.75 (s, 3 H, OCH ₃); 6.50–8.30 (m, 14 H, ArH)
5g	4-ClC ₆ H ₄	4-CH ₃ O-C ₆ H ₄	4-NO ₂ C ₆ H ₄	60	a	82–85	3040	1598	1510	988	
5h	4-ClC ₆ H ₄	3-NO ₂ C ₆ H ₄	4-CH ₃ OC ₆ H ₄	65	c	140–42	3030	1585	1510	985	3.85 (s, 3 H, OCH ₃); 7.15–8.25 (m, 13 H, ArH)
5i	4-ClC ₆ H ₄	2-C ₄ H ₃ S	3-CH ₃ OC ₆ H ₄	70	f	70–72	3045	1590	1520	975	
5j	4-ClC ₆ H ₄	2-C ₄ H ₃ S	4-ClC ₆ H ₄	60	e	128–30	3015	1588	1510	1010	3.85 (s, 3 H, OCH ₃); 7.15–8.25 (m, 13 H, ArH)
5k	4-ClC ₆ H ₄	2-C ₄ H ₃ O	4-ClC ₆ H ₄	60	a	126–28	3050	1615	1500	980	
5l	4-ClC ₆ H ₄	3-NO ₂ C ₆ H ₄	4-ClC ₆ H ₄	65	c	84–85	3080	1600	1520	985	3.85 (s, 3 H, OCH ₃); 7.15–8.25 (m, 13 H, ArH)
5m	4-ClC ₆ H ₄	4-CH ₃ OC ₆ H ₄	2-C ₄ H ₃ O	60	c	96–98	3025	1618	1516	990	
5n	4-ClC ₆ H ₄	4-FC ₆ H ₄	2-C ₄ H ₃ S	60	a	50–55	3050	1590	1500	980	

^a All compounds gave satisfactory elemental analysis for C, H, N. ^b IR spectra were recorded on Perkin Elmer Infracord spectrometer using KBr phase. ^c NMR spectra were run on Varian A-60 spectrometer using Me₄Si as internal standard. ν = stretching vibrations, ϕ = bending vibrations (out-of-plane vibrations), s = singlet, m = multiplet. Recrystallization solvent: a = C₅H₅N-MeOH; b = EtOH; c = C₅H₅N-EtOH; d = C₆H₆-AcOEt; e = C₅H₅N-CHCl₃; f = CHCl₃-C₆H₆.

Scheme I



acyldimethylsulfonium bromide (**1b**) and their sulfuranes (**2a-b**) were prepared as reported in the literature (7-9). All α,β -unsaturated ketones (**3**) were synthesized by using the Claisen Schmidt method (10) (Scheme I).

Preparation of 2,4,6-Triarylpyridine (5). To a stirred mixture of salt (**1a-b**) (3 mmol) and unsaturated ketone (**3a**) (3

mmol) was added ammonium acetate (3 g) in 20 mL of glacial acetic acid. The resulting mixture was refluxed for 5-8 h. and left overnight at room temperature. The mixture was then poured into ice cold water (50 mL) to precipitate a solid which was filtered and washed twice with water and then with methanol. The solid mass was dried and recrystallized from pyridine-methanol or other suitable solvents shown in Table I.

Registry No. **1a**, 5667-47-0; **1b**, 19158-57-7; **3a**, 399-10-0; **3b**, 7397-23-1; **3c**, 28081-14-3; **3e**, 959-33-1; **3f**, 614-48-2; **3g**, 6552-62-1; **3h**, 68063-55-8; **3i**, 105372-58-5; **3j**, 6028-91-7; **3k**, 14385-65-0; **3l**, 22027-92-5; **3m**, 19430-55-8; **3n**, 367-00-0; **5a**, 91232-30-3; **5b**, 105372-47-2; **5c**, 105372-48-3; **5d**, 105372-49-4; **5e**, 74918-89-1; **5f**, 105372-50-7; **5g**, 105372-51-8; **5h**, 105372-52-9; **5i**, 105372-53-0; **5j**, 105372-54-1; **5k**, 105372-55-2; **5l**, 105372-56-3; **5m**, 105372-57-4; **5n**, 85957-62-6.

Literature Cited

- (1) Tewari, R. S.; Awasthi, A. K. *Synthesis* **1981**, 314.
- (2) Gupta, K. C.; Srivastava, N.; Nigam, R. K. *J. Org. Chem.* **1981**, *104*, 91.
- (3) Gupta, K. C.; Srivastava, N.; Nigam, R. K. *Curr. Sci.* **1983**, *52*, 719.
- (4) Gupta, K. C.; Srivastava, N.; Nigam, R. K. *Indian J. Chem.* **1981**, *20B*, 802.
- (5) Tewari, R. S.; Gupta, K. C. *Indian J. Chem.* **1976**, *14B*, 829.
- (6) Gupta, K. C.; Nigam, R. K.; Malik, S.; Srivastava, N. *Indian J. Chem.* **1981**, *21B*, 241.
- (7) Johnson, A. W.; Amel, R. T. *J. Org. Chem.* **1969**, *34*, 1240.
- (8) Ratts, K. W.; Yao, A. N. *J. Org. Chem.* **1966**, *31*, 1185.
- (9) Trost, B. M. *J. Am. Chem. Soc.* **1962**, *89*, 138.
- (10) Gulman, H.; Blatt, A. H. *Org. Synth.* **1958**, *1*, 78.

Received for review November 15, 1985. Revised March 4, 1986. Accepted August 22, 1986.