


ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: Y. Dudognon, M. Presset, J. Rodriguez, Y. Coquerel, X. Bugaut and T. Constantieux, *Chem. Commun.*, 2016, DOI: 10.1039/C5CC10217K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Chemical Communications

COMMUNICATION

Addition of Silylated Nucleophiles to α -Oxoketenes

Yohan Dudognon,^a Marc Presset,^{a,b} Jean Rodriguez,^a Yoann Coquerel*,^a Xavier Bugaut*^a and Thierry Constantieux*^a

Received 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

www.rsc.org/chemcomm

A general evaluation of silylated nucleophiles to intercept transient α -oxoketenes generated by microwave-assisted Wolff rearrangement of 2-diazo-1,3-dicarbonyl compounds is presented. Original scaffolds and synthetic intermediates are accessed in a rapid, efficient and easy-to-handle way. Mechanistic studies by DFT calculations and various post-functionalizations are discussed.

The 1,2-Wolff rearrangement, i.e. the conversion of easily available α -diazocarbonyl derivatives into ketenes with extrusion of dinitrogen, is a venerable and useful synthetic tool (Scheme 1).^{1,2} It has been extensively used for the functionalization and variation of carbon backbones, especially for the ring contraction of cyclic molecules and the Arndt-Eistert homologation of carboxylic acids.³ Starting from 2-diazo-1,3-dicarbonyl compounds **1**,⁴ the Wolff rearrangement produces strongly electrophilic α -oxoketene species **2**,⁵ which are usually not isolable and must be generated in situ. α -Oxoketenes can react with a variety of heteroatomic or carbon-

centered nucleophiles to produce the corresponding 1,3-dicarbonyl products,^{6,7} and with various unsaturated compounds to afford stereodefined polycyclic molecules following pericyclic processes (Scheme 1, left).⁸ Microwave-assisted heating was shown to be highly beneficial in terms of efficiency and practicability in these transformations.⁹ However, the chemoselective interception of α -oxoketenes with simple nucleophiles such as dihydrogen, hydrogen azide or ammonia remains unknown (Scheme 1, upper right). We identified the following issues as possible explanations: (i) reactivity and chemoselectivity, since the nucleophile has to react with the α -oxoketene but not with the ketones of the starting materials; (ii) practicability for reactions at elevated temperature at the lab-scale, as most of those nucleophiles are gases, often with high toxicities; (iii) product stability since the resulting naked functionalities in the products (e.g. aldehydes and acyl azides), might be unstable under the reaction conditions. The alternative anionic species (such as



Scheme 1. Previous and new applications of the microwave-assisted Wolff rearrangement

^a Aix Marseille Université, Centrale Marseille, CNRS, iSm2 UMR 7313, 13397, Marseille, France. E-mail: yoann.coquerel@univ-amu.fr, xavier.bugaut@univ-amu.fr, thierry.constantieux@univ-amu.fr; Fax: +33 491 289 187; Tel: +33 491 282 874

^b Present address: Université Paris Est, ICMPE (UMR 7182), CNRS, UPEC, F-94320 Thiais, France.

†Electronic Supplementary Information (ESI) available: All spectroscopic data and

View Article Online

mineral cyanide or azide sources) can barely be used since they are hardly soluble in the unpolar organic reaction media commonly used to carry out the Wolff rearrangement.

Despite the widespread application of silylated nucleophiles in organic chemistry (hydrosilylation,¹⁰ Mukaiyama aldol reaction,¹¹ Peterson elimination,¹² Hiyama-Denmark cross-coupling¹³ etc...), they have hardly been reacted with ketene species. Indeed, there are only a single example of trapping of a ketene with hexamethyldisilazane (HMDS) to afford the corresponding primary amide in a context of total synthesis,¹⁴ and a study on the hydrosilylation of bench-stable polyfluoroketenes under Pt-catalysis.¹⁵ Additionally, a recent report presents the coupling of 1,3-dioxin-2-ones with silyl enol ethers.¹⁶ In the present work, we describe the first chemoselective addition of a variety of easy-to-handle silylated nucleophiles to α -oxoketenes generated by microwave-assisted Wolff rearrangement of either acyclic or cyclic 2-diazo-1,3-dicarbonyl compounds, to prepare building blocks that are hard or impossible to access by alternative existing methods (Scheme 1, bottom right). Triethylsilane and trimethylsilyl azide were used as sources of hydride and azide, to synthesize masked β -ketoaldehydes **3** and a β -ketoacyl azide **5**, respectively. In addition, HMDS acted as a liquid anhydrous ammonia surrogate and trimethylsilyldiazomethane as a source of methylene to afford upon desilylation primary β -ketoamides **4** and fused bicyclic furan-3-ones **6**, respectively. In these reactions, the silyl group remarkably enabled the in situ protection of sensitive functionalities avoiding their degradation during the reaction.

We started our investigation by mixing triethylsilane and 2-diazo-5,5-dimethylcyclohexane-1,3-dione in toluene at 160 °C under microwave irradiation during 3 minutes (Table 1).

Table 1. Synthesis of masked β -ketoaldehydes by triethylsilane addition^a

 3a , 98% (>20:1 dr) ^b	 3b , 91% (>20:1 dr) ^c	 3c , 95% (>20:1 dr) ^c
 3d , 97% (>20:1 dr)	 3e , 92% (1.6:1 dr)	 3f , 98% (1.5:1 dr) ^d
 3g , 96% (10:1 dr) ^d	 3h , 85% (3.5:1 dr)	 3i , 81% (>20:1 dr)

^a A solution of diazo compound (0.25 mmol, 1 equiv) and triethylsilane (1 equiv) was stirred at 160 °C for 3 min in toluene (2 mL). ^b The reactions were conducted on a 1 mmol scale. ^c The reactions were conducted on a 0.5 mmol scale. ^d The reactions were conducted at 200 °C for 3 min.

Pleasingly, analysis of the clean crude reaction mixture showed only the masked β -ketoaldehyde **3a** as a single diastereomer (98% yield), which was identified by comparison with known related compounds.¹⁷ This methodology could be applied to the synthesis of various cyclic masked β -ketoaldehydes by ring contraction (products **3b** to **3g**), such as functionalized thiolanone **3e**, oxindole **3f** and benzoxofurane **3g**. The acyclic products **3h** and **3i** could also be prepared from the corresponding acyclic diazo compounds. Notably, all those compounds were obtained with good to excellent yields in a practical and expeditious manner (reaction time = 3 min, no work-up, no purification). Generally, only the (*E*)-diastereomer was observed, but for several examples, the (*Z*) diastereomer could also be detected. The use of less nucleophilic silicium hydrides (Ph₃SiH, Ph₂SiH₂ and (EtO)₃SiH) resulted in complex reaction mixtures.¹⁸ Remarkably, with this methodology, the position of the protected aldehyde functionality in compounds **3a-c** and **3e** is fully controlled since the starting diazo compounds are symmetric molecules, whereas other existing methods to prepare this motif, such as the formylation of enolate, are plagued with problems of regioselectivity.^{17h}

The outcome of the reaction raised mechanistic considerations, which were investigated by computational DFT methods in the case of the model reaction between the conformationally constrained α -oxoketene **A** and trimethylsilane (Figure 1).¹⁹ In early calculations, both the *endo* and *exo* direct 1,2-hydrosilylation of the ketene carbonyl group reaction paths were computed, and the formation of (*E*) isomer **C** was found both kinetically and thermodynamically favored over the (*Z*) isomer **B** at the studied level of theory (Figure 1a). However, in both cases activation barriers are high (134 and 110 kJ.mol⁻¹, respectively), and an alternative reaction pathway was investigated (Figure 1b). The hydrosilylation reactions described in Table 1 are believed to occur via a three-step mechanism: (i) a concerted kinetically favored 1,4-hydrosilylation of the α -oxoketene produces intermediate **D** through a six-membered transition state **TS_{AD}**;²⁰ (ii) a 1,5-shift of the silyl group from the ketone enol ether oxygen atom to the aldehyde one in **D** is possible through another six-membered transition state **TS_{DB}** to give the (*Z*) isomer **B**; (iii) the thermodynamic (*E*) isomer product **C** can be obtained by an isomerization of the (*Z*) isomer **B**. The unusual **B** → **C** isomerization has a reasonable energy barrier due to the marked polarization of the enol ether double bond with a “push-pull” motif allowing for a transition state similar to an enolate/silyl-stabilized oxonium system.^{21,22}

We then turned our attention to nitrogen-centered silylated nucleophiles, using HMDS and trimethylsilyl azide (TMSN₃). The reactions of representative α -oxoketenes with HMDS afforded relatively complex crude reaction mixtures, which after purification by silica gel chromatography delivered the corresponding cyclic and acyclic primary β -ketoamides **4a-4e** in satisfactory yields (Table 2).²³






Table 2. Synthesis of β -ketoamides by HMDS addition^a



^aA solution of diazo compound (0.1 mmol, 1 equiv) and HMDS (1 equiv) was stirred at 160 °C for 3 min in toluene (2 mL). Yields of isolated pure products after purification by flash chromatography on silica gel. ^bThis reaction was conducted on a 0.5 mmol scale. ^c This reaction was conducted on a 0.25 mmol scale.

The reaction with TMSN₃ was less straightforward but, after optimization of the reaction conditions involving the portionwise addition of the diazo compound obtained from dimedone, the previously unknown masked β -ketoacyl azide **5** was obtained diastereoselectively with 86% yield (Scheme 2).¹⁹




Scheme 2. Addition of TMSN₃

To complete the study, carbon-centered silylated nucleophiles were investigated through the reactions with trimethylsilyl cyanide (TMSCN), trimethyl(trifluoromethyl)silane (TMSCF₃) and trimethylsilyldiazomethane (TMSCHN₂). With TMSCN, the expected product could be observed but it was not possible to optimize the reaction to a reasonable level of selectivity. As to the reaction with


TMSCF₃, only the α -oxoketene dimer was obtained, presumably because of its too weak nucleophilicity (not depicted). On the opposite, TMSCHN₂ behaved as a convenient source of methylene (Table 3): indeed, after initial nucleophilic trapping, the second diazo function underwent an insertion reaction to deliver original fused bicyclic furan-3-ones **6a-c**, whose synthesis has scarcely been reported,²⁴ with satisfactory yield after purification by silica gel chromatography. It is worth noting that related preliminary evaluation of the coupling between two different diazo species can be found in the literature but yields of products were low because of other competing pathways.²⁵ The present results highlight the beneficial effect of the silyl group to prevent over-reaction of the product under the reaction conditions. Moreover this transformation represents a selective cross-coupling of two different diazo compounds.

Table 3. Synthesis of bicyclicfuran-3-ones by TMSCHN₂ addition^a



^aA solution of diazo compound (0.1 mmol, 1 equiv) and TMSCHN₂ (1 equiv) was stirred at 160 °C for 3 min in toluene (2 mL). Yields of isolated pure products after purification by flash chromatography on silica gel. ^bThis reaction was conducted on a 0.5 mmol scale. ^c This reaction was conducted on a 0.25 mmol scale.

With this set of original structures in hand, we briefly investigated their potential post-functionalization to obtain more structurally diverse scaffolds. Starting from masked β -ketoaldehyde **3a**, Michael addition on methyl vinyl ketone in slightly basic hydro-organic reaction conditions¹⁹ delivered with good yield product **7**, which could be further functionalized into spiro-cyclohexenone **8** by Robinson annulation, also with good yield.



Scheme 3. Post-functionalization of masked β -ketoaldehyde **3a**

In conclusion, we have successfully developed a versatile method for the addition of silylated nucleophiles to α -oxoketenes generated by microwave-assisted Wolff rearrangement of 2-diazo-1,3-dicarbonyl compounds to synthesize valuable building blocks that are hard or impossible to access by existing methods. This methodology is a rapid, efficient and user-friendly way to prepare original scaffolds and synthetic intermediates that are likely to find interesting applications in synthetic organic chemistry. Additionally, DFT calculations are in agreement with an initial 1,4-hydrosilylation with participation of the ketone's oxygen atom. Our current efforts aim to find applications for these promising substrates in enantioselective organocatalysis.

We thank R. Rosas, Dr. N. Vanthuyne and M. Jean, and the whole team of Spectropole (www.spectropole.fr). Financial support from Aix Marseille Université, Centrale Marseille and the CNRS is acknowledged.

Notes and references

- 1 L. Wolff, *Justus Liebigs Ann. Chem.*, 1902, **325**, 129-195.
- 2 (a) W. Kirmse, *Eur. J. Org. Chem.*, 2002, 2193-2256; (b) Y. Coquerel and J. Rodriguez, in *Molecular Rearrangements in Organic Synthesis*, Ed.: C. Rojas, John Wiley & Sons, Hoboken, NJ, **2015**, Chap. 3, pp. 59-84.
- 3 F. Arndt, B. Eistert, *Ber.*, 1935, **68b**, 200-208.
- 4 M. Pisset, D. Mailhol, Y. Coquerel, J. Rodriguez, *Synthesis*, 2011, **16**, 2549-2552.
- 5 Reviews: (a) C. Wentrup, W. Heilmayer and G. Kollenz, *Synthesis*, 1994, **12**, 1219-1248; (b) G. Kollenz and S. Ebner, in *Science of Synthesis: Houben-Weyl Methods of Molecular Transformations*, Danheiser, R., Ed., Georg Thieme Verlag, Stuttgart, Germany, **2006**, Vol. 23, pp. 271-349; (c) K. P. Reber, S. D. Tilley and E. J. Sorensen, *Chem. Soc. Rev.*, 2009, **38**, 3022-3034; (d) A. Ford, H. Miel, A. Ring, C. N. Slattey, A. R. Maguire and M. A. McKervey, *Chem. Rev.*, 2015, **115**, 9981-10080.
- 6 (a) M. Pisset, Y. Coquerel and J. Rodriguez, *J. Org. Chem.*, 2009, **74**, 415-418; (b) T. Boddaert, Y. Coquerel and J. Rodriguez, *Eur. J. Org. Chem.*, 2011, 5061-5070; (c) J.-C. Castillo, M. Pisset, R. Abonia, Y. Coquerel and J. Rodriguez, *Eur. J. Org. Chem.*, 2012, 2338-2345.
- 7 Y. Coquerel, K. Mohanan, M. Pisset, D. Mailhol and J. Rodriguez, *Chem. Eur. J.*, 2012, **18**, 9217-9220.
- 8 (a) M. Pisset, Y. Coquerel and J. Rodriguez, *Org. Lett.*, 2009, **11**, 5706-5709; (b) M. Pisset, Y. Coquerel and J. Rodriguez, *Org. Lett.*, 2010, **12**, 4212-4215; (c) M. Pisset, K. Mohanan, M. Hamann, Y. Coquerel and J. Rodriguez, *Org. Lett.*, 2011, **13**, 4124-4127; (d) J. Galvez, J.-C. Castillo, J. Quiroga, M. Rajzmann, J. Rodriguez and Y. Coquerel, *Org. Lett.*, 2014, **16**, 4126-4129; (e) P. Neupane, L. Xia and Y. R. Lee, *Adv. Synth. Catal.*, 2014, **356**, 2566-2574.
- 9 (a) C. O. Kappe, D. Dallinger and S. S. Murphree, *Practical Microwave Synthesis for Organic Chemists*, Wiley-VCH, Weinheim, 2009; (b) Y. Coquerel, E. Colacino, J. Rodriguez, J. Martinez and F. Lamaty, in *Stereoselective Synthesis of Drugs and Natural Products*, eds. V. Andrushko and N. Andrushko, John Wiley & Sons, Hoboken, NJ, 2013, Chap. 5, pp. 145-166.
- 10 Y. Nakajima and S. Shimada, *RSC Adv.*, 2015, **5**, 20603-20616.
- 11 (a) T. Mukaiyama, K. Narasaka and K. Banno, *Chem. Lett.*, 1973, 1011-1014; (b) J. Matsuo and M. Murakami, *Angew. Chem. Int. Ed.*, 2013, **52**, 9109-9118.
- 12 (a) D. J. Peterson, *J. Org. Chem.*, 1968, **33**, 780-784; (b) L. F. v. Staden, D. Gravestock and D. J. Ager, *Chem. Soc. Rev.*, 2002, **31**, 195-200.
- 13 (a) Y. Hatanaka and T. Hiyama, *J. Org. Chem.*, 1988, **53**, 918-920; (b) H. F. Sore, W. R. J. D. Galloway and D. R. Spring, *Chem. Soc. Rev.*, 2012, **41**, 1845-1866.
- 14 L. N. Mander and M. M. McLachlan, *J. Am. Chem. Soc.*, 2003, **125**, 2400-2401.
- 15 (a) A. Y. Volkonskii, E. M. Kagramanova, E. I. Mysov and N. E. Mysova, *Russ. Chem. Bull.*, 2004, **53**, 1693-1699. (b) A. Y. Volkonskii, E. M. Kagramanova, E. I. Mysov and N. D. Kagramanov, *Russ. Chem. Bull.*, 2010, **59**, 569-576.
- 16 Q. Wang and B. List, *Synlett*, 2015, **26**, 1525-1527.
- 17 (a) P. J. Stang and W. L. Treptow, *J. Med. Chem.*, 1981, **24**, 468-472; (b) M. J. Suto, K. M. Trampusch, M. Wierzb, A. J. Solo and W. Duax, *J. Org. Chem.*, 1987, **52**, 2263-2273; (c) L. F. Tietze, A. Bergmann, G. Brill, K. Brüggemann, U. Hartfiel and E. Voß, *Chem. Ber.*, 1989, **122**, 83-94; (d) E. Krawczyk and A. Skowrońska, *Heteroat. Chem.*, 2000, **11**, 353-361; (e) P. T. O'Sullivan, W. Buhr, M. A. M. Fuhry, J. R. Harrison, J. E. Davies, N. Feeder, D. R. Marshall, J. W. Burton and A. B. Holmes, *J. Am. Chem. Soc.*, 2004, **126**, 2194-2207; (f) L. Wan and M. A. Tius, *Org. Lett.*, 2007, **9**, 647-650; (g) E. M. B. I. Janke, S. Schlund, A. Paasche, B. Engels, R. d. Dede, I. Hussain, P. Langer, M. Rettig and K. Weisz, *J. Org. Chem.*, 2009, **74**, 4878-4881; (h) N. Höttecke, H. Reinke, C. Fischer and P. Langer, *Z. Naturforsch.*, 2009, **64b**, 699-706; (i) T. S. I. Franczyk II, D. R. Hill, A. R. Haight, M. A. McLaughlin, S. Shekhar, S. Yu, J. Mei and L. Wang, WO 2009/155386(A1), 2009; (j) R. Dede, A. Riahi, M. Shkoo, M. A. Yawer, I. Hussain, N. Kelzhanova, Z. A. Abilov, A. Falodun, H. Görls and P. Langer, *Z. Naturforsch.*, 2013, **68b**, 1021-1030; (k) F. Erben, V. Specowius, J. Wölfling, G. Schneider and P. Langer, *Helv. Chim. Acta*, 2013, **96**, 924-930.
- 18 M. Horn, L. H. Schappele, G. Lang-Wittkowski, H. Mayr and A. R. Ofial, *Chem. Eur. J.*, 2013, **19**, 249-263.
- 19 See supporting information for further details.
- 20 The possible pseudopericyclic character of this process could explain its low activation barrier. For a review, see: P. v. R. Schleyer, J. I. Wu, F. P. Cossio and I. Fernandez, *Chem. Soc. Rev.*, 2014, **43**, 4909-4921.
- 21 (E/Z) isomerization of silyl enol ether has already been documented. See for example: J. L. Duffy, T. P. Yoon and D. A. Evans, *Tetrahedron Lett.*, 1995, **36**, 9245-9248.
- 22 The possible biradical character of TS_{BC} has been evaluated via a UB3LYP calculation and found unfavorable (+43.1 kJ.mol⁻¹).
- 23 A small amount (5-10%) of the corresponding β -ketonitriles was also isolated from these reactions.
- 24 R. Dolmazon, *J. Heterocycl. Chem.*, 1988, **25**, 751-757 and related articles from the same group.
- 25 (a) L. Capuano, R. Zander and P. Zenner, *Chem. Ber.*, 1979, **112**, 3753-3758; (b) L. Capuano and T. Tammer, *Chem. Ber.*, 1981, **114**, 456-467.