

View Article Online View Journal

# **RSC Advances**

This article can be cited before page numbers have been issued, to do this please use: S. Jadhav, A. Jagdale, S. Kamble, R. S. Salunkhe and A. Kumbhar, *RSC Adv.*, 2015, DOI: 10.1039/C5RA20680D.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This Accepted Manuscript will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/advances

## Palladium nanoparticles supported on titanium dioxide cellulose composite (PdNPs@TiO<sub>2</sub>-Cell) for ligand-free carbon-carbon cross coupling reactions

Sanjay Jadhav<sup>a</sup>, Ashutosh Jagdale<sup>b</sup>, Santosh Kamble<sup>c</sup>, Arjun Kumbhar<sup>b\*</sup>, Rajshri Salunkhe<sup>a\*</sup>

<sup>a</sup>Department of Chemistry, Shivaji University, Kolhapur, 416004, M.S., India

<sup>b</sup>Department of Chemistry, Padmabhushan Dr. Vasantraodada Patil College, Tasgaon, (Sangli) Maharashtra, 416312, India

<sup>c</sup>Department of Chemistry, Yashvantrao Chavan Institute of Science, Satara, Maharashtra, 415001, India

E-mail: arjun2win@yahoo.co.in

**Abstract**: Well-dispersed non-spherical PdNPs with a diameter of 39-45 nm supported on TiO<sub>2</sub>-cellulose composite (PdNPs@TiO<sub>2</sub>-Cell) can be synthesized by a simple and clean route. The catalyst was well characterized by XRD, FE-SEM, EDS, and TEM techniques. The PdNPs have good dispersity on the particles of support. This results in excellent catalytic activities for the synthesis of biphenyls, acrylates, acetylenes and prochiral ketones using low Pd loading (1 mol %) at comparatively low temperature. The effects of nature and amount of bases, nature of solvents, amount of catalyst and the reaction temperature on the activity of PdNPs@TiO<sub>2</sub>-Cell were thoroughly investigated. The catalyst showed at least four times reusability without decrease in catalytic activity.

**Keywords:** Cellulose, Pd nanoparticles, biopolymeric-inorganic hybrid material, TiO<sub>2</sub>, crosscoupling reactions.

#### 1. Introduction

Transition metal catalysis especially palladium catalyzed cross coupling reactions of aromatic halides in the presence of various nucleophiles is strategically important in organic synthesis. It has been widely used for the synthesis of a diverse array of biphenyls, acrylates, acetylenes and prochiral ketones by C-C cross coupling reactions. These compounds have profound importance in chemical, pharmaceutical and biochemical industries<sup>1</sup>. Additionally, such compounds are also present in many natural as well as biologically active compounds<sup>2</sup>, and are especially interesting in applications for organic light-emitting diodes and chemiluminescence detection systems<sup>3</sup>. These compounds have been mostly synthesized by palladium catalyzed Suzuki-Miyaura<sup>4</sup>, Mizoroki-Heck<sup>5</sup>, Heck-Matsuda<sup>6</sup>, Sonogashira-Hagihara<sup>7</sup>, and carbonylative cross-coupling reactions<sup>8</sup>. Recently, this area of research has attracted great interest because of its high compatibility to a wide variety of functional groups under mild reaction conditions.

Though, most of these transformations have been extensively investigated by homogeneous palladium complexes in solution<sup>9</sup>. The separation of metal catalysts from the reaction mixture and their reuse is highly desirable from economical and environmentally point of view<sup>10</sup>. Additionally, the homogeneous Pd complexes also undergo deactivation due to the aggregation of Pd during the reactions. In this context, heterogeneous catalysts particularly, the PdNPs supported on suitable solid support has found immense importance for many cross coupling reactions<sup>11</sup>. This strategy increases the catalytic activity of Pd and also reduces the amount of metal required for the reaction<sup>12</sup>. Several oxides have been used as a support for PdNPs<sup>13</sup>, because moderate to high dispersions was obtained on these oxides due to favorable metal–support interactions<sup>14</sup>. Out of these oxides TiO<sub>2</sub> based materials have found potential applications across many different areas<sup>15</sup>. In recent years much like the noble metal nanoparticles, PdNPs supported TiO<sub>2</sub> and Pd supported TiO<sub>2</sub> core shell catalysts have seen an extensive amount of research in methanol reforming<sup>16</sup>, hydrogenation<sup>17</sup>, and photocatalysis<sup>18</sup>.

Biopolymers such as alginate, chitosan, starch, and cellulose has been developed as a most attractive support for immobilization of many Pd catalysts<sup>19</sup>. The extensive number of -OH groups present in cellulose can facilitate the complexation of TiO<sub>2</sub> to the molecular matrix, and play a significant role in guiding the organization of TiO<sub>2</sub> among cellulose molecules. In addition to this, the use of cellulose has several key advantages. No additional reducing agents are required<sup>20</sup>. Cellulose also avoids the aggregations of PdNPs, as it acts as the protecting agent similar to other biopolymers<sup>21</sup>. There is binding interaction between cellulose and the metal nanoparticles which provides a platform to PdNPs and helps to stabilize Pd as that of hydroxyl group containing solvents<sup>22-24</sup>. As a result of this there is very less leaching of metal into solution. In addition it is possible to carry out reactions in aqueous medium due to the fine dispersibility of cellulose in water<sup>25</sup>. Hybrid organic-inorganic composites have received much attention during recent years because of their unique properties<sup>26</sup>. Due to these properties there is easy diffusion of reactants and reagents inside the pores of materials. They can be modified in the form of hybrid composites by easy processing with conventional techniques with good thermal and chemical stability.

Recently we showed that Pd supported on alumina cellulose composite (Pd@Al<sub>2</sub>O<sub>3</sub>-Cell catalyst) yields various biphenyls in good to excellent yields in aqueous medium<sup>27</sup>. In continuation of our efforts to use natural feedstock such as carbon<sup>28</sup>, agrose<sup>29</sup> and chitosan<sup>30</sup> for immobilization of Pd catalyst, herein we report an efficient method for the synthesis of PdNPs dispersed on TiO<sub>2</sub> cellulose composite (PdNPs@TiO<sub>2</sub>-Cell). The prepared catalyst was well characterized by different techniques and applied for the synthesis of various biphenyls, acrylates, prochiral ketones and acetylenes using low Pd loading (1 mol %) at comparatively low temperatures.

#### 2. Results and discussion

#### 2.1 The preparation of PdNPs@TiO<sub>2</sub>-Cell catalyst

The preparation of PdNPs@TiO<sub>2</sub>-Cell catalyst is quite straightforward. TiO<sub>2</sub>-Cell composite<sup>31</sup> obtained from cellulose and TiCl<sub>4</sub> was stirred with Pd(OAc)<sub>2</sub> in ethanol at 50 °C temperature as described in the experimental procedure affording the immobilized PdNPs on TiO<sub>2</sub>-Cell composite. The composite turned light gray after 4 h indicating the formation of PdNPs.

#### 2.2 The characterization of PdNPs@TiO2-Cell catalyst

The amount of Pd in the catalyst was 2.9 wt % as given by ICP-AES. The field emission scanning electron microscopy (FE-SEM) images of PdNPs@TiO<sub>2</sub>-Cell is shown in Fig. 1a and 1b. The SEM image shows the fibrous morphology of the catalyst. In EDS images (Fig.1c and 1d) the bright points on the surface are due to the characteristic emission of the Ti. The EDS metal mapping shows that  $TiO_2$  was uniformly covered on the fibrous surface of the cellulose. EDS spectrum (Fig. 1e) also shows that palladium is exists in the composite along with titanium, oxygen. This also confirmed the successful grafting of the Pd and  $TiO_2$  on cellulose.



**RSC Advances Accepted Manuscript** 

Fig. 1. FE-SEM images of PdNPs@TiO<sub>2</sub>-Cell, SEM images; (a, b), back scattering images (c, d), and EDS spectrum (e).

The TEM micrographs of the catalyst are depicted in Fig. 2. It can be seen that nonspherical shaped PdNPs of 39-45 nm were homogeneously dispersed on composite. As shown in figure, the PdNPs are non-spherical and contain a large number of sharp corners, and edge sites. The surface atoms at these sites are physically unstable and are chemically active.



Fig. 2. TEM analysis of PdNPs@TiO<sub>2</sub>-Cell (a) fresh catalyst, and (b) catalyst after 4<sup>th</sup> recycle.

The XRD analysis of cellulose and catalyst (Fig. 3) shows that the coating of oxide is exclusively in amorphous. This was indicated by no other diffraction peaks in XRD spectrum of the catalyst. The X-ray diffraction pattern also confirmed the formation of highly crystalline PdNPs.



Fig. 3. XRD pattern of cellulose and PdNPs@TiO2-Cell

#### 2.3 The activity of PdNPs@TiO<sub>2</sub>-Cell catalyst for Suzuki-Miyaura coupling reaction:

To test the activity of catalyst and effect of various parameters, we first carried out Suzuki-Miyaura cross-coupling reaction between 4-bromobenzophenone and phenylboronic acid (Scheme 1).



Scheme 1. Suzuki-Miyaura reaction catalyzed by PdNPs@TiO2-Cell

To examine the effect of bases, a series of bases were taken into consideration for the coupling between 4-bromobenzophenone and phenylboronic acid in 95 % ethanol at 80°C temperature (Table 1) using 1 mol % catalyst. The most common and inexpensive base K<sub>2</sub>CO<sub>3</sub> was found to be most effective (Table 1, entry 7). Surprisingly, K<sub>3</sub>PO<sub>4</sub>, NaOAc, Na<sub>2</sub>CO<sub>3</sub>, and Cs<sub>2</sub>CO<sub>3</sub> were found to be less effective (Table 1, entries 1-4). Though organic base Et<sub>3</sub>N showed good conversion (Table 1, entry 6), pyridine was completely failed to convert 4-bromobenzophenone into corresponding coupling product (Table 1, entry 5). We then studied the effect of quantity of base by carrying reactions using 0.5 to 3 equivalents of K<sub>2</sub>CO<sub>3</sub> (Table 1, entries 7-10). When the reaction is carried out without K<sub>2</sub>CO<sub>3</sub>, no product was observed indicating that the base is known to activate<sup>32</sup> phenylboronic acid (Table 1, entry 11). However, our results indicated that for this catalytic system base is not needed for the activation of phenylboronic acid, and its only role is to neutralize the boric acid<sup>33</sup> as one equivalent base showed good conversion (90 % yield) in 1 h (Table 1, entry 8). Thus, either a surface hydroxyl group of TiO<sub>2</sub> hydrolyzes the boronic acid to boric acid as reported for gold<sup>34</sup> and/or to the cerium oxide<sup>35</sup>.

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.

Entry	Base (mmol)	Time (h)	Yield <sup>b</sup> (%)
1	K <sub>3</sub> PO <sub>4</sub> (2)	6	74 (55)
2	NaOAc (2)	5	70(50)
3	$Na_2CO_3(2)$	5	82(70)
4	$Cs_2CO_3$ (2)	6	80 (65)

**Table 1.**Screening of various bases for Suzuki–Miyaura coupling reaction<sup>a</sup>

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.

5	Pyridine (2)	10	Trace
6	Et <sub>3</sub> N (2)	7	85(75)
7	$K_2CO_3(2)$	5	90 (90)
8	$K_2CO_3(1)$	5	90(90)
9	K <sub>2</sub> CO <sub>3</sub> (0.5)	6	50(50)
10	K <sub>2</sub> CO <sub>3</sub> (3)	6	90(90)
11	Base free	12	no reaction

<sup>a</sup>Reaction conditions: 4-bromobenzophenone (1.0 mmol), phenyl boronic acid (1.1 mmol), catalyst (1 mol %), base (0 to 3 mmol), 95 % ethanol (5 mL), at 80 °C. Values in the parentheses are yields after 1 h. <sup>b</sup>Isolated yields after column chromatography.

The screening of the solvents for model reaction showed that catalyst gave quantitative yield (90 %) in 95 % ethanol within 1 h (Table 2, entry 1). Unfortunately catalysts was less active in water (Table 2, entry 4), but showed moderate activity in EtOH:H<sub>2</sub>O mixture (Table 2, entries 2 and 3). We then examined the activity of catalyst in DMF and DMF:H<sub>2</sub>O mixture. The results showed that PdNPs@TiO<sub>2</sub>-Cell was less active as compared to our previous reported Pd@Al<sub>2</sub>O<sub>3</sub>-Cell catalyst<sup>27</sup> in DMF and DMF:H<sub>2</sub>O mixture. Use of 95 % ethanol renders this protocol quite practical to large scale synthesis considering its economic and environmental point of view.

Entry	Solvent	Time (h)	Yield <sup>b</sup> (%)
1	95% EtOH	1	90
2	EtOH:H <sub>2</sub> O (80:20)	6	70
3	EtOH:H <sub>2</sub> O (50:50)	6	40
4	H <sub>2</sub> O	6	25
5	DMF	6	80
6	DMF:H <sub>2</sub> O (80:20)	6	60
7	DMF:H <sub>2</sub> O (50:50)	6	34

Table 2. Effect of solvents on Suzuki–Miyaura coupling reaction<sup>a</sup>

<sup>a</sup>Reaction conditions: 4-bromobenzophenone (1.0 mmol), phenyl boronic acid (1.1 mmol), catalyst (1 mol %),  $K_2CO_3$  (1.0 mmol), solvent (5 mL) at 80 °C. <sup>b</sup>Isolated yields after column chromatography.

In order to evaluate the potential of the catalyst, the model reaction was run with changing the amount of catalyst. Lowering the palladium concentration might effectively suppress agglomeration of Pd as observed in ligand-free Suzuki–Miyaura coupling reaction by Pd supported zeolite<sup>36</sup>. The results shown in Table 3 indicated that among the different loading 1 mol% proved to be the best and was chosen as the optimum loading for further study.

Entry	Amount of catalyst	Temperature	Time	Yield <sup>b</sup>
	(mol %)	(°C)	(h)	(%)
1	0.25	80	1	10
2	0.50	80	1	70
3	0.75	80	1	70
4	1.00	80	1	90
5	1.25	80	1	90
6	1.50	80	1	90
7	2.00	80	1	90
8	1.00	r.t.	12	20
9	1.00	40	12	45
10	1.00	60	12	80
11	1.00	80	12	20 <sup>c</sup>
12	1.00	80	12	15 <sup>d</sup>

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.

**Table 3.**Optimization of catalyst loading for Suzuki–Miyaura coupling reaction<sup>a</sup>

<sup>a</sup>Reaction conditions: 4-bromobenzophenone (1.0 mmol), phenyl boronic acid (1.1 mmol), catalyst (0.25 to 2 mol %),  $K_2CO_3$  (1.0 mmol), 95 % ethanol (5 mL), room temperature to 80 °C. <sup>b</sup>Isolated yields after column chromatography. <sup>c</sup>Reaction using Pd@TiO<sub>2</sub>. <sup>d</sup>Reaction using Pd@cellulose.

To study the effect of temperature on catalytic activity, the model reaction was carried out at different temperatures (Table 3, entries 8-10). The results indicated that the reaction temperature is crucial to the catalytic efficiency. Increasing the reaction temperature from room temperature to 60 °C, the yield of the product gradually increases

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.

from 20 to 80%. While, significant acceleration in rate was observed at 80 °C, affording a quantitative yield of the desired product in 1h (Table 3, entry 4). It was observed that PdNPs@TiO<sub>2</sub>-Cell is more active as compared to Pd/TiO<sub>2</sub> and Pd/Cellulose (Table 3, entries 11 and 12). This may be due to more diffusion of substrate and distribution of PdNPs over composite as compared to pure support.

To expand the substrate scope of the arylation, we performed a number of corresponding transformations using substituted aryl halides and aryl boronic acids using K<sub>2</sub>CO<sub>3</sub> as base at 80 °C in 95 % ethanol using 1 mol % catalyst. Several functional groups such as OCH<sub>3</sub>, COCH<sub>3</sub>, PhCO, and heterocyclic moieties were reacted smoothly giving the orresponding coupling products in good vields (Table 4). The position of substituents on he aromatic ring of aryl halides did not have any appreciable effect on progress of the eaction. We next turned our attention towards ortho substitution by coupling sterically indered bromomesitylene with phenyl boronic acid. In this case good yield (72 %) could e achieved in 6 h (Table 4, entry 5).The C–X bond strength affects the activity towards the uzuki-Miyaura reaction as the general cross coupling reactions follows reactivity order of -I>C-Br > C-Cl. The main drawback of the many Pd mediated Suzuki-Miyaura crossoupling reaction is that only aryl iodides and aryl bromides can be used efficiently. While ryl chlorides are either inert or gave very less conversion, because of the stronger C-Cl ond which is in fact responsible for the slower reaction rate of aryl chlorides<sup>37</sup>. nfortunately, coupling of chlorobenzene as well as 4-chlorobenzophenone with phenyl oronic acid was inefficient under optimized reaction conditions (Table 4, entries 14 and 5).

Entry	Aryl halide	Arylboronic acid	Product	Time (h)	Yield <sup>b</sup> (%)
1	O Br	B(OH) <sub>2</sub>		1	90

Table 4 The Suzuki-Miyaura	reactions of various arv	l halides with arvl horonic acids <sup>a</sup>	
<b>I ADIC T.</b> I IIC SUZUKI-MIYAUIA	I Cacholis of various arv	I Hallues with all vi bui bille actus"	

L	)	(
+	1	ŀ
ι	.1	
r	•	f
•		
ł	1	j
ł	)	(
<i>_</i>		
	)	l
(	-	
ſ		
r	•	
Ľ	-	
	1	1
2	1	1
ł	( 1 )	
	ן ז ז	
e t	ן ז ן	
	ן י י	
	) ) )	
	ַ ן ן	
	ן 1 1	

View Article Online DOI: 10.1039/C5RA20680D



Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.



<sup>a</sup>Reaction conditions: aryl halides (1.0 mmol), arylboronic acids (1.1 mmol), catalyst (1 mol %), K<sub>2</sub>CO<sub>3</sub> (1.0 mmol), 95 % ethanol (5 mL) at 80 °C. <sup>b</sup>Isolated yields after column chromatography.

## 2.4 The activity of PdNPs@TiO<sub>2</sub>-Cell catalyst for Suzuki-Miyaura coupling reactions of arenediazonium salt

There are two main problems associated with use of aryl halides in many Pd catalyzed cross-coupling reactions. Firstly, most of aryl halides are insoluble in water hence it is very difficult to carry out reactions in aqueous medium. Secondly aryl halides undergo homocoupling<sup>38</sup> many times hence separation and purification of products become tedious. One of the best ways to overcome these drawbacks is the use of diazonium salts as substitute for aryl halides as they showed excellent reactivity towards many cross coupling reactions<sup>39</sup>. The diazonium salts are generally crystalline solids and show high solubility in water. They can be easily prepared from aromatic amines by diazotization in water<sup>40</sup>. Furthermore arenediazonium salts undergo easy oxidative addition<sup>41</sup> during the

mechanistic cycle to generate active cationic aryl-Pd(II) intermediate. In addition the ligand-free palladium-catalyzed coupling reactions of diazonium salts do not undergo  $N_2$  elimination as observed in triphenylphosphine assisted cross-coupling reactions<sup>42</sup>. In this connection we used arenediazonium salts for ligand-free Suzuki-Miyaura reactions using 1 mol % catalyst,  $K_2CO_3$  as a base in water at room temperature (Table 5). Regardless of electron-rich or electron-poor substituents present on the phenyl rings, all the substrates could react smoothly to give the desired cross-coupling products with good to excellent yields (84-90 %) in 2 to 5 h.

**Table 5.**The Suzuki-Miyaura coupling reactions of various arenediazonium salts with

#### arylboronic acids<sup>a</sup>

	$N_2^+BF_4^-$	B(OH) <sub>2</sub> ↓	PdNPs@TiOColl	\	<b>`</b>
	FG +	FG -	$H_2O, K_2CO_3, r. t.$	<b>}</b> _{	⟩ ≻ <sub>FG</sub>
Entry	Arenediazoniu	Aryl boronic	e Product	Time	Yield <sup>b</sup>
	m salt	acid		(h)	(%)
1	N <sup>+</sup> BF <sub>4</sub> <sup>+</sup>	B(OH) <sub>2</sub>		2	88
2	CI N <sup>+</sup> BF <sub>4</sub> <sup>+</sup>	B(OH) <sub>2</sub>		3	90
3	CI N <sup>+</sup> BF <sub>4</sub> <sup>-</sup>	K <sup>S</sup> B(OH)		3.5	86
4	$0_2N$ $N^+BF_4^-$	B(OH) <sub>2</sub>	0 <sub>2</sub> N	2.5	90
5	O <sub>2</sub> N N <sup>+</sup> BF <sub>4</sub> <sup>+</sup>	KS B(OH)	$^{2}$ $0_{2}N$ $\sim$ $S$	3	85



<sup>a</sup>Reaction conditions: arenediazonium salts (1 mmol), aryl boronic acids (1.1 mmol), catalyst (1 mol %), K<sub>2</sub>CO<sub>3</sub> (1.0 mmol), water (5 mL) at room temperature. <sup>b</sup>Isolated yields after column chromatography.

# 2.5 The activity of PdNPs@TiO<sub>2</sub>-Cell catalyst for Mizoroki-Heck and Heck-Matsuda coupling reactions:

Encouraged by above satisfying results of Suzuki-Miyaura reaction, we studied the applicability of this catalytic system for Mizoroki-Heck and Heck-Matsuda cross-coupling reactions. Since, 1968 the Mizoroki-Heck coupling reaction<sup>43</sup> has been emerged as excellent catalytic tool for the formation of carbon–carbon bonds, since it has enormous efficiency to tolerate different functional groups<sup>44</sup>. We explored this catalytic system for Mizoroki-Heck coupling reaction of various aryl bromides with different olefins in DMF using K<sub>2</sub>CO<sub>3</sub> as a base at 100 °C. The results shown in Table 6 indicated that all substrates could smoothly react with different olefins in good to excellent yields (80-95%) in 3.5 to 9 h.

**RSC Advances Accepted Manuscript** 



#### Table 6. The Mizoroki-Heck coupling reactions of various aryl bromides with olefins<sup>a</sup>

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.



<sup>a</sup>Reaction conditions: aryl bromides (1.0 mmol), olefins (1.1 mmol), catalyst (1 mol %), K<sub>2</sub>CO<sub>3</sub> (1.0 mmol), DMF (5 mL) at 100 °C. <sup>b</sup>Isolated yields after column chromatography.

It has been known that Pd-catalyzed coupling reaction of arenediazonium salts by Heck-Matsuda reaction can easily carried out in water under base-free conditions<sup>45</sup>. The first use of arenediazonium salt for Mizoroki-Heck reaction was reported by Matsuda<sup>46</sup> and hence known as 'Heck-Matsuda reaction'. This reaction has been emerged as an alternative to aryl halides for Pd catalyzed synthesis of olefinic compounds. Considering the possible use of water as universal solvent and excellent reactivity of diazonium salts as arylating agent herein we report use of PdNPs@TiO<sub>2</sub>-Cell catalyst for base-free Heck-Matsuda reaction in water at room temperature. The results shown in Table 7 clearly demonstrates the utility of this system in Heck-Matsuda reaction in which different arenediazonium salts were coupled with varies olefin acceptors under specified reaction conditions without base.





Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31



<sup>a</sup>Reaction conditions: arenediazonium salts (1 mmol), olefins (1.1 mmol), catalyst (1 mol %), water (5 mL) at room temperature. <sup>b</sup>Isolated yields after column chromatography.

# 2.6 The activity of PdNPs@TiO<sub>2</sub>-Cell catalyst for carbonylative Suzuki-Miyaura coupling reaction:

Carbonyl compounds including prochiral ketones are of considerable interest due to their existence in a wide range of biologically active molecules<sup>47</sup>. It also acts as a promising synthetic fragment for the construction of structural core of many pharmacologically active heterocycles<sup>48</sup>. Palladium-catalyzed carbonylative cross-coupling of benzoyl chloride with aryl boronic acid is opened as alternative way to the classical Friedel–Crafts reaction<sup>49</sup> to synthesize carbonyl compounds. Thus using above catalyst range of corresponding ketone derivatives were prepared from the different benzoyl chlorides in good to excellent yields using 1 mol % catalyst, K<sub>2</sub>CO<sub>3</sub> as a base in 95 % EtOH at room temperature (Table 8).

**Table 8.** The Suzuki-Miyaura coupling reactions of various benzoyl chlorides witharylboronic acidsa



**RSC Advances Accepted Manuscript** 

DOI: 10.1039/C5RA20680D



<sup>a</sup>Reaction conditions: benzoyl chlorides (1 mmol), arylboronic acids (1.1 mmol), catalyst (1 mol %), K<sub>2</sub>CO<sub>3</sub> (1 mmol), 95 % EtOH (5 mL) at room temperature. <sup>b</sup>Isolated yields after column chromatography.

#### 2.7. The activity of PdNPs@TiO2-Cell catalyst for Sonogashira coupling reaction

Compounds containing acetylene motifs are prevalent in many important natural products and have been prepared by Sonogashira reaction. Such compounds have several important biological as well as pharmaceutical properties<sup>50</sup>. Besides these compounds,

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31.

oligomers and polymers of acetylenes have also been prepared via the Sonogashira reaction. Considering the importance of this reaction in synthesis of such important compounds<sup>51</sup>, we next attempted to perform the Cu-free Sonogashira coupling<sup>52</sup> of various aryl bromides with terminal alkynes using 1 mol % catalyst, K<sub>2</sub>CO<sub>3</sub> as a base in DMF at 80 °C. Thus we are succeeded to carry out the reactions of terminal acetylenes with number of aryl bromides under Cu-free conditions (Table 9). The results showed that the nature of the substituents did not exert significant effect on the yield and reaction time. The reactions were rapid giving good to excellent yields (85-95%) of the corresponding products in 1 to 7h.

Table 9.Cu-free Sonogashira coupling reactions of aryl bromides with terminal alkynes<sup>a</sup>



**RSC Advances Accepted Manuscrip** 



<sup>a</sup>Reaction conditions: aryl bromides (1.0 mmol), acetylenes (1.1 mmol), catalyst (1 mol %), K<sub>2</sub>CO<sub>3</sub> (1 mmol), DMF (5 ml), at 80 °C. <sup>b</sup>Isolated yields after column chromatography.

#### 2.8. Leaching and Heterogeneity

To ascertain the heterogeneous nature of the catalyst we performed hot filtration and leaching test. This study was carried out by coupling 4-bromobenzophenone and phenyl boronic acid under optimized reaction condition of Suzuki-Miyaura reaction. After 30% conversion of the aryl halide (analyzed by HPLC), the catalyst was separated at the

reaction temperature and the reaction mixture was continued to stirred at that temperature for another 2 h. There was no further increase in the product concentration, indicated heterogeneous nature of catalyst. This was further supported by Pd leaching test. Possible Pd leaching during the reaction was tested by analyzing Pd concentration in the reaction mixture by ICP-AES. It was observed that only 3 ppm of Pd was detected. The activity of the leached Pd was found to be negligible by carrying out an experiment in 3 ppm of Pd in the form of Pd(OAc)<sub>2</sub> as catalyst and P(Ph<sub>3</sub>)<sub>3</sub> ligand.

#### 2.9. Recyclability of catalyst

The feasibility of recycling the catalyst was also examined for coupling 4bromobenzophenone with phenylboronic acid. After completion of reaction, the supported catalyst was separated by centrifugation washed with ethyl acetate, water, dried under vacuum and used directly for the next cycle without further treatment. The recycling results shown in Table 10 indicated that the catalyst was recycled for at least four times without decrease in product yield. TEM analysis of reused catalysts (Fig. 2b) showed particle size of PdNPs are 45 nm this indicated that TiO<sub>2</sub>-Cell composite avoids agglomeration of PdNPs.

	Table	10.Recyclability	of catalyst in	Suzuki-Miyaura	coupling reaction
--	-------	------------------	----------------	----------------	-------------------

Cycle	Yield (%)
Fresh catalyst	90
1 <sup>st</sup> cycle	90
2 <sup>ed</sup> cycle	89
3 <sup>rd</sup> cycle	89
4 <sup>th</sup> cycle	89

#### 3. Conclusions

In summary the Pd nanoparticles dispersed on the  $TiO_2$  particles supported on cellulose was prepared by a simple procedure. The immobilization of Pd and  $TiO_2$  on

cellulose was confirmed by FE-SEM and EDS as well as XRD analyses. Furthermore the formation of PdNPs of 39-45 nm was confirmed by TEM. This methodology allows the synthesis of various substituted biphenyls and acrylates from aryl halides as well diazonium salts in environmentally benign solvents like 95% ethanol and water at low temperature in good to excellent yields. We have also demonstrated that the catalyst showed good activity for Sonogashira coupling of aryl bromides in DMF under Cu-free conditions. In almost all conversions 1 equivalent K<sub>2</sub>CO<sub>3</sub> was required. Also, efficient cross-coupling could be accomplished with benzoyl chlorides for the synthesis of ketones at room temperature. The use of this catalytic system has many advantages such as easy separation, recyclability without decrease in yield, and the reaction does not require an inert atmosphere.

#### Supporting Information.

<sup>1</sup>H and <sup>13</sup>C NMR data of representative compounds.

#### Author information

#### **Corresponding authors**

Department of Chemistry, Padmabhushan Dr. Vasantraodada Patil College, Tasgaon, 416312, M.S., India.

E-mail: arjun2win@yahoo.co.in. Tel. +91 2346 250575; Fax: +91 2346 250665.

#### Acknowledgment

This work was supported by the Science and Engineering Research Board, Department of Science and Technology (SERB-DST), Government of India, New Delhi, under the scheme of Start-Up research grants for Young Scientists (SB/FT/CS-153/2013).

#### References

- 1. K. C. Nicolaou, P. G. Bulger, and D. Sarlah, *Angew. Chem. Int. Ed.*, 2005, **44**, 4442–4489.
- 2. J. D. Sellars and P. G. Stee, *Chem. Soc. Rev.*, 2011, **40**, 5170–5180.
- 3. S. Xu, E. H. Kim, A. Wei and E.-I. Negishi, *Sci. Technol. Adv. Mater.*, 2014, **15**, 044201.
- 4. N. Miyaura, K. Yamada and A. Suzuki, *Tetrahedron Lett.*, 1979, **20**, 3437-3440.
- 5. K. Mori, T. Mizoroki and A. Osaki, Bull. Chem. Soc. Jpn. 1973, 46, 1505-1508.
- 6. J. G. Taylor, A. V. Moro, C. Roque and D. Correia, *Eur. J. Org. Chem.*, 2011, 1403–1428.

Published on 11 December 2015. Downloaded by Chinese University of Hong Kong on 12/12/2015 04:06:31

- 7. K. Sonogashira, Y. Tohda and N. Hagihara, *Tetrahedron Lett.*, 1975, **16**, 4467–4470.
- 8. Y. Li, D. Xue, C. Wang, Z.-T. Liu and J. Xiao, *Chem. Commun.*, 2012, 48, 1320-1322.
- 9. Q.-A. Chen, Z.-S. Ye, Y. Duan and Y.-G. Zhou, *Chem. Soc. Rev.*, 2013, **42**, 497-511.
- 10. B. H. Lipshutz, A. R. Abela, Z. V. Boskovic, T. Nishikata, C. Duplais and A. Krasovskiy, *Top. Catal.*, 2010, **53**, 985–990.
- 11. A. Fihri, M. Bouhrara, B. Nekoueishahraki, J.-M. Basset and V. Polshettiwar, *Chem. Soc. Rev.*, 2011, **40**, 5181-5203.
- 12. C. Deraedt and D. Aatruc, Acc. Chem. Res., 2014, 47, 494–503.
- 13. L. Djakovitch and K. Koehler, J. Am. Chem. Soc., 2001, **123**, 5990-5999.
- 14. L. M. Neal, D. Hernandez and H. E. Hagelin-Weaver, J. Mol. Catal. A, 2009, **307**, 29-36.
- 15. M. Dahl, Y. Liu and Y. Yin, *Chem. Rev.*, 2014, **114**, 9853–9889.
- L. S. Al-Mazroai, M. Bowker, P. Davies, A. Dickinson, J. Greaves, D. James and L. Millard, *Catal. Today*, 2007, **122**, 46-50.
- 17. B. Tapin, F. Epron, C. Especel, B. K. Ly, C. Pinel and M. Besson, ACS Catal. 2013, **3**, 2327-2335.
- M. Maicu, M. C. Hidalgo, G. Colón and J. A. Navío, J. Photochem. Photobiol. A, 2011, 217, 275-283..
- 19. A. Kumbhar and R. Salunkhe, *Current Organic Chemistry*, 2015, **19**, 2275-2121.
- 20. Y. Xu, L. Zhang and Y. Cui, *J. Appl. Polym. Sci.*, 2008, **110**, 2996-3000.
- 21. M. J. Gronnow, R. Luque and D. J. Macquarrie and J. H. Clark, *Green Chem.*, 2005, **7**, 552-557.
- 22. B. Zhang, Y. Yuan, K. Philippot and N. Yan, *Catal. Sci. Technol.*, 2015, 5, 1683-1692.
- 23. E. E. Zvereva, S. Grimme, S. A. Katsyuba, V. V. Ermolaev, D. A. Arkhipova, N. Yan, V. A. Miluykov, O. G. Sinyashina and A. Aleksandrove, *Phys. Chem. Chem. Phys.*, 2014, 16, 20672-20680
- 24. X. Yuan, G. Sun, H. Asakura, T. Tanaka, X. Chen, Y. Yuan, G. Laurenczy, Y. Kou, P. J. Dyson, and N. Yan, *Chem. Eur. J.*, 2013, 19, 1227–1234
- 25. F. Quignard and A. Choplin, *Chem. Commun.*, 2001, 21-22.
- 26. K. Kanamori and K. Nakanishi, *Chem. Soc. Rev.*, 2011, **40**, 754–770.
- A. Kumbhar, S. Jadhav, S. Kamble, G. Rashinkar and R. Salunkhe, *Tetrahedron Lett.*, 2013, 54, 1331–1337.

- 28. S. Jadhav, A. Kumbhar, C. Rode and R. Salunkhe, *Green Chem.*, 2015, DOI: 10.1039/c5gc02314a.
- 29. S. Jadhav, A. Kumbhar, S. Kamble, P. More and R. Salunkhe, *C. R. Chimie.*, 2013, **16**, 957-961.
- 30. S. Jadhav, A. Kumbhar and R. Salunkhe, *Appl. Organomet. Chem.*, 2015, **29**, 339-345.
- 31. L.-Z. Meng, C.-Q. Du, Y.-Y. Chen and Y.-B. Ghe, *J. Appl. Polym. Sci.* 2002, **84**, 61–66.
- 32. A. de Meijere and F. Diederich, Metal-Catalyzed Cross-Coupling Reactions, Vol. 1–2, *Wiley-VCH, Weinheim*, 2004.
- 33. S. Carrettin, J. Guzman and A. Corma, *Angew. Chem. Int. Ed.*, 2005, **44**, 2242 –2245.
- 34. E. D. Park and J. S. Lee, *J. Catal.*, 1999, **186**, 1–11.
- T. Tabakova, F. Boccuzzi, M. Manzoli and D. Andreeva, *Appl. Catal. A:Gen.*, 2003, 252, 385–397.
- A. Kumbhar, S. Kamble, A. Mane, R. Jha and R. Salunkhe, *J. Organomet. Chem.*, 2013, 738, 29-34.
- S. R. Chemler, D. Trauner and S. J. Danishefsky, *Angew. Chem. Ed. Engl.*, 2001, 40, 4544-4568.
- 38. R. B. DeVasher, L. R. Moore and K. H. Shaughnessy, *J. Org. Chem.*, 2004, **69**, 7919-7927.
- A. Roglans, A. Pla-Quintana, and M. Moreno-Manas, *Chem. Rev.*, 2006, **106**, 4622–4643.
- 40. B. S., Hannaford, A. J. Smith, P. W. G. Tatchell, A. R. Vogels Textbook of Practical Organic Chemistry, 5<sup>th</sup> ed.; *Eds.; Longman: Singapore*, 1989.
- 41. N. Taccardi, R. Paolillo, V. Gallo, P. Mastrorilli, C. F. Nobile, M. Räisänen and T. Repo, *Eur. J. Inor. Chem.*, 2007, 4645–4652.
- 42. J. P. Knowlesa and A. Whiting, Org. Biomol. Chem., 2007, 5, 31-44.
- 43. R. F. Heck, J. Am. Chem. Soc., 1968, **90**, 5518-5526.
- 44. A. B. Dounay and L. E. Overman, *Chem. Rev.*, 2003, **103**, 2945–2964.
- 45. F. Felpin, K. Miqueu, J. Sotiropoulos, E. Fouquet, O. Ibarguren and J. Laudien, *Chem. Eur. J.* **2010**, 16, 5191-5204.
- 46. K. Kikukawa, K. Nagira, F. Wada and T. Matsuda, *Tetrahedron*, **1980**, 37, 31-36.
- 47. J. Li, Y. Jiang and P.-F. Tu, *J. Nat. Prod.*, **2005**, 68, 1802-1804.

- 48. Y. Deng, Y. W. Chin, H. Chai, W. J. Keller, and A. D. Kinghorn, *J. Nat. Prod.*, **2007**, 70, 2049-2052.
- 49. M. C. Wilkinson, *Org. Lett.*, **2011**,13, 2232-2235.
- 50. D. Mujahidin and S. Doye, *Eur. J. Org. Chem.*, **2005**, 2689–2693.
- 51. R. Chinchilla and C. Najera, *Chem. Soc. Rev.*, **2011**, 40, 5084-5121.
- 52. F. Ngassa, E. Lindsey and B. Haines, *Tetrahedron*, **2009**, 65, 4085–4091.

#### Palladium nanoparticles supported on titanium dioxide cellulose composite (PdNPs@TiO<sub>2</sub>-Cell) for ligand-free carbon-carbon cross coupling reactions

Sanjay Jadhav<sup>a</sup>, Ashutosh Jagdale<sup>b</sup>, Santosh Kamble<sup>c</sup>, Arjun Kumbhar<sup>b</sup>\*, Rajshri Salunkhe<sup>a</sup>\*

<sup>a</sup>Department of Chemistry, Shivaji University, Kolhapur, 416004, M.S., India

<sup>b</sup>Department of Chemistry, Padmabhushan Dr. Vasantraodada Patil College, Tasgaon, (Sangli) Maharashtra, 416312, India

<sup>c</sup>Department of Chemistry, Yashvantrao Chavan Institute of Science, Satara, Maharashtra, 415001, India

E-mail: arjun2win@yahoo.co.in

# $R_{1}$ $R_{2}$ $R_{3}$ $R_{3$

#### **Graphical Abstract**