

NOTE

Hydrothermal Synthesis of 1D Zn(II) Coordination Polymer

BIN LI

Institute of Chemical Engineering and Environment, Wei Fang University of Science and Technology, Shandong 262700, P.R. China

Corresponding author: E-mail: whyull@yeah.net

(Received: 15 October 2012;

Accepted: 15 April 2013)

AJC-13242

One new Zn(II) compound $[Zn(L)(1,10\text{-phen})]_n [H_2L = 4,4\text{-(isopropylidene)-bis(benzoic acid)}$ and $1,10\text{-phen} = 1,10\text{-phenanthroline}]$ has been successfully synthesized. This compound shows a one-dimensional chain. The carboxylate groups in present compound adopt a *bis*-chelating coordination mode.

Key Words: Coordination polymer, Crystal structure.

The current interest in polymeric coordination networks is rapidly expanding not only for their potential applications in catalysis, molecular adsorption, magnetism, non-linear optics and molecular sensing, but also for their intriguing variety of topologies¹⁻⁴. Although the rapid progress in metal-organic frameworks has been made, it is also a great challenge to rationally prepare and control the structures and composition of target products in crystal engineering because of the difficult prediction of either the composition or the structure of the reaction product.

All reagent and solvents employed were commercially available and used as received without further purification.

Preparation of compound: A mixture of 4,4-(isopropylidene)-*bis*(benzoic acid) (1 mmol), $Zn(OAc)_2 \cdot 2H_2O$ (1 mmol), 1,10-phenanthroline (1 mmol) and distilled water (15 mL) was heated in a 25 mL stainless steel reactor with a Teflon liner 160 °C for 96 h, followed by slow cooling to room temperature. Yellow crystals of the compound formed.

X-crystallography: Suitable single crystals were selected under a polarizing microscope and fixed with epoxy cement on fine glass fibers which were mounted on a Bruker Smart 1000 CCD diffractometer with a MoK_α radiation ($\lambda = 0.71073 \text{ \AA}$) at 293(2) K. The hydrogen atoms bound to carbon were located by geometrically calculations. All non-hydrogen atoms were refined by full-matrix least-squares techniques. All calculations were performed by the SHELXTL 97 program⁵. The crystallographic data and experimental details of structural analyses for coordination polymers are summarized in Table-1.

Structure description: In compound **1**, Zn (II) atom is coordinated to four oxygen atoms from two L ligands and two nitrogen atoms from 1,10-phenanthroline ligand. In compound the carboxylate groups adopt a *bis*-chelating coordinated modes which lead to form a 1D chain structure (Fig. 1).

TABLE-1
CRYSTALLOGRAPHIC DATA AND STRUCTURE REFINEMENT SUMMARY FOR COMPLEX

Empirical formula	$C_{29}H_{22}N_2O_4 Zn$	Z, Calculated density (mg/m^3)	8,1.584
Formula weight	527.88	Absorption coefficient (mm^{-1})	1.869
Crystal system space group	Orthorhombic, Pbc _a	F(000)	2017
Unit cell dimensions	$a = 13.828(5) \text{ \AA}$	Limiting indices	$-17 \leq h \leq 17$
	$b = 10.833(5) \text{ \AA}$		$-10 \leq k \leq 14$
	$c = 28.728(5) \text{ \AA}$		$-37 \leq l \leq 37$
	$\beta = 90^\circ$		
Volume ($\text{\AA}^3$)	4303(3)	Largest diff. peak and hole ($e/\text{\AA}^3$)	0.341 and -0.727
θ Range for data collection	1.42-27.53	Goodness-of-fit on F^2	0.897
Final R indices [$I > 2 \sigma(I)$]	$R_1 = 0.0470$, $wR_2 = 0.1387$	R indices (all data)	$R_1 = 0.0804$, $wR_2 = 0.1660$




Fig. 1. Molecular structure of 1-D Zn(II) coordination polymer at 30 % probability displacement ellipsoids

REFERENCES

1. B.O. Patrick, C.L. Stevens, A. Storr and R.C.Thompson, *Polyhedron*, **24**, 2242 (2005).
2. Y.H. Wen, J.K. Cheng, Y.L. Feng, J. Zhang, Z.L. Li and Y.G. Yao, *Inorg. Chim. Acta*, **358**, 3347 (2005).
3. X.L. Wang, C. Qin, E.B. Wang and L. Xu, *J. Mol. Struct.*, **749**, 45 (2005).
4. S.R. Batten and K.S. Murray, *Coord. Chem. Rev.*, **246**, 103 (2003).
5. G.M. Sheldrick, SHELXTL97, Program for the Refinement of Crystal Structure, University of Gottingen, Germany (1997).