



Accepted Manuscript

View Article Online View Journal

This article can be cited before page numbers have been issued, to do this please use: S.-F. Yin, X. Chen, T. Chen, Y. Zhou and F. Ji, *Catal. Sci. Technol.*, 2015, DOI: 10.1039/C4CY01618A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/catalysis

Published on 14 January 2015. Downloaded by University of Huddersfield on 15/01/2015 08:14:51

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxx

ARTICLE TYPE

Iron-Catalyzed Aerobic Oxidative Functionalization of *sp*³C-H Bonds: a Versatile Strategy for the Construction of *N*-Heterocycles

Xiuling Chen,[§] Tieqiao Chen,[§] Fangyan Ji, Yongbo Zhou,* and Shuang-Feng Yin*

Received (in XXX, XXX) Xth XXXXXXXX 200X, Accepted Xth XXXXXXXX 200X 5 DOI: 10.1039/b000000x

An iron-catalyzed aerobic oxidative functionalization of $sp^{3}C$ -H bonds has been developed for the construction of *N*heterocycles from easily available carboxylic acid derivatives and *o*-substituted anilines. This transformation represents a ¹⁰ widely applicable protocol to *N*-heterocycles using biofriendly iron as catalyst in combination with molecular oxygen or air as the sole oxidant.

The application of iron salts as catalysts in organic synthesis has attracted much attention due to their abundance, low price, ¹⁵ less toxic and biofriendly properties.¹ For example, iron salts have been utilized extensively to promote the traditional cross-coupling,² Friedel–Crafts benzylation,³ carbonylation⁴ and other processes.⁵⁻⁸ Iron-catalyzed oxidations for C-H bonds have also been extensively developed⁶⁻⁸ with Gif ²⁰ chemistry⁶ and Fenton chemistry being the most famous.⁷ However, these oxidations have commonly depended on hazardous peroxides. Nowadays, the improvement emphasizes on the development of synthetic models, using a "green" oxidant such as readily available and nontoxic O_2 or air.⁹ 25 Therefore, the iron-catalyzed aerobic oxidation of C-H bonds is highly desired. In 2012, Maes et al realized iron-catalyzed aerobic oxidation of benzylic C-H bonds in diarylmethanes for the preparation of diarylketones.¹⁰ Subsequently, Kappe et al developed gas-liquid continuous-flow technology to improve 30 Maes' iron-catalyzed aerobic oxidation system.¹¹ Despite these advances, further Iron-catalyzed aerobic oxidative functionalization of sp³C-H bonds is rare.¹² Currently, the decarboxylation reactions are also emerging as the powerful methodology for the construction of carbon-carbon bonds and 35 carbon-heteroatom bonds in organic synthesis due to the readily available substrates, simple operation and clean byproduct (only CO₂ as the byproduct).^{13,14} Herein, we communicate an efficient strategy of iron-catalyzed aerobic

- oxidative functionalization of sp^{3} C-H bonds to construct *N*-40 heterocycles from easily accessible carboxylic acid derivatives and *o*-substituted anilines (eq 1). In the present catalytic system, we achieved aerobic oxidation of sp^{3} C-H bonds, decarboxylation and oxidative cyclization in one pot. Compared to conventional methods, this procedure is 45 distinguished by using biofriendly iron catalyst in
- combination of clean dioxygen oxidant. This new method provides a general and environmentally friendly access to *N*-

heterocyclic compounds which are ubiquitous core units of various biologically active drugs and natural products.^{15,16}



o-Aminobenzamide 1a and phenylacetic acid 2a were chosen as the model substrates for optimization of the present iron-catalyzed aerobic oxidative functionalization of sp³C-H 55 bonds to construct N-heterocycles and the results are compiled in Table 1. Preliminary screening shows that various iron catalysts can catalyze the reaction alone without any ligands, bases or additives, and FeCl₃ shows the highest catalytic efficiency among the examined iron catalysts (Table 1, entries 60 1-7). In the presence of 10 mol% FeCl₃, this tandem reaction takes place smoothly at 100 °C in DMF to produce the corresponding 2-phenyl quinazolin-4(3H)-one compound 3a in 92% yield (Table 1, entry 7). Worth noting is that iron catalyst is essential for the current catalytic system. In the 65 absence of iron complex, this reaction cannot work and no product is detectable at all (Table 1, entry 8). Other metal complexes such as PdCl₂, NiCl₂.6H₂O and CuCl₂ (Note: Cu salts are used as good catalyst for the oxidative decarboxylation of the sp³C-H bonds^{9e-9g}) cannot mediate the 70 present reaction (Table 1, entries 9-11). The solvents also play an important role, and only DMF serves as the good solvent (Table 1, entries 12-15). Under the similar reaction conditions, replacement of dioxygen with other oxidants such as TBHP, DTBP, mCPBA and K₂S₂O₈ results in no yield of the desired 75 product (Table 1, entries 16-17). This transformation also depends on the reaction temperature. For example, catalyzed by 10 mol% FeCl₃, the reaction doesn't occur at 80 °C (Table 1, entry 18), while the yield dramatically increases to 92% at 100 °C (Table 1, entry 7). However, further increase of 80 reaction temperature to 120 °C leads to no evident improvement on product yield (Table 1, entry 19).



Table 1. Optimization of the reaction conditions

^aReaction conditions: 1a (0.2 mmol), 2a (0.22 mmol), catalyst (0.02 mmol), sovent (1 mL), O₂ (1 atm) in a Schlenk tube (10 mL) at 100 °C, 12 h, recharging dioxygen after 6 h. ^bGC yields based on 1a using dodecane as an internal standard. ^c0.5 mmol *m*-CPBA (*m*-chloroperbenzoic acid) or K₂S₂O₈ was employed as oxidant under N₂ (1 atm) atmosphere. ^d0.5 mmol
 ¹⁰ TBHP (*tert*-butylhydroperoxide) or DTBP (di-*tert*-butyl peroxide) was used as oxidant under N₂ (1 atm) atmosphere. ^e80 °C. ^f120 °C.

Published on 14 January 2015. Downloaded by University of Huddersfield on 15/01/2015 08:14:51

This iron-catalyzed aerobic oxidative functionalization of sp^{3} C-H bonds can be successfully applied to other substrates, showing that this reaction is a general method for the ¹⁵ preparation of *N*-heterocyclic compounds, i.e. quinazolin-4(3*H*)-ones (Table 2), quinazolines (Table 3), benzimidazoles and benzothiazoles (Table 4). As shown in Table 2, *o*-aminobenzamide **1a** reacts readily with different kinds of arylacetic acids bearing both electron-donating groups and ²⁰ electron-withdrawing groups on the aromatic cycles to give

- ²⁰ electron-withdrawing groups on the aromatic cycles to give the corresponding quinazolin-4(3*H*)-ones. Various valuable functional groups such as OMe **2b**, OH **2c**, NH₂ **2d**, NO₂ **2e**, Br **2f** and Cl **2g** all survive in the present catalytic system and the expected products are produced in high yields. Especially,
- ²⁵ the iodine substituted phenylacetic acid also reacts readily with *o*-aminobenzamide **1a** under similar reaction conditions, generating the desired product **3h** in 68% yield. Catalyzed by 10 mol% FeCl₃, both 1-naphthylacetic acid and 2naphthylacetic acid undergo aerobic oxidation tandem
- 30 reaction with 1a and the corresponding products 3i and 3j are afforded in 82% and 85% yields, respectively. Furthermore, different kinds of heterocycles can be introduced into





^a Reaction conditions: **1a-1d** (0.2 mmol), **2a-2o** (0.22 mmol), FeCl₃ (0.02 mmol, 10 mol%), DMF (1 mL), O₂ (1 atm) in a Schlenk tube (10 mL) at 100 °C, 12 h, recharging dioxygen after 6 h. ^bChlorobenzene as solvent, 3 equivs propionic acid, 140 °C, 12 h, recharging dioxygen after 40 6 h.

quinazolin-4(3*H*)-one molecule using the current aerobic oxidative functionalization of *sp*³C-H bonds. Under dioxygen atmosphere, the reaction of 2-thiopheneacetic acid and **1a** takes place smoothly in the presence of 10 mol% ⁴⁵ FeCl₃ at 100 °C in DMF, furnishing the expected product **3k** in 80% yield. 3-Pyridylacetic acid **2l** is also an efficient Published on 14 January 2015. Downloaded by University of Huddersfield on 15/01/2015 08:14:51

substrate and can be converted to **31** in 80% yield. 2-(1Hindol-3-yl)quinazolin-4(3*H*)-one **3m** bearing indolyl has been synthesized from the reaction of 2-(1H-indol-3yl)acetic acid and **1a**. In addition to *o*-aminobenzamide **1a**, substituted *s o*-aminobenzamides also serve as good substrates and react with arylacetic acids to afford the corresponding quinazolin-4(3*H*)-one derivatives **3n-3p**. Using ethyl 2-phenylacetate **2n** instead of 2-phenylacetic acid, the tandem reaction proceeds smoothly, giving the desired product **3a** in 69% ¹⁰ yield. In addition, 2-methylquinazolin-4(3*H*)-one **3q** is obtained in 60% yield from propionic acid **2o** and *o*aminobenzamide.





^aReaction conditions: **1e** (0.2 mmol), **2a-2k** (0.22 mmol), FeCl₃ (0.02 mmol, 10 mol%), DMF (1 mL), air (1 atm) in a Schlenk tube (25 mL) at 100 $^{\circ}$ C, 18 h, recharging air after 6 h.

To our delight, quinazoline derivatives **4** have been efficiently ²⁰ synthesized using the current aerobic oxidative functionalization of sp^{3} C-H bonds (Table 3). It is noted that the reaction of 2-aminobenzylamine with arylacetic acid can occur smoothly under air atmosphere catalyzed by 10 mol% FeCl₃ at 100 °C in DMF to give the expected product **4a** in 90% yield. As shown in Table 3,

- $_{25}$ similar to the substrate scope for the synthesis of quinazolin-4(3*H*)-one derivatives, arylacetic acids holding both electrondonating groups and electron-withdrawing groups on the aromatic cycles work well and react readily with 2aminobenzylamine to give the corresponding quinazolines in high
- ³⁰ yields. Noteworthy, valuable functional groups such as OMe, OH, NH₂, NO₂, Br and Cl all are compatible under the present reaction conditions. The heterocycle exemplified by **4i** having thiophenyl has also been introduced into quinazoline framework by this strategy.
- ³⁵ Under the optimal reaction conditions, the protocol can be applied to preparation of the bioactive five-membered

compounds like benzimidazoles and benzothiazoles (Table 4). Benzothiazole **5a** is obtained from similar reaction of *o*aminothiophenol **1f** with **2b**. Other substituted phenylacetic ⁴⁰ acid also serves as good substrates, affording the corresponding benzothiazoles **5b-5d** in high yields. Besides, bioactive benzimidazoles **5e** and **5f** have been prepared from the reaction of *o*-phenylenediamine **1g** with phenylacetic acid and 2-(thiophen-2-yl)aceticacid under similar reaction ⁴⁵ conditions and the yields are 80% and 81%, respectively.

Table 4. Fe-catalyzed aerobic oxidative amidation of $sp^{3}C$ -H bonds for the synthesis benzimidazoles and benzothiazoles^{*a*}



^{*a*}Reaction conditions: **1f-1g** (0.2 mmol), **2a-2i** (0.22 mmol), FeCl₃ (0.02 mmol, 10 mol%), DMF (1 mL), O₂ (1 atm) in a Schlenk tube (10 mL) at 100 $^{\circ}$ C, 12 h, recharging dioxygen after 6 h.

To clarify the reaction mechanism, several control 55 experiments were performed. In the nitrogen atmosphere, phenylacetic acid 2a is recovered completely and no decarboxylative products are detectable under similar condition (eq 2). Thus it is reasonable that the aerobic oxidation of sp³C-H bonds takes place prior to ⁶⁰ decarboxylation.¹⁷ In addition, α -hydroxyphenylacetic acid **2p** reacts with o-aminobenzamide 1a smoothly under the optimal condition to give the corresponding product 3a in 88% yield (eq 3). When 2-oxo-2-phenylacetic acid 2q is used as substrate without the aid of anhydrous FeCl₃, the reaction with 65 1a takes place to give 3a in 81% yield (eq 4). While in the absence of 1a, benzaldehyde is generated in 99% yield from the iron-catalyzed aerobic oxidation of phenylacetic acid, and the resulting benzaldehyde can further react with 1a to furnish **3a** in 97% yield (eq 5). Therefore, it is speculated that α -70 hydroxyphenylacetic acid, 2-oxo-2-phenylacetic acid and benzaldehyde probably serve as the efficient intermediates in the current iron-catalyzed aerobic oxidation-decarboxylationcyclization system. When phenylacetic acid 2a was replaced by 1,2-propanediol **2r**, the 2-methylquinazolin-4(3*H*)-one **3q** was obtained in 75% yield (eq 6). Moreover, when the radical trapper BHT (2,6-di-tert-butyl-4-methylphenol) is loaded, the reaction of **1a** with **2a** does not occur and no product **3a** or ⁵ benzaldehyde can be detected, indicating that this iron-catalyzed sp^{3} C-H bonds aerobic oxidation and decarboxylation of arylacetic acids proceeds via a radical process (eq 7).



¹⁰ On the basis of results described above and in literatures, 9,10,16,18 a plausible reaction process for the present iron-catalyzed aerobic oxidative functionalization of sp^{3} C-H bonds is proposed as shown in Scheme 1. In the Fe/O₂ system, the C-H bonds are first oxidized to form α -hydroxycarboxylic ¹⁵ acid via a radical path, followed by dehydrogenation to give

2-oxo-2-carboxylic acid.^{9,10,18} Subsequently, the resulting 2-







20 oxo-2-carboxylicacetic acid undergoes decarboxylation

generating aldehyde, which is captured by *o*-substituted anilines to produce *N*-heterocyclic compounds under oxygen atmosphere.¹⁶

In summary, we have developed an efficient strategy of 25 iron-catalyzed sp³C-H bonds aerobic oxidative functionalization to construct N-heterocycles. The transformation occurs via a tandem sequence involving ironcatalyzed *sp*³C-H bonds aerobic oxidation, decarboxylation and subsequent oxidation cyclization. This method also 30 provides an environmentally friendly protocol to Nheterocyclic compounds from easily accessible carboxylic acid derivatives and o-substituted anilines with wide substrate scope.

Acknowledgements

³⁵ This work was supported by the NSFC (U1162109, 21273066, 21172062, 21273067), Program for Changjiang Scholars and Innovative Research Team in University (IRT1238), and the Fundamental Research Funds for the Central Universities (Hunan University).

40 Notes and references

- [§]These authors contributed equally to this work.
- * State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Chemistry and Chemical Engineering, Hunan University, Changsha, 410081, P. R. China. Fax: (+) 86-731-88821171.; E-mail: 45 zhouyb@hnu.edu.cn; sf_yin@hnu.edu.cn

^{*} Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

- For some reviews: a) C. Bolm, J. Legros, J. Le Paih and L. Zani, 1 Chem. Rev. 2004, 104, 6217-6254; b) C. Sun, B. Li and Z. Shi, 50 Chem. Rev. 2011, 111, 1293-1314; c) F. Jia and Z. Li, Org. Chem. Front. 2014, 1, 194-214; d) J. E. M. N. Klein and B. Plietker, Org. Biomol. Chem. 2013, 11, 1271-1279; e) K. C Majumdar, N. De, T. Roy, Ghosh and B. Tetrahedron. 2014. doi: 10.1016/j.tet.2014.04.025. 55
- a) R. Chinchilla and C. Nájera, *Chem. Rev.* 2007, **107**, 874-922; b)
 B. D. Sherry and A. Fürstner, *Acc. Chem. Res.* 2008, **41**, 1500-1511;
 c) W. Liu, H. Cao and A. Lei, *Angew. Chem. Int. Ed.* 2010, **49**, 2004-2008; d) J. P. Knowles and A.Whiting, *Org. Biomol. Chem.* 2007, **5**, 31-44 and references therein.
- a) M. Bandini, M. Tragni and A. Umani-Ronchi, *Adv. Synth. Catal.* 2009, **351**, 2521-2524; b) I. Iovel, K. Mertins, J. Kischel, A. Zapf
 and M. Beller, 3981-3985; *Angew. Chem. Int. Ed.* 2005, **44**, 3913-3917.
- 65 4 K. M. Driller, H. Klein, R. Jackstell and M. Beller, Angew. Chem. Int. Ed. 2009, 48, 6041-6044.
- 5 a) W. Han and A. R. Ofial, *Chem. Commun.* 2009, 5024-5026; b) S. Prateeptongkum, I. Jovel, R. Jackstell, N. Vogl, C. Weckbecker and M. Beller, *Chem. Commun.* 2009, 1990-1992; c) K. Kohno, K.
- Nakagawa, T. Yahagi, J. C. Choi, H. Yasuda and T. Sakakura, J. Am. Chem. Soc. 2009, 131, 2784-2785; d) J. Kischel, I. Jovel, K. Mertins, A. Zapf and M. Beller, Org. Lett. 2006, 8, 19-22; e) P. Spannring, I. Prat, M. Costas, M. Lutz, P. C. A. Bruijnincx, B. M. Weckhuysend and R. J. M. Klein Gebbink, Catal. Sci. Technol. 2014, 4, 708–716;
 f) Y. Song, X. Tang, X. Hou and Y. Bai, Chin. J. Org. Chem. 2013, 33, 76-89 and references therein.
 - 6 M. J. Perkins, Chem. Soc. Rev. 1996, 25, 229-236.
- 7 S. Goldstein, and D. Meyerstein, Acc. Chem. Res. 1999, 32, 547-550.
- 8 a) W. Nam, Acc. Chem. Res. 2007, 40, 522-531; b) S. Shaik, H.
- Hirao and D. Kumar, Acc. Chem. Res. 2007, 40, 532-542; c) Z. Li, L.
 Cao and C. Li, Angew. Chem. Int. Ed. 2007, 46, 6505-6507; d) Y.

70

Zhang and C. Li, *Eur. J. Org. Chem.* 2007, 4654-4657; e) C. Song, G. Cai, T. R. Farrell, Z. Jiang, H. Li, L. Gan and Z. Shi, *Chem. Commun.* 2009, 6002-6004; f) H. Li, Z. He, X. Guo, W. Li, X. Zhao and Z. Li, *Org. Lett.* 2009, **11**, 4176-4179; g) Z. Wang, Y. Zhang, H.

- Fu, Y. Jiang and Y. Zhao, Org. Lett. 2008, 10, 1863-1866; h) F. Szabó, B. Pethő, Z. Gonda and Z. Novák, RSC Adv. 2013, 3, 4903-4908; i) T. Wang, W. Zhou, H. Yin, J. Ma and N. Jiao, Angew. Chem. Int. Ed. 2012, 51, 10823-10826; j) S. Grupe and A. Jacobi von Wangelin, ChemCatChem, 2013, 5, 706-710; k) S. Iwata, T. Hatt and H. Harb, Adv. Surf. Control 2012, 254, 2480, 2480, 4484, h) M.
- Hata and H. Urabe, *Adv. Synth. Catal.* 2012, **354**, 3480-3484; 1) M.
 Nakanishi and C. Bolm, *Adv. Synth. Catal.* 2007, **34**9, 861-864.
- 9 a) Z. Shi, C. Zhang, C. Tang and N. Jiao, *Chem. Soc. Rev.* 2012, 41, 3381-3430; b) W. Wu and H. Jiang, *Acc. Chem. Res.* 2012, 45, 1736-1748; c) S. E. Allen, R. R. Walvoord, R. Padilla-Salinas and Aller All
- M. C. Kozlowski, *Chem. Rev.* 2013, **113**, 6234-6458; d) H. Guo, A. Al-Hunaiti, M. Kemell, S. Rautiainen, M. Leskelä and T. Repo, *ChemCatChem*, 2011, **3**, 1872-1875; e) Q. Song, Q. Feng, and K. Yang, *Org. Lett.* 2014, **16**, 624-627; f) Q. Feng and Q. Song, *J. Org. Chem.* 2014, **79**, 1867-1871; g) Q. Feng and Q. Song, *Adv. Synth. Catal.* 2014, **356**, 1697-1702.
- 10 J. D. Houwer, K. A. Tehrani and B. U. W. Maes, *Angew.Chem. Int. Ed.* 2012, **51**, 2745 -2748.
- 11 B. Pieber and C. O. Kappe, Green Chem. 2013, 15, 320-324.
- 12 A. Gonzalez-de-Castro, C. M. Robertson, and J. Xiao, *J. Am. Chem.* 25 Soc. 2014, **136**, 8350–8360.
- a) L. J. Goossen and K. Goossen, in *Topics in Organometallic Chemistry Vol.* 44 (Ed.: L. J. Goossen), Springer, Heidelberg, 2013, pp. 121–141; b) N. Rodr guez and L. J. Goossen, *Chem. Soc. Rev.* 2011, 40, 5030-5048; b) S. M. Bonesi and M. Fagnoni, *Chem. Eur. J.* 2010, 16, 13572-13589.
- a) H. Bi, W. Chen, Y. Liang and C. Li, *Org. Lett.* 2009, **11**, 3246-3249; b) H. Yang, H. Yan, P. Sun, Y. Zhu, L. Lu, D. Liu, G. Rong and J. Mao, *Green Chem.* 2013, **15**, 976-981; c) K. Yang, C. Zhang, P. Wang, Y. Zhang and H. Ge, *Chem. Eur. J.* 2014, **20**, 7241-7244.
- ³⁵ 15 a) R. W. DeSimone, K. S. Currie, S. A. Mitchell, J. W. Darrow and D. A. Pippin, *Comb. Chem. High Throughput Screen*. 2004, **7**, 473-493; b) P. D. Leeson and B. Springthorpe, *Nat. Rev. Drug Discov*. 2007, **6**, 881-890; c) X. Wu, L. He, H. Neumann and M. Beller, *Chem. Eur. J.* 2013, **19**, 12635-12638.
- ⁴⁰ 16 There are several methods to prepare N-heterocyclic compounds, in which the coupling of 2-substituted aniline with aldehydes was the popular choice. Although aldehydes can serve as the efficient starting material for the preparation of N-heterocyclic compounds, the aldehydes are chemically unstable for preparation and suffer
- 45 from difficult storage. Thus, efficient alternative methods are still desired. For selected examples using aldehyde as starting material see: a) G. Wang, C. Miao, H. Kang and Bull. Chem. Soc. Jpn. 2006, 79, 1426-1430; b) C. Balakumar, P. Lamba, D. P.Kishore, B. L. Narayana, K. V. Rao, K. Rajwinder, A. R. Rao, B. Shireesha and B.
- ⁵⁰ Narsaiah, *Eur. J. Med. Chem.* 2010, **45**, 4904-4913; c) C. Balakumar, P. Lamba, D. P. Kishore, B. L. Narayana, K. V. Rao, K. Rajwinder, A. R. Rao, B. Shireesha and B. Narsaiah, *Eur. J. Med. Chem.* 2010, **45**, 4904-4913; For selected examples of Ullmann-type *N*-arylation, see: d) D. S. Yang, Y. Y. Wang, H. J. Yang, T. Liu and
- H. Fu, Adv. Synth. Catal. 2012, 354, 477-482; for selected examples using alcohol as starting material, see: e) M. Heidary, M. Khoobi, S. Ghasemi, Z. Habibi and M. Ali Faramarzi, Adv. Synth. Catal. 2014, 1789-1794; for CO insertion and intramolecular cyclization, see: f) B. Ma, Y. Wang, J. L. Peng and Q. Zhu, J. Org. Chem. 2011, 76,
- 60 6362-6366; for selected examples using amines as starting material, see: g) T. B. Nguyen, L. Ermolenko and A. Al-Mourabit, *Green Chem.* 2013, **15**, 2713-2717.
- 17 When ethyl 2-phenylacetate **2n** (efficient substrate for the current catalytic system, see Table 2) was loaded in the absence of **1a** under
- 65 similar condition, ethyl 2-oxo-2-phenylacetate was produced in 31% yield after 12 hours (eq 8). It should be noted that the hydrolysis of ethyl 2-phenylacetate does not occur during the reaction process,

because 100% 2n was recovered from the reaction system under N_2 atmosphere (eq 9).



18 M. Kamitani, M. Ito, M. Itazaki and H. Nakazawa, Chem. Commun. 2014, 7941-7944.