ue

Cyclic and open-chain derivatives of bis(trihydrosilyl)benzenes[†]

Robert Schröck, Karl-Heinz Dreihäupl, Alexander Sladek and Hubert Schmidbaur*

Anorganisch-chemisches Institut der Technischen Universität München, Lichtenbergstrasse 4, D-85747 Garching, Germany



bis(trifluoromethylsulfonyloxysilyl)benzene, obtained from 1,2-bis(p-tolylsilyl)benzene by treatment with 2 equivalents of triflic acid (CF₃SO₃H), at -10 °C. Treatment of this intermediate with primary amines, or with H₂C=P(NMe₂)₃ in a transylidation process, afforded the heterocycles in high yields. The molecular structure of N-(p-bromophenyl)-1,3-disilaisoindoline has been determined by X-ray diffraction. The molecules are almost completely flat with only a minor curvature. Open-chain disiloxanes $(m,p-H_3SiC_6H_4SiH_2)_2O$ were prepared from the corresponding arylsilyl(silyl)benzenes, which were converted into the trifluoromethylsulfonyloxysilyl-(silyl)benzenes (stable up to -10 °C) by triflic acid and hydrolysed by water-NEt₃.

Cyclic and open-chain siloxanes, silazanes and carbosilanes are important starting materials for the preparation of silicone polymers as well as ceramics based on silicon nitride and silicon carbide systems.¹ The large variety of methods that can be employed for the formation of siloxane and silazane polymers have recently been summarised in a detailed review.²

1,3-Disila-isoindolines (A), -indans (B) and their siloxane analogues (C) are expected to be versatile precursors for the preparation of polymers with the general formula (-XSiR₂- $C_6H_4SiR_2$, $(X = NR, O \text{ or } CR_2)$. While Si-alkylated compounds of this type were extensively studied in the past,³⁻⁸ the knowledge on systems based on o-bis(trihydrosilyl)benzene as a prototype is still rather limited. Recent investigations included a platinum(IV) complex $[Pt[C_6H_4(SiH_2)_2]_2(PEt_3)_2]$, synthesized by Tanaka and co-workers,⁹ which showed great promise as both an intermediate and a catalyst for dehydrogenative or desilanative coupling of arylsilanes. Our own studies in the chemistry of various poly(trihydrosilyl)-alkanes, 10,11 -alk-enes, 12 and -arenes $^{13-17}$ suggested new pathways also for the preparation of related compounds with -SiH₂XSiH₂units.^{18,19} The preparation of cyclic disiloxanes and disilazanes derived from 1,8-disilylnaphthalene described in a previous paper is an example of the synthetic strategies employed in this context.20

In this report we present a convenient synthesis for cyclic derivatives of o-disilylbenzene (D) and open-chain disiloxanes still containing 'free' silyl groups (E) suitable for dehydrogenative or desilanative coupling and related reactions.

Results and Discussion

The preparative method employed successfully in our studies of cyclic 1,8-disilylnaphthalene derivatives is also applicable for the synthesis of 1,3-disilaisoindolines. Treatment of 1,2-bis(ptolylsilyl)benzene 1 with 2 equivalents of triflic acid (CF_3SO_3H) at -10 °C gives the corresponding 1,2-bis(trifluoromethylsulfonyloxysilyl)benzene 2 as the key intermediate. This precursor is extremely sensitive to moisture and thermally labile and should not be isolated, but rather used in situ in all further reactions. For example, cyclisation is readily accomplished with one equivalent of a primary amine in the presence of two equivalents of triethylamine to trap the resulting triflic acid (Scheme 1).



If the reaction temperature is carefully controlled, the 1,3disilaisoindolines 3-8 are obtained in good yields as colourless liquids (3, 7) or solids (4-6, 8), which can be characterised by standard spectroscopic methods and elemental analysis. The formation of oligo- or poly-meric side products has not been observed. The benzyl-substituted isoindolines 7 and 8 are quite sensitive to hydrolysis, while the alkyl and aryl derivatives are more robust owing to the steric protection of the disilazane linkage.

Single crystals of N-(p-bromophenyl)-1,3-disilaisoindoline 6 (from hexane) were investigated by X-ray diffraction. Compound 6 crystallises in the monoclinic space group $P2_1/n$ with four formula units in the unit cell. The molecules have no crystallographically imposed symmetry, but the coordinates of the atoms closely obey mirror symmetry (Fig. 1). The five-membered ring containing the silazane unit is planar, and the configuration of the nitrogen atom is also planar (sum of angles 359.9°). The molecule as a whole is slightly curved, and the isoindoline and phenyl rings are tilted by 13° (Fig. 2). The endocyclic angles at the silicon atoms (99°) are clearly smaller than the ideal tetrahedral angle, and the Si(1)-N-Si(2) angle (113.8°) is smaller than the 120° reference. These details suggest a significant strain of the five-membered ring, which seems to be even more pronounced in the corresponding siloxane and silathiane, which polymerise spontaneously and cannot be isolated.

[†] Non-SI units employed: $eV \approx 1.60 \times 10^{-19}$ J, Torr ≈ 133 Pa



Scheme 1 $R' = CF_3SO_3$. (i) $2CF_3SO_3H$, toluene; (ii) NRH_2 , $2NEt_3$



Fig. 1 Molecular structure of compound 6 with atomic numbering (ORTEP, 21 50% probability ellipsoids for non-hydrogen atoms; arbitrary radii for hydrogen atoms). Selected bond lengths (Å) and angles (°): Si(1)–N 1.746(3), Si(2)–N 1.742(3), Si(1)–C(1) 1.858(3), Si(2)–C(2) 1.858(3), N-C(7) 1.409(4) and Si–H (average) 1.42; Si(1)–N-Si(2) 113.8(1), Si(1)–N-C(7) 122.9(2), Si(2)–N-C(7) 123.2(2), N-Si(1)–C(1) 98.9(1), N-Si(2)–C(2) 99.1(1), C(1)–C(2)–Si(2) 113.9(2), C(2)–C(1)–Si(1) 114.1(2), H(01)–Si(1)–H(02) 108(2), H(03)–Si(2)–H(04) 107(2), C(6)–C(1)–Si(1) 126.5(3), C(6)–C(1)–C(2) 119.4(3), C(3)–C(2)–Si(2) 126.5(3) and C(6)–C(1)–Si(1) 126.5(3)



Fig. 2 Molecular structure of compound 6, side projection

These structural features also explain the lability of the corresponding ylide 9 in solution [as compared with the peribridged naphthalene $(Me_2N)_3P=C(SiH_2)_2C_{10}H_6^{20}]$. Compound 9 is easily synthesized by a transylidation reaction 22-24of three equivalents of tris(dimethylamino)phosphonio methanide and one equivalent of 1,2-bis(trifluoromethylsulfonyloxysilyl)benzene 2 (Scheme 2). In this process, two equivalents of the ylide function as a base to trap the triflic acid which is produced in the cyclisation. The product 9 is a colourless solid which can be isolated by recrystallisation from hexane. In solution it decomposes rapidly even at low temperature, whereas it is stable in the solid state under an inert atmosphere at room temperature. The NMR spectra are typical for ylidic species. The carbon atom in the bridgehead position between the phosphorus and the two silicon atoms has its ¹³C resonance at very high field ($\delta - 12.6$) with a large ${}^{1}J(PC)$ coupling constant of 155 Hz, indicating sp² hybridisation.25,26



The access to open-chain derivatives (type E) of the disilylbenzenes is accomplished by regioselective and stepwise protodearylation reactions, in which only one *p*-tolyl group is cleaved from the bis(*p*-tolylsilyl)arenes. The resulting *p*-tolylsilyl(trifluoromethylsulfonyloxysilyl)benzenes can be reduced with LiAlH₄ to give *p*-tolylsilyl(silyl)arenes. This reaction sequence shows a surprisingly high selectivity.²⁷ The resulting arylsilyl(silyl)arenes (*e.g.* 10 and 11) are ideal precursors for the preparation of monofunctional silyl triflates of polysilylarenes in a second protodearylation round. The trifluoromethylsulfonyloxysilylarenes generated thereby are treated *e.g.* with water to yield the corresponding siloxanes 12 and 13, which can serve as model compounds for a large group of derivatives with this structural principle (Scheme 3).

1,3-bis(3-silylphenyl)disiloxane 12 1,3-bis(4-silylphenyl)disiloxane 13

The corresponding 1,3-bis(2-silylphenyl)disiloxane could not be detected in the analogous reaction mixture, and only polymeric products and 1,2-disilylbenzene were found. The siloxanes 12 and 13 were isolated in moderate yields as colourless oils by vacuum condensation. Their composition was confirmed by standard analytical and spectroscopic techniques. The ¹⁷O NMR signals could be detected with chemical shifts of $\delta - 2.7$ (12) and -3.5 (13) in a range typical for this type of siloxanes.^{18,20}

Conclusion

The present studies have thus given access to (1) heterocyclic molecules of the 1,3-disila-isoindoline and -indan series, and (2) open-chain siloxanes with $-H_2SiXSiH_2$ - and SiH_3 units. These functions allow further reactions including dehydrogenative or desilanative coupling. Future studies will focus on Si–Si coupling reactions using these molecules and their derivatives containing other heteroatoms.

Experimental

All experiments were carried out under dry purified nitrogen. Solvents and glassware were dried and kept under nitrogen. NMR: JEOL GX 400 spectrometer. Mass spectra: Mat 311 (electron impact, EI; 70 eV) or gas chromatograph—mass spectrometer with HP 5971 A mass-selective detector (EI, 70 eV).



1,3-Disilaisoindolines 3-8, general procedure

A solution of 1,2-bis(p-tolylsilyl)benzene 1¹⁶ (2.00 g, 6.3 mmol) in toluene (40 cm³) was cooled to -10 °C and triffic acid (1.11 cm³, 12.6 mmol) added. The solution was stirred for 1 h at this temperature and then a mixture of the amine (6.3 mmol) and NEt₃ (1.81 cm³, 12.6 mmol) was added. After 30 min the reaction mixture was warmed to room temperature and the oily triethylammonium triflate separated. The solvent was removed under vacuum and the residue purified by distillation, sublimation or recrystallisation.

N-tert-Butyl-1,3-disilaisoindoline 3. Yield 73% (0.95 g), b.p. 45 °C (0.01 Torr) (Found: C, 57.2; H, 7.9; N, 6.9; Si, 27.4. C₁₀H₁₇NSi₂ requires C, 57.9; H, 8.2; N, 6.8; Si, 27.1%). NMR $(C_6D_6, 20 \text{ °C})$: ¹H, δ 1.40 (9 H, s, CH₃), 5.29 [4 H, s, ¹J(SiH) = 210, SiH₂], 7.24 and 7.55 (4 H, m each, C_6H_4); ¹³C-{¹H}, δ 32.1 (CH₃), 51.3 (CN), 129.6 (C^{5,6}), 132.7 (C^{4,7}), 141.8 (C_{ipso}, $C^{8,9}$); ²⁹Si, δ -34.5 [tm, ¹J(SiH) = 211 Hz, SiH₂]. IR: \tilde{v}_{max} 2112 (SiH) and 978 cm⁻¹ (SiN). Mass spectrum: m/z = 207 (M^+) , 192 $(M^+ - CH_3)$, 151 $(M^+ - C_4H_8)$, 133 $(M^+ NH_2C_4H_9$) and 105 ($C_6H_5SiH_3$).

N-Cyclohexyl-1,3-disilaisoindoline 4. Yield 83% (1.22 g), m.p. 18-20 °C (Found: C, 61.2; H, 7.9; N, 6.1; Si, 24.4. C₁₂H₁₉NSi₂ requires C, 61.7; H, 8.1; N, 6.0; Si, 24.1%). NMR $(C_6D_6, 20 \text{ °C})$: ¹H, $\delta 0.93-2.00$ (10 H, m, C_6H_4), 2.89 (1 H, m, NCH), 5.18 [4 H, s, ${}^{1}J(SiH) = 208$, SiH₂], 7.15 and 7.46 (4 H, m each, C_6H_4); ¹³C-{¹H}, δ 25.8 (C_y), 26.1 (C_{δ}), 36.6 (C_{β}), 54.2 (C_{α}), 129.6 (C^{5,6}), 132.9 (C^{4,7}), and 141.5 (C_{*i*pso}, C^{8.9}); ²⁹Si, δ 31.7 [tm, ${}^{1}J(SiH) = 208$ Hz, SiH_{2}]. IR: \tilde{v}_{max} 2115 (SiH) and 981 (SiN) cm⁻¹. Mass spectrum: $m/z = 233 (M^+)$, 190 (M^+ – C_3H_7), 177 ($M^+ - C_4H_8$), 150 ($M^+ - C_6H_{11}$), 135 and 105.

N-Phenyl-1,3-disilaisoindoline 5. Yield 80% (1.13 g), m.p. 108-110 °C (Found: C, 63.0; H, 6.1; N, 6.1; Si, 24.8. C₁₂H₁₃NSi₂ requires C, 63.4; H, 5.7; N, 6.1; Si, 24.6%). NMR $(C_6D_6, 20 \text{ °C})$: ¹H, δ 5.27 [4 H, s, ¹J(SiH) = 217, SiH₂], 6.83 (1 H, m, H¹³), 7.09 and 7.34 (4 H, m each, H⁴⁻⁷) and 7.15 (4 H, m, $H^{11,12,14,15}$; ¹³C-{¹H}, δ 119.8 (C^{11,15}), 121.0 (C¹³), 129.9 $(C^{12.14})$, 130.1 $(C^{5.6})$, 133.0 $(C^{4.7})$, 139.9 $(C_{ipso}, C^{8.9})$ and 147.0 (C_{ipso}, C^{10}) . ²⁹Si, δ – 30.6 [tdm, ¹J(SiH) = 217, ³J(SiH) = 6 Hz, SiH₂]. IR (Nujol): \tilde{v}_{max} 2139 (SiH) and 943 cm⁻¹ (SiN). Mass spectrum: m/z = 227 (M^+), 149 ($M^+ - C_6H_5$), 134 ($M^+ - NH_2C_6H_5$), 105 ($M^+ - SiH_2NC_6H_5$) and 93.

N-(p-Bromophenyl)-1,3-disilaisoindoline 6. Yield 75% (1.44 g), m.p. 144-146 °C (Found: C, 47.0; H, 3.9; Br, 25.4; N, 4.7; Si, 18.8. C₁₂H₁₂BrNSi₂ requires C, 47.1; H, 3.9; Br, 26.1; N, 4.6; Si, 18.3%). NMR (C_6D_6 , 20 °C): ¹H, δ 4.88 [4 H, s, ¹J(SiH) = 217, SiH₂], 6.76 and 7.21 [4 H, d each, ³J(HH) = 9, NC₆H₄Br], 7.13 and 7.33 (4 H, m each, C₆H₄Si₂); 13 C-{¹H}, δ 113.6 (C¹³), 121.3 (C^{11,15}), 130.2 (C^{5.6}), 132.7 (C^{12,14}), 133.0 $(C^{4,7})$, 139.4 $(C_{ipso}, C^{8,9})$ and 146.0 (C^{10}) ; ²⁹Si, $\delta - 30.0$ [tdm, ¹J(SiH) = 217, ³J(SiH) = 4 Hz, SiH₂]. IR (Nujol): \tilde{v}_{max} 2142 (SiH) and 942 cm⁻¹ (SiN). Mass spectrum: m/z = 305/307 (M^+) , 225 $(M^+ - Br)$, 148 $(M^+ - C_6H_4Br)$ and 105 $(M^+ - C_6H_4Br)$ $SiH_2NC_6H_4Br$).

N-Benzyl-1,3-disilaisoindoline 7. Yield 58% (0.88 g), m.p. -5 °C (Found: C, 64.5; H, 6.2; N, 5.9; Si, 23.4. C₁₃H₁₅NSi₂ requires C, 64.7; H, 6.2; N, 5.8; Si, 23.3%). NMR/(C₆D₆, 20°C): ¹H, δ 4.02 (2 H, s, CH₂), 5.08 [4 H, s, ¹J(SiH) = 210, SiH₂], 7.10 and 7.36 (4 H, m each, C_6H_4), 7.12 and 7.16 (5 H, m each, C_6H_5); ¹³C-{¹H}, δ 48.9 (CH₂), 127.4 (C¹⁴), 128.2 (C^{12.16}), $128.7 (C^{13,15}), 129.7 (C^{5,6}), 132.9 (C^{4,7}), 141.1 (C_{ipso}, C^{11}) and$ 141.6 (C_{ipso} , $C^{8,9}$); ²⁹Si, $\delta - 25.3$ [tm, ¹J(SiH) = 210 Hz, SiH₂]. IR: \tilde{v}_{max} 2121 (SiH) and 973 cm⁻¹ (SiN). Mass spectrum: m/z =241 (M^+) , 163 $(M^+ - C_6H_5)$, 150 $(M^+ - CH_2C_6H_5)$, 135 $(M^+ - H_2 NCH_2 C_6 H_5)$, 105 and 91.

N-(p-Bromobenzyl)-1,3-disilaisoindoline 8. Yield 65% (1.32 g), m.p. 42-44 °C (Found: C, 48.8; H, 4.5; N, 4.3; Si, 17.4. C13H15NSi2 requires C, 48.7; H, 4.4; N, 4.4; Si, 17.5%). NMR $(C_6D_6, 20 \text{ °C})$: ¹H, δ 3.82 (2 H, s, CH₂), 4.99 [4 H, s, ¹J(SiH) = 216, SiH₂], 6.80 and 7.21 (4 H, m each, C₆H₄Br), 7.12 and 7.37 (4 H, m each, C_6H_4); ¹³C, δ 48.9 [tt, ¹J(CH) = 138, ³J(CH) = 4, CH₂], 121.3 [tt, ³J(CH) = 11, ²J(CH) = 4, C¹⁴], 129.7 $[dm, {}^{1}J(CH) = 157, C^{12,16}], 129.8 [dm, {}^{1}J(CH) = 159, C^{5,6}], 131.8 [dd, {}^{1}J(CH) = 166, {}^{3}J(CH) = 6, C^{13,15}], 133.0 [dm,]$ ${}^{1}J(CH) = 156, C^{4,7}], 140.0 \text{ (m, } C_{ipso}, C^{11}) \text{ and } 141.3 \text{ (m, } C_{ipso}, C^{11})$ $C^{8,9}$; ²⁹Si, $\delta - 24.9$ [tdm, ¹J(SiH) = 210, ³J(SiH) = 8 Hz, SiH₂]. IR: $\tilde{\nu}_{max}$ 2123 (SiH) and 975 cm⁻¹ (SiN). Mass spectrum: m/z 319/321 (M^+), 240 (M^+ – Br), 163 (M^+ – C₆H₄Br), 150 $(M^+ - CH_2C_6H_4Br)$, 136 $(M^+ - NCH_2C_6H_4Br)$ 105 and 90.

2-[Tris(dimethylamino)phosphoranylidene]-1,3-disilaindan 9

To a solution of 1,2-bis(p-tolylsilyl)benzene 1^{16} (2.16 g, 6.8 mmol) in toluene (100 cm³) was added triffic acid (1.20 cm³, 13.6 mmol) at -10 °C. The reaction mixture was stirred for 1 h at this temperature and subsequently H₂C=P(NMe₂)₃ (3.88 g, 20.4 mmol) diluted with toluene (10 cm³) was added and the mixture slowly warmed to room temperature. After 30 min the solvent was removed in vacuo, and the residue washed three times with hexane (30 cm³ each). The hexane solution was concentrated in vacuo to yield the crude product. The ylide was purified by recrystallisation from hexane to give colourless crystals (75 %, 1.5 g), m.p. > 200 °C (decomp.) NMR (C_6D_6 , 20 °C): ¹H, δ 2.40 [18 H, d, ${}^{3}J(PH) = 10$, $P(NMe_{2})_{3}$], 5.50 [4 H, s, ${}^{1}J(SiH) = 187$, SiH_{2}], 7.30 and 7.90 (4 H, m each, $C_{6}H_{4}$); ${}^{13}C{}^{1}H$ }, $\delta - 12.6$ [d, ${}^{1}J(PC) = 155$, C^{2} , P=C], 37.7 [d, ${}^{2}J(PC) = 4$, $P(NMe_{2})_{3}$], 133.2 ($C^{5.6}$), 146.7 ($C^{8.9}$) and 146.8 ($C^{4.7}$); ${}^{31}P$ -{¹H}, δ 64.7 (s); ²⁹Si, δ - 33.0 [tdm, ¹J(SiH) = 188, ²J(SiP) = 12 Hz (observed in the ²⁹Si-{¹H} spectrum), SiH₂]. Mass spectrum: $m/z = 311 (M^+)$, 268 $(M^+ - NMe_2)$, 244 $(M^+ - 2)$ NMe_2), 179 (M^+ – 3 NMe_2), 148 [M^+ – P(NMe_2)₃], 135 $[M^+ - C = P(NMe_2)_3], 119 [P(NMe_2)_2]$ and 106.

1,3-Bis(silylphenyl)disiloxanes 12 and 13, general procedure

To a solution of 1-(p-tolylsilyl)-3-silylbenzene 10 or 1-(ptolylsilyl)-4-silylbenzene 11²⁷ (0.80 g, 3.5 mmol) in toluene (50 cm³) was added triflic acid (0.31 cm³, 3.5 mmol) at -10 °C. After 1 h at this temperature a mixture of water (0.031 cm³, 1.7 mmol) and NEt₃ (2.1 cm³, 3.5 mmol) was added. The oily [NEt₃H][CF₃SO₃] was separated at room temperature, the solvent evaporated and the colourless oil recondensed in a vacuum (110 °C, 0.01 Torr).

1,3-Bis(3-silylphenyl)disiloxane 12. Yield 25% (0.25 g) (Found: C, 49.3; H, 6.3; Si, 39.1. C₁₂H₁₈OSi₄ requires C, 49.6; H, 6.2; Si, 38.7%). NMR (C₆D₆, 20 °C): ¹H, δ 4.17 [6 H, s, ${}^{1}J(SiH) = 200, SiH_{3}], 5.22 [4 H, s, {}^{1}J(SiH) = 217, SiH_{2}], 7.01$ $[2 H, t, {}^{3}J(HH) = 7, H^{5}], 7.36 \text{ and } 7.43 [4 H, d each,$ ${}^{3}J(\text{HH}) = 7, \text{H}^{4,6}$ and 7.75 (2 H, s, H²); ${}^{13}\text{C}-\{{}^{1}\text{H}\}, \delta$ 133.8 (C³), 135.7 (C¹), 137.4 and 137.5 (C^{4,6}), 138.2 (C⁵) and 141.9 (C^2) ; ²⁹Si, $\delta - 59.7$ [qtm, ¹J(SiH) = 201, ³J(SiH) = 7, SiH₃] and $-24.9 \text{ [tm } {}^{1}J(\overline{SiH}) = 220 \text{ Hz}, \text{ SiH}_{2}\text{]}; {}^{17}O, \delta -2.7. \text{ Mass}$ spectrum: $m/z = 290 (M^+)$, 259 $(M^+ - \text{SiH}_3)$, 227 $(M^+ - 2)$ SiH_3 , 181 ($M^+ - C_6H_4SiH_3$), 151 [$M^+ - SiH_2(C_6H_4)SiH_3$],

Table 1 Crystallographic data for compound 6

Formula	$C_{12}H_{12}BrNSi_2$
М	306.32
Crystal system	Monoclinic
Crystal size/mm	$0.18 \times 0.24 \times 0.40$
Space group	$P2_1/n$ (no. 14)
a/Å	8.777(1)
b/Å	8.080(1)
c/Å	18.914(2)
β/°	102.20(1)
$U/Å^3$	1331.5(3)
$D_{2}/g \mathrm{cm}^{-3}$	1.552
Z	4
μ (Mo-K α)/cm ⁻¹	32.90
<i>T/</i> °C	-62
Absorption correction	DIFABS ³⁰
T_{\min}, \hat{T}_{\max} (%)	34.1, 100
Scan mode	ω
Scan width/° in ω	1.30
hkl Ranges	- 10 to 10, 0-10, 0-24
Measured data	2917
Unique data	2835
Observed data $[F] \ge 4\sigma(F)$]	2049
No parameters	193
$w R^2$	0.097
R based on $F(OMIT4)$	0.036
Weighting scheme w	$[\sigma^2(F^2) \pm (0.0367n)^2 \pm 1.3312n^{-1}$
Weighting scheme, w	$n = [max(F^2 0) + 2F^2]/3$
Pesidual electron density/	$p = [max(r_0, 0) + 2r_c]/3$
λ^{-3}	+ 0.37, - 0. 4 7
CA	

137 $[M^+ - \text{OSiH}_2(\text{C}_6\text{H}_4)\text{SiH}_3]$ and 105 $[M^+ - \text{SiH}_2\text{O-SiH}_2(\text{C}_6\text{H}_4)\text{SiH}_3]$.

1,3-Bis(4-silylphenyl)disiloxane 13. Yield 35% (0.35 g) (Found: C, 49.8; H, 6.3; Si, 38.8. $C_{12}H_{18}OSi_4$ requires C, 49.6; H, 6.2; Si, 38.7%). NMR (C_6D_6 , 20 °C): ¹H, δ 4.17 [6 H, s, ¹J(SiH) = 200, SiH₃], 5.23 [4 H, s, ¹J(SiH) = 220, SiH₂], 7.34 and 7.41 [8 H, d each, ³J(HH) = 7, C_6H_4]; ¹³C-{¹H}, δ 131.3 (C⁴), 133.8 (C^{2.6}), 135.4 (C¹) and 135.7 (C^{3.5}); ²⁹Si, δ - 59.8 [qtt, ¹J(SiH) = 200, ³J(SiH) = 6, ⁴J(SiH) = 1, SiH₃] and -25.1 [ttm, ¹J(SiH) = 220, ³J(SiH) = 6 Hz, SiH₂]; ¹⁷O, δ -3.5. Mass spectrum: m/z = 290 (M^+), 259 (M^+ - SiH₃), 227 (M^+ - 2 SiH₃), 181 (M^+ - C₆H₄SiH₃), 151 [M^+ - SiH₂(C₆H₄)SiH₃], 137 [M^+ - OSiH₂(C₆H₄)SiH₃] and 105 [M^+ - SiH₂OSiH₂-(C₆H₄)SiH₃].

Crystallography

A suitable crystal of compound 6 was sealed into a glass capillary and examined on an Enraf-Nonius CAD 4 diffractometer. Graphite-monochromated Mo-K α radiation ($\lambda = 0.710$ 69 Å) was used. The structure was solved by direct methods (SHELXTL²⁸) and refined against F^2 (SHELXL 93²⁹). All heavy atoms were refined with anisotropic displacement parameters. All hydrogen atoms were located. The final unit-cell parameters and specific data collection parameters are summarised in Table 1.

Atomic coordinates, thermal parameters, and bond lengths

and angles have been deposited at the Cambridge <u>Grystalloc</u> graphic Data Centre (CCDC). See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1996, Issue 1. Any request to the CCDC for this material should quote the full literature citation and the reference number 186/230.

References

- N. Auner and J. Weis, Organosilicon Chemistry II, VCH, Weinheim, 1996.
- M. Birot, J.-P. Pillot and J. Dunoguès, *Chem. Rev.*, 1995, 95, 1443.
 M. Laguerre, J. Dunogues, R. Calas and N. Duffaut, *J. Organomet.*
- Chem., 1976, 112, 49. 4 R. J. P. Corriu, J. J. E. Moreau and M. Pataud-Sat, J. Organomet.
- Chem., 1982, **228**, 301.
- 5 K. Shiina, J. Organomet. Chem., 1986, 310, C57.
- 6 R. P. Bonar-Law, A. P. Davis, B. J. Dorgan, M. T. Reetz and A. Wehrsig, *Tetrahedron Lett.*, 1990, **31**, 6725.
- 7 Y. Uchimaru, A. Mouneer M. El Sayed and M. Tanaka, Organometallics, 1993, 12, 2065.
- 8 K. Tamao, G.-Ri Sun and A. Kawachi, J. Chem. Soc., Chem. Commun., 1995, 2079.
- 9 S. Shimada, M. Tanaka and K. Honda, J. Am. Chem. Soc., 1995, 117, 8289.
- 10 R. Hager, O. Steigelmann, G. Müller, H. Schmidbaur, H. E. Robertson and D. W. H. Rankin, Angew. Chem., 1990, 102, 204; Angew. Chem., Int. Ed. Engl., 1990, 29, 201.
- 11 H. Schmidbaur, J. Zech, D. W. H. Rankin and H. E. Robertson, Chem. Ber., 1991, 124, 1953.
- 12 C. Rüdinger, H. Beruda and H. Schmidbaur, Z. Naturforsch., Teil B, 1994, 49, 1348.
- 13 C. Rüdinger, P. Bissinger, H. Beruda and H. Schmidbaur, Organometallics, 1992, 11, 2867.
- 14 C. Rüdinger, H. Beruda and H. Schmidbaur, Chem. Ber., 1992, 125, 1401.
- 15 R. Schröck, K. Angermaier, A. Sladek and H. Schmidbaur, Organometallics, 1994, 13, 3399.
- 16 R. Schröck, A. Sladek and H. Schmidbaur, Z. Naturforsch., Teil B, 1994, 49, 1036.
- 17 R. Schröck, K. Angermaier, A. Sladek and H. Schmidbaur, J. Organomet. Chem., 1996, 509, 85.
- 18 S. Bommers and H. Schmidbaur, Chem. Ber., 1994, 127, 1359.
- 19 S. Bommers and H. Schmidbaur, *Phosphorus Sulfur, Silicon Relat. Elem.*, 1995, 105, 171.
- 20 R. Schröck, K.-H. Dreihäupl, A. Sladek, K. Angermaier and H. Schmidbaur, *Chem. Ber.*, 1996, **129**, 495.
- 21 C. K. Johnson, ORTEP, Report ORN2-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- 22 H. J. Bestmann, Chem. Ber., 1962, 95, 58.
- 23 N. W. Mitzel, A. Schier, H. Beruda and H. Schmidbaur, *Chem. Ber.*, 1992, **125**, 1053.
- 24 K.-H. Dreihäupl, K. Angermaier, J. Riede and H. Schmidbaur, Chem. Ber., 1994, 127, 1599.
- 25 H. Schmidbaur, Adv. Organomet. Chem., 1976, 14, 205.
- 26 A. W. Johnson, Ylides and Imines of Phosphorus, Wiley, New York, 1993.
- 27 R. Schröck, K. Angermaier and H. Schmidbaur, Z. Naturforsch., Teil B, 1995, 50, 613.
- 28 SHELXTL-PLUS, Release 4.1, Siemens Analytical X-Ray Instruments, Madison, WI, 1990.
- 29 G. M. Sheldrick, SHELXL 93, Program for the Refinement of Structures, University of Göttingen, 1993.
- 30 R. F. Stewart, E. R. Davidson and W. T. Simpson, J. Chem. Phys., 1965, 42, 3175.

Received 8th July 1996; Paper 6/04752A