Site-Specific Incorporation of the Alkaline Labile, Oxidative Stress Product (5R)-5,6-Dihydro-5-hydroxythymidine in an Oligonucleotide

Tracy J. Matray and Marc M. Greenberg*

Department of Chemistry, Colorado State University Fort Collins, Colorado 80523

Received April 18, 1994

The nucleobases within nucleic acids are alkylated by electrophiles such as oxygenated benzo[a]pyrenes and are oxidized by activated oxygen species.^{1,2} The damage resulting from such chemical reactions can be mutagenic and/or cytotoxic. In order to understand the origin of mutagenicity, it is important to correlate changes induced by mutated nucleosides on the function and structure of nucleic acids. Determination of the effects that mutated nucleosides impart on the function of polymerase and repair enzymes in vitro and in vivo, as well as spectroscopic characterization of structural alterations induced by these molecules, is made possible by their site-specific incorporation in oligonucleotides.^{1,3-7} In some instances, mutations are introduced in the oligonucleotide postsynthetically.1,5 This strategy is limited by the structural variation of the biopolymers containing the mutated nucleoside that can be prepared, as well as the types of mutations that one can introduce. Incorporation of the mutated monomer as its nucleotide triphosphate or phosphoramidite provides the most versatile synthesis of modified oligonucleotides.⁴ However, site-specific incorporation of some mutated nucleosides in biopolymers during solid-phase synthesis protocols is prohibited by their lability to the alkaline conditions typically used to deprotect chemically synthesized nucleic acids.5 We wish to report the first chemical synthesis of an oligonucleotide containing alkaline labile (5R)-5,6-dihydro-5-hydroxythymidine (1), in which Pd⁰ labile protecting groups are utilized for carrying out the synthesis on a photolabile solid-phase synthesis support.



(5R)-5,6-Dihydro-5-hydroxythymidine (1) is one of several mutated forms of thymidine that are believed to be mutagenic and/or cytotoxic. These mutated nucleosides are formed as a result of the interaction between reactive oxygen species and pyrimidine nucleosides.^{2a} Oligonucleotides containing 5-(hydroxymethyl)-2'-deoxyuridine (2) and 5-hydroxy-5-methylhydantoin (3) were synthesized using conventional protecting groups, because the modified nucleosides withstood deprotection with concentrated ammonium hydroxide.^{4a-c} Site-specific incorpora-

tion of a mixture of stereoisomers of alkali labile thymidine glycol (4) was achieved postsynthetically via osmylation of an oligonucleotide containing a single thymidine.⁵ To our knowledge, there is no chemical reaction that enables one to transform a nucleoside into 1 postsynthetically. Furthermore, 1 decomposes when exposed to concentrated ammonium hydroxide. Hence, the integrity of 1 cannot be maintained when it is incorporated into chemically synthesized oligonucleotides that require alkaline deprotection conditions.

In order to insure that oligonucleotides containing 1 are sufficiently pure for use as mechanistic probes in vitro and in vivo, we sought conditions that obviate the need for harsh base treatment during biopolymer synthesis. With this requirement in mind, allyloxy protecting groups were chosen in order to take advantage of their lability to Pd⁰ at pH 5.5.8 We also elected to utilize a solid-phase synthesis support from which oligonucleotides are cleaved under nonalkaline conditions. Oligonucleotides bound to solid-phase supports via disulfide linkages are cleaved under mild reductive conditions. However, the oligonucleotides released by this method contain 3'-termini that are not compatible with subsequent enzymatic manipulation unless they are subjected to further alkaline hydrolysis.9 We recently reported on a family of photolabile orthogonal linkers that release oligonucleotides containing 3'-hydroxyl groups.10 O-Nitrobenzyl support 5 releases oligonucleotides under conditions that result in less than 2% thymine-thymine dimer formation. The resulting 3'-succinato moiety undergoes hydrolysis under the mildly basic conditions (pH 8.3) encountered during denaturing polyacrylamide gel electrophoresis, yielding oligonucleotides containing 3'-termini that are suitable for enzymatic reactions.



Phosphoramidite 10 was synthesized using methodologies previously described for the synthesis of 1 and other (allyloxy)carbonyl phosphoramidites (Scheme 1).8,11 The stereochemistry at C5 is established during the dihydroxylation of 3'-O-(tertbutyldimethylsilyl)-5'-O-(dimethoxytrityl)thymidine by OsO4. Thymidine and hydroxy-protected derivatives are predisposed to favor formation of the respective 5R, 6S stereoisomer, such as 6, which was isolated as a single diastereomer in 87% yield.¹¹ Following esterification of 6 (78%), the protected thymidine C5hydrate (7) was isolated in 68% yield via photoinduced SET reduction.11,12 Bis-allyloxy nucleoside 8 was obtained via acylation of the dianion of 7 by diallyl pyrocarbonate. A number of deprotonation methods were investigated, but t-BuMgCl in THF was found to be the most effective. In order to prevent cleavage of the allyloxy carbonate protecting groups, desilylation was carried out using buffered TBAF (0.5 M AcOH, 0.5 M TBAF).13

 ^{(1) (}a) Shibutani, S.; Margulis, L. A.; Geacintov, N. E.; Grollman, A. P. Biochemistry 1993, 32, 7531. (b) Shibutani, S.; Takeshita, M.; Grollman, A. P. Nature 1991, 349, 431. (c) Harris, C. M.; Zhou, L.; Strand, E. A.; Harris, T. M. J. Am. Chem. Soc. 1991, 113, 4328.
 (2) (a) von Sonntag, C. The Chemical Basis of Radiation Biology; Taylor

^{(2) (}a) von Sonntag, C. The Chemical Basis of Radiation Biology; Taylor and Francis: Philadelphia, PA, 1987. (b) Sies, H. Angew. Chem., Int. Ed. Engl. 1986, 25, 1058. (c) Imlay, J. A.; Linn, S. Science 1988, 240, 1302. (d) Feig, D. I.; Loeb, L. A. Biochemistry 1993, 32, 4466.

⁽³⁾ Kouchakdjian, M.; Bodepudi, V.; Shibutani, S.; Eisenberg, M.; Johnson, F.; Grollman, A. P.; Patel, D. J. Biochemistry 1991, 30, 1403.

^{(4) (}a) Sowers, L. C.; Beardsley, G. P. J. Org. Chem. 1993, 58, 1664. (b) Conte, M. R.; Galeone, A.; Avizonis, D.; Hsu, V. L.; Mayol, L.; Kearns, D. R. Bioorg. Med. Chem. Lett. 1992, 2, 79. (c) Guy, A.; Dubet, J.; Téoule, R. Tetrahedron Lett. 1993, 34, 8101. (d) Shibutani, S.; Bodepudi, V.; Johnson, F.; Grollman, A. P. Biochemistry 1993, 32, 4615. (e) Schulhof, J. C.; Molko, D.; Teoule, R. Nucleic Acids Res. 1988, 16, 319.

⁽⁵⁾ Clark, J. M.; Beardsley, G. P. Biochemistry 1987, 26, 5398.

^{(6) (}a) Dizdaroglu, M.; Laval, J.; Boiteux, S. *Biochemistry* 1993, 32, 12105.
(b) Mao, B.; Li, B.; Amin, S.; Cosman, M.; Geacintov, N. E. *Biochemistry* 1993, 32, 11785.

Scheme 1



Figure 1. Polyacrylamide gel electrophoresis of 5'-32P-labeled 12 and 13. Lanes 1 and 3, 12. Lanes 2 and 4, 13. Lanes 1 and 2, no alkali treatment. Lanes 3 and 4, treatment with 1 M NaOH, 6 h, 55 °C.

Phosphitylation of 9 was carried out in the final step using literature procedures.14

The stability of 1 to conditions typically employed for the deprotection of allyloxy protecting groups was examined prior to synthesizing an octadecamer containing (5R)-5,6-dihydro-5hydroxythymidine. Treatment of 11 with Pd₂(dba)₃CHCl₃ and PPh₃ in 0.5 M BuNH₂/HCO₂H at 55 °C for 1 h resulted in complete conversion of 11 to 1, as analyzed by TLC and reversephase HPLC.⁸ Neither thymidine nor any other nucleoside product was detected when 11 was subjected to these reaction conditions.



An octadecamer (12) containing a single molecule of 1 was synthesized via automated synthesis using previously reported cycles.^{8,10a} The only restriction imposed upon the design of the oligonucleotide's sequence was the avoidance of consecutive thymidines, in order to guard against small amounts of photodimer

formation. The crude oligonucleotide was subjected to Pd⁰ deprotection, followed by band-pass photolysis ($\lambda_{max} = 400 \text{ nm}$) for 3 h in order to cleave the oligonucleotide from the controlled pore glass support. Following the prescribed workup, the oligonucleotide was purified by denaturing polyacrylamide gel electrophoresis (20% polyacrylamide) and desalted using a reversephase purification cartridge.10a

Purified 12 and an octadecamer (13) that differs from 12 solely by having a thymidine substituted for 1 were labeled with ³²P at their 5'-termini and shown to comigrate on a 20% denaturing polyacrylamide gel (Figure 1). Treatment of the labeled oligonucleotides with 1 M NaOH does not affect 13 but results in complete cleavage of 12 to yield a labeled fragment that moves faster than the Bromophenol Blue dye. The mobility of the fragment is consistent with the presence of an alkali labile site at the position where 1 is located in 12.15 Further evidence for the presence of 1 in the oligonucleotide was elicited by acidic hydrolysis of 12 and subsequent silvlation of the nucleobases released.^{16,17} The mixture of silvlated nucleobases was analyzed by GC/MS using selected ion monitoring (supplementary material).17 The peak corresponding to trisilylthymine C5hydrate coeluted with material produced from hydrolysis of independently synthesized 1.11a

The methodology discussed above will enable the investigation of the effects of alkali labile mutated nucleosides (e.g., 1) on the structure and biological function of the nucleic acids in which they are present. Furthermore, this methodology demonstrates that photolabile orthogonal solid-phase synthesis supports can be used to incorporate alkali labile nucleosides in oligonucleotides which contain 3'-termini suitable for enzymatic manipulation.¹⁰

Acknowledgment. Financial support of this research from the National Institutes of Health (GM-46534) is appreciated. The authors are grateful to Professor Robert M. Williams for helpful discussions.

Supplementary Material Available: Experimental procedures, spectral characterization for all new compounds, and GC/MS SIM trace of hydrolyzed 12 (7 pages). This material is contained in many libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.

(7) (a) Wood, M. L.; Esteve, A.; Morningstar, M. L.; Kuziemko, G. M.; Essigmann, J. M. Nucleic Acids Res. 1992, 20, 6023. (b) Dosanjh, M. K.; Singer, B.; Essigmann, J. M. Biochemistry 1991, 30, 7027.

(8) Hayakawa, Y.; Wakabayashi, S.; Kato, H.; Noyori, R. J. Am. Chem. Soc. 1990, 112, 1691

(9) (a) Gupta, K. C.; Kumar, P.; Sathyanarayana, S. Nucleic Acids Res. 1991, 19, 3019. (b) Asseline, U.; Bonfils, E.; Kurfurst, R.; Chassignol, M.; Roig, V.; Thuong, N. T. Tetrahedron 1992, 48, 1233. (c) Zuckermann, R.; Corey, D.; Schultz, P. Nucleic Acids Res. 1987, 15, 5305.

(10) (a) Greenberg, M. M.; Gilmore, J. L. J. Org. Chem. 1994, 59, 746.
 (b) Greenberg, M. M. Tetrahedron Lett. 1993, 34, 251.

(11) (a) Barvian, M. R.; Greenberg, M. M. J. Org. Chem. 1993, 58, 6151. (b) Vaishnav, Y.; Holwitt, E.; Swenberg, C.; Lee, H.-C.; Kan, L.-S. J. Biomol. Struct. Dyn. 1991, 8, 935

(12) (a) Barvian, M. R.; Greenberg, M. M. Tetrahedron Lett. 1992, 33, 6057. (b) Saito, I.; Ikehira, H.; Kasatani, R.; Watanabe, M.; Matsuura, T. J. Am. Chem. Soc. 1986, 108, 3115.

(13) Danishefsky, S. J.; McClure, K. F.; Randolph, J. T. Science 1993, 260, 1307.

 (14) Hamamoto, S.; Takaku, H. Chemistry Lett. 1986, 1401.
 (15) Cleavage of 5'-³²P-labeled 12 releases a labeled tetramer, which should migrate faster than Bromophenol Blue on a 20% denaturing polyacrylamide gel. Maniatis, T.; Fritsch, E. F.; Sambrook, J. Molecular Cloning, A Laboratory Manual; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, NY, 1982; p 185.

(16) Compositional analysis of oligonucleotides is often carried out via enzymatic digestion, followed by HPLC analysis of the liberated nucleosides. However, 1 does not absorb strongly above 214 nm, which hindered its detection via the UV detectors available to us. See: (a) Crain, P. F. Methods Enzymol. 1990, 193, 782. (b) Eadie, J. S.; McBride, L. J.; Efcavitch, J. W.; Hoff, L. B.; Cathcart, R. Anal. Biochem. 1987, 165, 442.

(17) (a) Dizdaroglu, M. Methods Enzymol. 1990, 193, 842. (b) Barvian, M. R.; Greenberg, M. M., in preparation.