

SYNTHESIS OF 3-AMINO-4-INDOLYL-2-PIPERIDONES: TRYPTOPHAN-DERIVED PSEUDODIPEPTIDES

Ricardo Rodríguez, Anna Díez, and Mario Rubiralta*

Laboratori de Química Orgànica. Facultat de Farmàcia. Universitat de Barcelona, 08028-Barcelona, Spain


Ernest Giralt

Departament de Química Orgànica. Facultat de Química. Universitat de Barcelona, 08028-Barcelona, Spain

Abstract – ($\alpha R, 3R^*, 4R^*$)-*N*-(2-Acetoxy-1-phenylethyl)-3-allyloxycarbonylamino-4-(1-benzyl-3-indolyl)-2-piperidone (1), a pseudodipeptide which contains a conformationally restricted analogue of tryptophan has been synthesized from anhydride (2) and (*R*)-(-)-phenylglycinol (3).

The study of pseudopeptides has two main goals: finding peptidomimetic molecules which can be used as drugs, and creating moieties, which once inserted in larger peptide chains of known activity, allow the study of the peptide structure-activity relationship.^{1,2} Since several endogenous peptides involved in the regulation of key physiological processes such as somatostatin,³ cholecystokinin (CCK),⁴ or the luteinizing hormone-releasing hormone (LH-RH)⁵ contain a tryptophan that is important in their receptor-binding site, we planned to prepare conformationally restricted analogues of this aminoacid. We thus designed structure (1), a Trp-phenylglycinol pseudodipeptide in which the tryptophan side chain (χ -1) is included in a 2-piperidone ring.⁶


In order to introduce a chiral center α with respect to the lactam nitrogen atom, the synthesis of compound



Scheme 1

(1) was planned by condensation of the conveniently functionalized anhydride (2) with (*R*)-(-)-phenylglycinol.

Anhydride (2) was prepared from triester (10), which in turn was synthesized in four steps as indicated in Scheme 2. Thus, condensation of protected indole-3-carbaldehyde (5) with sodium ethyl nitroacetate gave indolynitroacrylate (6), which was converted into the triester (7) by conjugate addition of the di-*tert*-butyl malonate monopotassium salt. At this point, the nitro group was reduced by hydrogenation in the presence of Raney Ni.⁷ The resulting primary amines (8) were protected as the allyl carbamate (Alloc), and amino triesters *erythro*-9 and *threo*-10⁸ were isolated by flash column chromatography in a 1:2 proportion.



Reagents and conditions: i) TiCl_4 (3 equivalents), pyridine (3 equivalents), THF, room temperature, 24 h (78%); ii) di-*tert*-butyl malonate (1.1 equivalents), $t\text{BuOK}$ (1.1 equivalents), CH_2Cl_2 , room temperature, 1 h, quenching with AcOH (87%); iii) W-2 Raney Ni, EtOH, H_2 (1 atm), 15 h (92%); iv) allyl chloroformate (1.2 equivalents), pyridine (1.2 equivalents), 0°C to room temperature, 1 h (87%).

Scheme 2


Treatment of triester (10) with TFA at 0°C allowed the selective hydrolysis of the *tert*-butyl esters in 80% yield (Scheme 3). The ethyl ester was cleaved subsequently with aqueous 5% KOH at room temperature and the decarboxylation was carried out by refluxing in 4N HCl to yield the *threo* diacid (11) (97% yield).⁹ Further mesylation and Et_3N treatment of diacid (11) led to anhydride (2)¹⁰ in 88% yield.

Condensation of racemic anhydride (2) with (*R*)-(-)-phenylglycinol¹¹ led to a complex mixture, from which only isomer (12)¹² was isolated by column chromatography, in 40% yield.¹³ The structural assignment of compound 12 was confirmed by 2D nOe experiments, thus assuring the stereochemistry of the starting triester (10) and of anhydride (2).

The $\text{NaBH}_4\text{-2N HCl}$ reduction of compound (**12**) occurred, as expected from glutarimides, in a regioselective manner upon the less hindered carbonyl group.¹⁴ However, despite the reaction was carried out at low temperature, only hydroxy amide (**13**) was obtained, resulting from the reduction of the aldehyde generated by ring opening.

Finally, mesylation of hydroxyamide (**13**) and subsequent cyclisation using DBU led to the target lactam (**1**),¹⁵ whose structural assignment was carried out by means of 2D nmr experiments.

With the synthesis of compound (**1**) as a model structure, a synthetic pathway which should be useful to prepare a great variety of tryptophan-derived pseudodipeptides has been established.



Reagents and conditions: i) TFA (5 equivalents), CH_2Cl_2 , 0°C to room temperature, 48 h (white precipitate); ii) 5% aqueous KOH, dioxane, room temperature, 30 min; iii) 4N HCl, reflux, 4 h (78% in three steps); iv) MsCl (1 equivalent), Et_3N (3 equivalents), THF, -20°C , 2 h (88%); v) a. (*R*)-(-)-phenylglycinol (1 equivalent), reflux, 24 h. b. AcCl (excess) reflux, 4 h (40%); vi) NaBH_4 (5.5 equivalents), EtOH-THF, 2N HCl (3 drops every 15 min), -40°C , 2 h (70%); vii) a. MsCl (1 equivalent), Et_3N (1 equivalent), THF, -40°C , 2 h, hexane, filtration. b. DBU (1.2 equivalents), MeCN, room temperature, 1 h (26%).

Scheme 3

ACKNOWLEDGEMENTS

Support for this research has been provided by the CIRIT (Generalitat de Catalunya) through grant QFN92-4303.

REFERENCES AND NOTES

1. D. Mendel, J. Ellman, and P.G. J. Schultz, *J. Am. Chem. Soc.*, 1993, **115**, 4359; A. Giannis and T. Kolter, *Angew. Chem., Int. Engl. Ed.*, 1993, **32**, 1244, and references cited therein.
2. G.R. Marshall, *Tetrahedron*, 1993, **49**, 3547; D.C. Rees, *Current Med. Chem.*, 1994, **1**, 145.

3. I. Lewis and C. Bruns, *Tetrahedron*, 1994, **50**, 7485; W. Kazmierski, W.S. Wire, G.K. Lui, R.J. Knapp, J.E. Shook, T.F. Burks, H.I. Yamamura, and V.J. Hruby, *J. Med. Chem.*, 1988, **31**, 2170; Y. Sasaki, W.A. Murphy, M.L. Heiman, V.A. Lance, and D.H. Coy, *J. Med. Chem.*, 1987, **30**, 1162.
4. K.E. Köver, D. Jiao, S. Fang, and V.J. Hruby, *J. Org. Chem.*, 1994, **59**, 991.
5. F. Haviv, C. A. Palabrica, E.N. Bush, G. Diaz, E.S. Johnson, S. Love, and J. Greer, *J. Med. Chem.*, 1989, **32**, 2340.
6. A similar variation had been described for phenylalanine rigidified analogues used in a study of renin inhibitors: S.E. Laszlo, B.L. Bush, J.J. Doyle, W.J. Greenlee, D.G. Hangauer, T.A. Halgren, R.J. Lynch, T.W. Schorn, and P.K.S. Siegl, *J. Med. Chem.*, 1992, **35**, 833.
7. When the reduction was carried out on the triethyl triester analogue of **7**, spontaneous intramolecular lactamization was observed.
8. (**10**): Ir (NaCl) 3430 (NH), 1740 and 1728 (C=O) cm^{-1} . ^1H Nmr 0.95 (s, 9H, CCH_3), 1.05 (t, $J = 7$ Hz, 3H, CH_3CH_2), 1.50 (s, 9H, CCH_3), 3.49 (d, $J = 5.3$ Hz, 1H, CHCO), 3.98 (q, $J = 7$ Hz, 2H, CH_3CH_2), 4.23 (dd, $J = 12$ and 6 Hz, 1H, CH-In), 4.58 (d, $J = 5.6$ Hz, 2H, $\text{CH}_2\text{CH=}$), 4.99 (dd, $J = 8.2$ and 6.4 Hz, 1H, CHNH), 5.27 (s, 2H, CH_2Ph), 5.20-5.40 (m, 2H, CH=CH_2), 5.90 (m, 1H, CH=CH_2), 7.0-7.4 (m, 9H, Ar-H), 7.76 (m, 1H, In-4H). ^{13}C Nmr 13.8 (CH_3), 27.1 and 27.8 (CCH_3), 37.9 (CH-In), 50.1 (CH_2Ph), 56.9 and 57.2 (CHNH and CHCO), 61.2 (CH_2CH_3), 65.7 ($\text{CH}_2\text{CH=}$), 81.2 and 81.9 ($\text{C}(\text{CH}_3)_3$), 109.6 (In-C7), 117.7 (In-C3a), 119.4 (In-C4), 119.6 (In-C5), 121.9 (In-C6), 126.8, 127.3, 127.6, 128.4, 128.6, 132.6, 135.8 (In-C7a), 137.1, 155.4 (CONH), 166.6 and 167.5 (COO), 171 (COOEt). Ms (EI) m/z (%) 620 (M^+ , 2), 562 (6), 491 (12), 434 (59), 278 (100), 91 (99), 57 (65). Anal. Calcd for $\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_8$: C, 67.72; H, 7.14; N, 4.51. Found: C, 67.82; H, 7.13; N, 4.26.
9. When the hydrolysis of triester (**10**) was assayed directly with 5% aqueous KOH, heating was necessary, and a diastereomeric mixture of the *threo* and *erythro* diacids (**11**) was obtained.
10. (**2**): Ir (NaCl) 1720 (C=O), 1650 cm^{-1} (C=C). ^1H Nmr 3.00-3.40 (m, 2H, CH_2CO), 3.95 (m, 1H, CH-In), 4.23 (d, $J = 7$ Hz, 2H, $\text{CH}_2\text{CH=}$), 4.68 (d, $J = 6$ Hz, 1H, CHNH), 5.00-5.50 (m, 4H, $\text{CH}_2=\text{CH}$ and CH_2Ph), 5.80 (m, 1H, $\text{CH}_2=\text{CH}$), 6.90-7.30 (m, 9H, Ar-H), 7.60 (d, $J = 8$ Hz, 1H, In-4H). ^{13}C Nmr 31.8 (CHIn), 37.2 (CH_2CO), 49.9 (CH_2Ph), 55.8 (CHNH), 65.8 ($\text{CH}_2\text{CH=}$), 110.1 (In-C3a), 111.2 (In-C7), 117.4 (CH=CH_2), 118.5 (In-C4), 119.6 (In-C5), 122.14 (In-C6), 126.6, 127.3, 127.5, 128.5, 128.6 and 132.1 (Ph), 137.0 (In-C7a), 156.1 (CONH), 173.1 (CO), 174.2 (CO). Ms (EI) m/z (%) 418 (M^+ , 7), 334 (14), 278 (58), and 91 (100). Ms (CI) 447 (M^++29) 432 (M^++14), 419 (M^++1).

11. J. Royer, H.-P. Husson, *Heterocycles*, 1993, **36**, 1493.
12. (**12**): $[\alpha]_D = -22.4$ ($c = 0.8$, CHCl_3), Ir (NaCl) 1744 and 1696 (CO), 1539 ($\text{C}=\text{C}$) cm^{-1} . $^1\text{H Nmr}$ 2.01 (s, 3H, COCH_3), 2.75 (dd, $J = 17.5$ and 4 Hz, 1H, 5-H), 3.21 (dd, $J = 17.5$ and 8.5 Hz, 1H, 5-H), 3.88 (m, 1H, 4-H), 4.24 (dd, $J = 11.5$ and 4.5 Hz, 1H, $\beta\text{-H}_A$), 4.38 (dd, $J = 11.5$ and 7.5 Hz, 1H, $\beta\text{-H}_B$), 4.59 (d, $J = 3$ Hz, 1H, 3-H); 4.68 (dd, $J = 5.5$ and 1 Hz, 2H, $\text{CH}_2\text{CH=}$), 5.22 (m, 1H, $\text{CH}_2=\text{CH}$), 5.24 (s, 2H, CH_2Ph), 5.34 (m, 2H, $\alpha\text{-H}$ and $\text{CH}_2=\text{CH}$), 5.83 (m, 1H, $\text{CH}=\text{CH}_2$), 6.95 (s, 1H, In-2H), 7.00-7.30 (m, 13H, Ar-H) and 7.55 (d, $J = 8$ Hz, 1H, In-4H). $^{13}\text{C Nmr}$ 20.8 (COCH_3), 32.5 (C-4), 38.3 (C-5), 50.1 (CH_2Ph), 52.9 (C-3), 65.9 (C- β), 67.0 (C- α), 67.6 ($\text{CH}_2\text{CH=}$), 110.3 (In-C7), 118.6 ($\text{CH}=\text{CH}_2$), 119.2 (In-C4), 119.9 (In-C5), 122.7 (In-C6), 125.0 (In-C2), 126.7, 126.8, 127.8, 128.1, 128.8, 130.9 ($\text{CH}=\text{CH}_2$), 151.5 (CONH), 169.6 (CO), 171.4 (CO) and 173.1 (CO). Ms (EI) m/z (%) 579 (M^+ , 4), 418 (4), 233 (7), 331 (8), 521 (11), 289 (13), 260 (35), 91 (100). Anal. Calcd for $\text{C}_{34}\text{H}_{33}\text{N}_3\text{O}_6$: C, 70.45; H, 5.74; N, 7.25. Found: C, 70.38; H, 5.65; N, 7.15.
13. The diastereomer of compound **12** was detected in the crude reaction mixture by nmr, but we could never isolate it.
14. W.N. Speckamp and H. Hiemstra, *Tetrahedron*, 1985, **41**, 4367.
15. (**1**): $[\alpha]_D = -19.6$ ($c = 0.5$, CHCl_3), $\text{Ir (CHCl}_3\text{)}$ 3450-3200 (OH), 1750, 1700 and 1620 (CO) cm^{-1} . $^1\text{H Nmr}$ (500 MHz) 1.62 (s, 3H, CH_3CO), 2.19-2.28 (m, 1H, 5-H), 2.70 (br t, $J = 10.5$ Hz, 1H, 5-H), 2.77 (dd, $J = 10.5$ and 7 Hz, 1H, $\beta\text{-H}_A$), 3.57 (td, $J = 10.5$ and 6.5 Hz, 1H, 6-H), 3.66 (dd, $J = 10.5$ and 2.5 Hz, 1H, $\beta\text{-H}_B$), 3.84-3.98 (m, 2H, 4-H and 6-H), 4.50-4.60 (m, 3H, $\alpha\text{-H}$ and $\text{CH}_2\text{CH=}$), 4.68 (d, $J = 8$ Hz, 1H, 3-H), 5.26 and 5.35 (2d, $J = 14$ Hz, 1H each, CH_2Ph), 5.00-5.50 (m, 2H, $\text{CH}_2=\text{CH}$), 5.88-5.98 (m, 1H, $\text{CH}_2=\text{CH}$), 6.78 (br s, 2H, Ph-H), 7.01 (s, 1H, In-2H), 7.11-7.31 (m, 10H, Ar-H), 7.34 (d, $J = 7$ Hz, 1H, In-7H), 7.62 (d, $J = 7$ Hz, 1H, In-4H). $^{13}\text{C Nmr}$ 20.5 (CH_3), 27.3 (C-5), 39.0 (C-4), 45.8 (C-6), 50.1 (CH_2Ph), 51.6 (C-3), 63.9 (C- α), 64.9 (C- β), 65.9 ($\text{CH}_2\text{CH=}$), 110.6 (In-C7), 117.3 ($\text{CH}_2=\text{CH}$), 118.2 (In-C4), 119.9 (In-C5), 122.4 (In-C6), 126.1 (In-C2), 126.5, 126.9, 127.6, 127.7, 128.5, 128.5 and 128.6 (Ar and CH=), 169.5, 169.6 and 170.2 (CO). Ms (EI) m/z (%) 565 (M^+ , 8), 374 (8), 359 (30), 318 (41), 246 (15), 163 (9), 91 (100). Anal. Calcd for $\text{C}_{34}\text{H}_{35}\text{N}_3\text{O}_5$: C, 72.18; H, 6.24; N, 7.43. Found: C, 72.21; H, 5.98; N, 7.21.

Received, 17th October, 1995