

View Article Online View Journal

RSC Advances

This article can be cited before page numbers have been issued, to do this please use: Q. liu, K. Cai and Z. Zhao, *RSC Adv.*, 2015, DOI: 10.1039/C5RA11089K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This Accepted Manuscript will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/advances

Synthesis, Structure and Catalysis of NHC-Pd(II) Complex Based on Tetradentate Mixed Ligand

Qing-Xiang Liu*, Kang-Qing Cai, Zhi-Xiang Zhao

Tianjin Key Laboratory of Structure and Performance for Functional Molecules; Key Laboratory of Inorganic-Organic Hybrid Functional Material Chemistry, Ministry of Education; College of Chemistry, Tianjin Normal University, Tianjin 300387, China. * Corresponding author, E-mail: tjnulqx@163.com

(S)-2,2'-bis[2"-(N-picolyl-benzimidazoliumyl)ethoxy]-1,1'-binaphthyl

Bis-benzimidazolium

salt

hexafluorophosphate $[(S)-LH_2]\cdot(PF_6)_2$ and its NHC palladium(II) complex $[(S)-LPd](PF_6)_2$ (1) have been prepared and characterized. Complex 1 is formed by one tetradentate mixed ligand (S)-L and one Pd(II) ion, in which one 15-membered ring and two 6-membered rings are contained. In this complex, the intramolecular π - π interactions between naphthalene rings and benzimidazole rings are observed. Additionally, the catalytic activity of complex 1 in three types of C-C coupling reactions (Suzuki-Miyaura, Heck-Mizoroki and Sonogashira reactions) was investigated. The results show that complex 1 is an effective catalyst in these coupling reactions.

Keywords: N-heterocyclic carbene, palladium(II) complex, structure, catalysis

Introduction

N-Heterocyclic carbenes (NHCs) have become a type of important ligands of organometallic chemistry and catalysis.¹ NHC ligands can form strong carbon-metal bonds with transition metals due to the strong σ -donating ability of carbene carbon.² So far, numerous successful methods have been developed to synthesize NHC complexes with different metal centers. Among these NHC-metal complexes, NHC palladium(II) complexes play a very important role in the N-heterocyclic carbene

chemistry, ^{3, 4} and they have turned out to be efficient catalysts for some C-C coupling reactions, such as Suzuki-Miyaura, Heck-Mizoroki and Sonogashira reactions.⁵ Compared with palladium(II) complexes based on phosphine ligands, NHC palladium(II) complexes have better stability to air and moisture and catalytic activity.⁶ Some NHC palladium(II) complexes bearing monodentate or polydentate ligands have been prepared and studied, and they show good catalytic activity in the construction of C-C, C-N and other C-heteroatom bonds.^{4, 7-9} In order to improve further catalytic activity of NHC palladium(II) complexes, some heteroatoms (P,¹⁰ N,^{10a, 11} or O¹²) were introduced into NHC ligands. The research results show that the catalytic activity of palladium(II) complexes bearing mixed ligands of NHC and heteroatoms can be improved remarkably, and the reason may be that strong coordinating NHC and weak coordinating heteroatoms easily form catalytically active species with vacant coordination sites. Therefore, these NHC palladium(II) complexes bearing mixed ligands may have potential applications in organic reactions.¹³

We are interested in NHC palladium(II) complexes with mixed ligands. Herein, we report the preparation of bis-benzimidazolium salt (S)-2,2'-bis[2''-(N-picolyl-benzimidazoliumyl)ethoxy]-1,1'-binaphthyl hexafluorophosphate $[(S)-LH_2]\cdot(PF_6)_2$, as well as the preparation and structure of N-heterocyclic carbene palladium(II) complex $[(S)-LPd](PF_6)_2$ (1). Additionally, the catalytic activity of complex 1 in three types of C-C coupling reactions (Suzuki-Miyaura, Heck-Mizoroki and Sonogashira reactions) was investigated.

Results and discussion

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41

Synthesis and characterizations of precursor $[(S)-LH_2]\cdot(PF_6)_2$. As shown in Scheme 1, the (S)-2,2'-dihydroxy-1,1'-binaphthyl as starting material was reacted with 2-chloroethanol to afford (S)-2,2'-di(2"-hydroxyethoxy)-1,1'-binaphthyl, followed by chlorination of hydroxyl groups with thionyl chloride to form (S)-2,2'-di(2"-chloroethoxy)-1,1'-binaphthyl, which was reacted with N-picolyl-benzimidazole in toluene to afford the bis-benzimidazolium salt $[(S)-LH_2]\cdot Cl_2$, and subsequent anion exchange with ammonium hexafluorophosphate



hexafluorophosphate ([(S)-LH₂]·(PF₆)₂). This precursor is stable toward air and moisture, and soluble in organic solvents, such as CH₂Cl₂, CH₃CN and DMSO, however, the solubility is poor in water, petroleum ether and diethyl ether. In the ¹H NMR spectra of [(S)-LH₂]·(PF₆)₂, the proton signals (NC*H*N) of benzimidazolium appear at $\delta = 9.43$ ppm, which is consistent with the chemical shifts of reported benzimidazolium salts.¹⁴



Scheme 1 Preparation of precursor [(S)-LH₂]·(PF₆)₂ and complex 1

Synthesis, characterization and structure of complex 1. As shown in Scheme 1, complex $[(S)-LPd](PF_6)_2$ (1) was prepared via the reaction of $[(S)-LH_2]\cdot(PF_6)_2$ with Pd(OAc)₂ in the presence of K₂CO₃ in acetonitrile/ethylene glycol solution. The single crystals of complex 1 suitable for X-ray diffraction were grown by slow diffusion of diethyl ether into its solution. Complex 1 is stable to heat, air and moisture, soluble in CH₃CN and DMSO, but scarcely soluble in diethyl ether and hydrocarbon solvents. In the ¹H NMR spectra of 1, the disappearance of the resonances for the benzimidazolium protons (NCHN) shows the formation of the expected metal carbene complexes, and the chemical shifts of other hydrogen atoms are similar to those of corresponding precursor. In ¹³C NMR spectra, the signals for the carbene carbon of 1 appear at $\delta = 179.2$ ppm, which are similar to known metal carbene complexes.¹⁵

Complex 1 is formed by one tetradentate mixed ligand (S)-L and one Pd(II) ion, in which one 15-membered ring and two 6-membered rings are contained (Fig. 1). The Pd(II) ion is tetra-coordinated with two carbene carbon atoms and two nitrogen atoms from two pyridine rings to adopt a square-planar geometry. The bond distances of Pd-C and Pd-N are 1.966(3)-1.976(3) Å and 2.095(3)-2.098(3) Å, respectively. The bond angles of C(7)-Pd(1)-C(38) and N(1)-Pd(1)-N(6) are 96.0(1)° and 94.4(1)°, respectively. The bond angles of C-Pd-N are from 84.9(1)° to 175.0(1)°. These values are similar to those of reported NHC-Pd(II) complexes.¹⁶ The axial chirality of complex 1 is consistent with corresponding precursor because its chiral binaphthyl unit (S configuration) was not changed during the preparation of metal complex. In this complex, two naphthalene rings form the dihedral angle of $69.2(3)^{\circ}$, and the dihedral angle between two benzimidazole rings is 87.2(2)°. The dihedral angle between benzimidazole ring and adjacent pyridine ring is $71.5(3)^{\circ}$, and the dihedral angle between two pyridine rings is $50.7(4)^{\circ}$. The internal ring angles (N-C-N) at the carbene centers are $107.1(3)^{\circ}$ and $107.6(3)^{\circ}$, and these values are similar to those of known NHC-metal complexes.¹⁵ Besides, the intramolecular π - π interactions between the naphthalene rings and the benzimidazole rings are observed¹⁷ with the face-to-face distance of 3.405(4) Å and the center-to-center distance of 3.858(3) Å.

In the crystal packing of **1**, 2D supramolecular layer is formed through three types of $\pi \cdots \pi$ stacking interactions as shown in Fig. S1. In three types of $\pi \cdots \pi$ stacking interactions, one is from intermolecular pyridine rings with the face-to-face distance of 3.761(7) Å and the center-to-center distance of 3.773(2) Å, and another is from intermolecular benzimidazole rings with the face-to-face distance of 3.654(5) Å and the center-to-center distance of 3.659(3) Å, and third is from naphthalene rings and pyridine rings with the face-to-face distance of 3.534(5) Å and the center-to-center distance of 3.901(3) Å.



Fig. 1. Perspective view of **1**. All hydrogen atoms were omitted for clarity. Selected bond lengths (Å) and angles (°): C(38)-Pd(1) 1.966(3), C(7)-Pd(1) 1.976(3), N(6)-Pd(1) 2.095(3), N(1)-Pd(1) 2.098(3), O(1)-C(15) 1.409(4), O(2)-C(36) 1.432(4); N(4)-C(38)-N(5) 107.6(3), N(2)-C(7)-N(3) 107.1(3), C(38)-Pd(1)-C(7) 96.0(1), C(38)-Pd(1)-N(6) 84.9(1), N(1)-Pd(1)-N(6) 94.4(1), C(7)-Pd(1)-N(1) 85.2(1), C(7)-Pd(1)-N(6) 172.3(1), C(38)-Pd(1)-N(1) 175.0(1).

Catalytic activity of NHC-Pd(II) complex 1 in the Suzuki-Miyaura reaction. The Suzuki-Miyaura reaction, a cross-coupling process between an organoboron reagent and an aryl halide, has become a successful and useful tools for forming C-C bonds.¹⁸ In order to examine the catalytic activity of NHC-Pd(II) complex **1** in the Suzuki-Miyaura reaction, we chose the cross-coupling reaction of 4-bromotoluene with phenylboronic acid as a model reaction in the air to test the effects of solvent and base (Table S1 in Supplementary Information). Use of K₂CO₃ as base and water as solvent gave 97% coupling yield at 40 °C over the course of 12 h in the presence of air (Table S1, Entry 1), whereas use of CH₃OH/H₂O (1:1) as solvent gave 98% yield in 6 h (Table S1, Entry 2). Use of K₂CO₃ as base and other solvents (such as MeOH, C₂H₅OH, 1,4-dioxane, THF or C₂H₅OH/H₂O (1:1)) gave yields of 7-80% (Table S1, Entries 3-7). Additionally, use of water as solvent and other common bases (such as 'BuOK, K₃PO₄·3H₂O and NaOAc) gave yields of 52-74% (Table S1, Entries 8-10). Under conditions similar to those of entry 1 (but the time being shorten from 12 h to 6 h), 10 mol% of TBAB were added to afford 99% yield (Table S1, Entry 11). The

result indicates that the reaction could be effectively improved in the presence of TBAB. It has been known that the addition of TBAB as a co-catalyst can enhance the activity of many catalytic systems. Besides, TBAB is able to activate and stabilize palladium(0) species generated in situ, which is believed to be the real catalytically active species.¹⁹ Based on this extensive screening, water and K₂CO₃ were found to be the efficient solvent and base in the presence of 10 mol% TBAB with a catalyst loading of 0.1 mol% of complex **1** at 40 °C in the air, and these conditions proved ideal. Reactions were monitored by GC at appropriate time intervals. Althouth the good yield can be also obtained using CH₃OH/H₂O (1:1) as solvent, we chose water as solvent for the purpose of environmental benignancy.

We attempted cross-coupling reactions of various aryl halides with phenylboronic acids under the optimal reaction conditions. The catalyst is highly efficient toward the coupling of aryl bromides (or aryl dibromides) with phenylboronic acid to afford excellent yields of 90-98% (Table 1, 2a, 2b, 2c(1)-2e(1) and 2f-2i, 2n(2)). The catalyst is also effective toward the coupling of aryl chlorides bearing electronic-donating or -withdrawing substituents with phenylboronic acid to give moderate to high yields of 64-89% (Table 1, 2c(2)-2e(2) and 2j-2m). Coupling reaction of chlorobenzene with phenylboronic acid gave relatively poor yields of 43% (Table 1, 2n(1)). The coupling of iodobenzene with phenylboronic acid gave quantitative yield (Table 1, 2n(3)).

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41

In order to further study the influence of the ligand on catalysis, the control experiment was performed using 4-bromotoluene and phenylboronic acid as substrates, in which $Pd(OAc)_2$ (or $PdCl_2$) and TBAB were added in the absence of the ligand with the optimized conditions. Only 39% for $Pd(OAc)_2$ and 32% for $PdCl_2$ desired coupling product were obtained, and these values are smaller than 99% yield in the case of complex **1** as catalyst. This result highlights the preponderant catalytic role of the ligand in these aqueous and aerobic coupling reactions.

6

Table 1 Suzuki-Miyaura Reaction of aryl halides with phenylboronic acid catalyzedby complex 1^[a]



- [a] Reaction conditions: Aryl halide (0.5 mmol), phenylboronic acid (0.6 mmol), K₂CO₃ (1.2 mmol), complex 1 (0.1 mol %), TBAB (10 mol %), H₂O (6 mL) at 40 °C in the air.
- [b] Reactions were monitored by TLC.
- [c] Isolated yield using aryl halide as a standard.

The above results show that complex **1** has high stability to both air and water, while also demonstrating excellent tolerance of various functional groups. Additionally, complex **1** can resolve partly in aqueous medium due to the existence of hydrophilic groups (such as oxygen atom and nitrogen atom). These qualities make **1** a valuable precatalyst for thermally sensitive substrates. This provides an effective phosphine-free catalyst system for facile coupling of aryl halides with aryl boronic acids in water under mild and aerobic conditions. These favorable reaction conditions of Suzuki-Miyaura reaction are important toward ensuring environmental responsibility while also reducing costs for fine chemical and pharmaceutical industries.

Catalytic activity of NHC-Pd(II) complex 1 in the Heck-Mizoroki reaction. The Heck reaction has evolved significantly from its original mode as the arylation of olefins with aryl iodides, and it is a powerful tool for the coupling of less reactive bromides, chlorides and pseudo-halides such as triflates, tosylates, mesylates, and aryl diazonium salts.²⁰ We chose bromobenzene and styrene as the coupling partners to test the solvent and base effects in the air (Table S2). Using 1,4-dioxane as solvent and NaOAc as base gave 92% coupling yields at 120 °C over the course of 24 h in the presence of air (Table S2, Entry 1), whereas using other bases (such as K_2CO_3 , K₃PO₄·3H₂O, Et₃N, ^tBuOK, KOH, NaHCO₃ and Cs₂CO₃) gave the yields of 15-83% (Table S2, Entries 2-8). Using NaOAc as base and other solvents (such as H_2O , DMF, CH₃CN, THF, DME or DMSO) gave the yields from trace to 61% (Table S2, Entries 9-15). Under conditions similar to those of entry 1 (but temperature being dropped from 120 °C to 100 °C), 10 mol% of PEG-400 was added to give coupling yields of 95% (Table S2, Entry 16). This result shows that the reaction could be dramatically improved in the presence of PEG-400. It has been known that the addition of PEG-400 as a co-catalyst can enhance the activity of many catalytic systems.²¹ Based on this extensive screening, 1,4-dioxane and NaOAc were found to be the efficient solvent and base in the presence of 10 mol% of PEG-400 as co-catalyst with a catalyst

8

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41

RSC Advances

loading of 0.1 mol% of complex **1** at 100 °C in the air, thus giving the optimal coupling result.

We attempted cross-coupling reactions of various aryl halides with styrene under the optimal reaction conditions. The cross-coupling of various aryl bromides with styrene gave good to excellent yields of 80-98% (Table 2, 3a, 3b, 3c(1)-3e(1) and **3f-3i**). The coupling of 4-chloronitrobenzene or 2-chloronitrobenzene with styrene gave good yields of 85% or 80%, respectively (Table 2, 3d(2) and 3j). The coupling of 4-chloroaniline. 4-chloroacetophenone, 2,4-dinitrochlorobenzene, 2-chlorobenzaldehyde or chlorobenzene with styrene gave relatively poor yields of 20-56% (Table 2, 3c(2), 3e(2), 3k, 3l and 3n(1)). And 2-chlorotoluene even gave trace coupling yield (Table 3: 3m). The catalyst is highly efficient in the coupling of iodobenzene with styrene to give almost quantitative yield (Table 2, 3n(3)). Notably, there is high selectivity for the *trans* product in this reaction system. Hardly any *cis* products could be detected after the reaction. The *trans* isomer appears to be the dominant conformation assumed during the course of these reactions.

Considering all these facts, complex **1** is an effective catalyst for Heck-Mizoroki reaction. In order to further study the influence of the ligand on catalysis, the control experiment was performed using bromobenzene and styrene as substrates, in which $Pd(OAc)_2$ (or $PdCl_2$) and PEG-400 were added in the absence of the ligand with the optimized conditions. Only 26% for $Pd(OAc)_2$ and 21% for $PdCl_2$ coupling product were obtained, and these values are smaller than 95% yield in the case of complex **1** as catalyst. This result highlights the preponderant catalytic role of the ligand in the coupling reactions.



Table 2 Heck-Mizoroki reaction of aryl halides with styrene catalyzed by complex 1^[a]

- [a] Reaction conditions: aryl halide (0.5 mmol), styrene (0.6 mmol), K₂CO₃(1.2 mmol), complex 1 (0.1 mol%), PEG-400 (10 mol%), 1,4-dioxane (6 mL), 100 °C in air.
- [b] Monitored by TLC.

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41.

[c] Isolated yield using aryl halide as a standard.

RSC Advances

Catalytic activity of NHC-Pd(II) complex 1 in the Sonogashira reaction. In order to prove that complex 1 would be a good precatalyst for the Sonogashira reaction, we chose the cross-coupling reaction of 4-bromoanisole with phenylacetylene as a model reaction to test the solvent, base and catalyst activity in the air.²² As illustrated in Table S3, using K_3PO_4 ·3H₂O as base and 1,4-dioxane/H₂O (1:1) as solvent gave 79% yield at 100 °C over the course of 24 h in the presence of air (Table S3, entry 1). In the same conditions, when 10 mol% of TBAB was added, the yield rose to 93% (Table S3, entry 2). Using Cs₂CO₃ as base and other solvents (such as 1,4-dioxane, DMF, DMSO, CH₃CN, H₂O, THF, DMF/H₂O (1:1) or 1,4-dioxane/H₂O (1:1)) gave yields from trace to 69% (Table S3, Entries 3-10). Using 1,4-dioxane/H₂O (1:1) as solvent in the presence of 10 mol% of PPh₃ or TBAB, the different bases (such as K_2CO_3 , KOH, NaOAc, Et₃N or NaHCO₃) were investigated to give the yields of 70-91% (Table S3, Entries 11-17). Based on this extensive screening, 1,4-dioxane/H₂O (1:1) and $K_3PO_4 \cdot 3H_2O$ were proved to be the efficient solvent and base in the presence of 10 mol% of TBAB with a catalyst loading of 0.1 mol% of complex 1, thus giving the optimal coupling result.

We attempted the cross-coupling reactions of various aryl halides with phenylacetylene under the optimized conditions. Various aryl bromides could be easily coupled to phenylacetylene to give the good to excellent yields of 86-96% (Table 3, 4a, 4b, 4c(1)-4e(1), 4f-4i and 4n(2)). Coupling reactions of 4-chloroaniline, 4-chloronitrobenzene or 2-chloronitrobenzene with phenylacetylene gave good yields 4c(2), 4d(2) and 4j). While the couplings of of 76-83% (Table 3. 2,4-dinitrochlorobenzene, 4-chloroacetophenone, 2-chlorobenzaldehyde, 2-chlorotoluene or chlorobenzene with phenylboronic acid gave poor yields of 10-45% (Table 3, 4e(2), 4k-4m and 4n(1)). The catalyst was highly efficient in the coupling of iodobenzene with phenylacetylene to give almost quantitative yield (Table 3, 4n(3)). The above results show complex 1 is an effective precatalyst for Sonogashira reaction. Additionally, a control experiment was also performed using 4-bromoanisole with phenylacetylene as substrates, in which $Pd(OAc)_2$ (or $PdCl_2$) and TBAB were added without the ligand with the optimized conditions. Only 31% for $Pd(OAc)_2$ and 29% for $PdCl_2$ coupling product were obtained, and these values are smaller than 93% yield in the case of complex **1** as catalyst. This result highlights the preponderant catalytic role in the coupling reaction.

 Table 3 Sonogashira reaction of aryl halides with phenylacetylene catalyzed by

 complex 1^[a]



[a] Reaction conditions: aryl halide (0.5 mmol), phenylacetylene (0.6 mmol), base (1.2 mmol),

complex 1 (0.1 mol%), TBAB (10 mol%), 1,4-dioxane/H₂O (6 mL), 100 $^\circ C$ in air. [b] Monitored by TLC.

[c] Isolated yield using aryl halide as a standard.

It is noteworthy that the good catalytic activity of NHC-Pd(II) complex **1** can be ascribed to the strong donating ability of NHC and the weak coordination of nitrogen. The weak coordinating abilities of nitrogen atoms allow them to readily dissociate from metal centers thus creating catalytically active species with vacant coordination sites. The strong Pd-C bond presumably facilitates oxidative addition of aryl halogens to the metal. The dynamic dissociation and association of nitrogen atoms allows the approach of substrate to the metal center, and the substituents on the benzimidazole rings favor stabilization of the active metal species after reductive elimination of the products.

Additionally, the reaction conditions (catalyst loading, temperature, solvents and bases) between the present paper and literatures are compared as shown in Table S4. The catalyst loading of complex **1** is all 0.1% in three types of C-C coupling reactions, which is same as the most case of reported data in Suzuki-Miyaura^{22(b), 23} and Heck-Mizoroki reactions,^{22(b), 23(k), 23(l), 24} whereas this value is smaller than reported data (1-4%) in Sonogashira reaction.^{22, 23m, 25} The reaction temperature for these reactions in the present paper is in the middle of reported data (in three types of C-C coupling reactions: 40 °C, 100 °C and 100 °C, respectively, in the present paper; 25-140 °C, 80-150 °C and 80-120 °C, respectively, in literatures). In the present paper, pure water, organic solvent and water/organic solvent are used, respectively, in three types of C-C coupling reactions, and the similar cases can be also found in reported results. In the use of bases, we selected inexpensive K₂CO₃, NaOAc and K₃PO₄·3H₂O, respectively, in three types of C-C coupling reactions, and the similar bases are also used in literatures.

On the whole, the results obtained using tetradentate Pd(II) complex **1** as catalyst in most Suzuki-Miyaura, Heck-Mizoroki and Sonogashira reactions are similar to or slightly better than those of related NHC Pd(II) complexes. By the analysis and comparison of complex **1** with other NHC metallic catalysts published,²⁶⁻²⁸ we can

know that the catalytic activity of a catalyst is related to a number of factors such as the steric hindrance and electronic effects of different substituents around Pd, as well as the stability and solubility of complexes. Besides, the reaction conditions are also unneglected factors.

Conclusion

In summary, bis-benzimidazolium salt $[(S)-LH_2]\cdot(PF_6)_2$ and its NHC palladium(II) complex 1 have been prepared and characterized. In the crystal structure of complex 1, one 15-membered ring and two 6-membered rings are contained. Additionally, the catalytic activity of the complex 1 in three types of C-C coupling reactions (Suzuki-Miyaura, Heck-Mizoroki and Sonogashira reactions) was studied. The results show that complex 1 can smoothly catalyze most Suzuki-Miyaura reactions of various aryl halides (iodide, bromide and chloride) with arylboronic acids using water as solvent and K₂CO₃ as base in the air at 40 °C to afford moderate to high yields. Likewise, most Heck-Mizoroki or Sonogashira reactions can also be performed efficiently at 100 °C in the air.

Compared with reported NHC Pd(II) complexes, complex 1 shows similar or slightly better catalytic activity in these coupling reactions. Complex 1 may be a valuable precatalyst for the preparation of diaryl derivatives, styrene derivatives or phenylacetylene derivatives. Further studies of new organometallic complexes from precursor $[(S)-LH_2]\cdot(PF_6)_2$ as well as analogous ligands are underway.

Experimental section

General procedures. N-Picolyl-benzimidazole was prepared according to the literature method.²⁹ All the reagents for synthesis and analyses were of analytical grade and used without further purification. Melting points were determined on an Digital Vision MP Instrument. ¹H and ¹³C NMR spectra were recorded at 400 MHz and 100 MHz, respectively. Chemical shifts, δ , are reported in ppm relative to the internal standard TMS for both ¹H and ¹³C NMR. *J* values are given in Hz. The elemental analyses of all compounds were obtained from the powder compounds

RSC Advances

recrystallised. GC analyses were carried out with a capillary column (CBPī or CBP-5, $0.25 \text{ mm i.d} \times 40 \text{ m}$).

Preparation of (S)-2,2'-di(2"-hydroxyethoxy)-1,1'-binaphthyl. A DMF (100 mL) suspension of (S)-2,2'-dihydroxy-1,1'-binaphthyl (5.000 g, 17.4 mmol) and potassium carbonate (9.646 g, 69.9 mmol) was stirred for 1 h at 50 °C. Then 2-chloroethanol (5.628 g, 69.9 mmol) was added dropwise, and stirred continually for 24 h under refluxing. The solvent was removed under vacuum, and 500 mL water was added. The solution was extracted with CH_2Cl_2 (3 × 30 mL) and the extracting solution was dried over anhydrous MgSO₄. After removing CH₂Cl₂, a pale yellow oil was obtained, and the crude product was recrystallized with MeOH to give (S)-2,2'-di(2"-hydroxyethoxy)-1,1'-binaphthyl as a white powder. Yield: 6.008 g (92%). M.p.: 112-114 °C. Anal. Calcd for C₂₄H₂₂O₄: C, 76.98; H, 5.92%. Found: C, 76.63; H, 5.88%. ¹H NMR (400 MHz, CDCl₃): δ 8.02 (d, J = 8.8 Hz, 2H, ArH), 7.93 (d, J = 8.0 Hz, 2H, ArH), 7.48 (d, J = 9.2 Hz, 2H, ArH), 7.41 (t, J = 7.4 Hz, 2H, ArH), 7.30 (t, J = 7.4 Hz, 2H, ArH), 7.17 (d, J = 8.4 Hz, 2H, ArH), 4.28 (m, 2H, CH₂), 4.08 (m, 2H, CH₂), 3.66 (m, 4H, CH₂), 2.31 (s, 2H, OH). ¹³C NMR (100 MHz, CDCl₃): δ 153.5 (PhC), 133.8 (PhC), 129.8 (PhC), 129.6 (PhC), 128.1 (PhC), 126.7 (PhC), 125.2 (PhC), 124.2 (PhC), 120.3 (PhC), 115.9 (PhC), 71.7 (OCH₂CH₂), 61.2 (OCH₂CH₂).

Preparation of (S)-2,2'-di(2''-chloroethoxy)-1,1'-binaphthyl. To a chloroform (100 mL) solution of (S)-2,2'-di(2''-hydroxyethoxy)-1,1'-binaphthyl (5.000 g, 13.4 mmol) and pyridine (5.288 g, 66.8 mmol), thionyl chloride (7.947 g, 66.8 mmol) was added dropwise at room temperature within 1 h. Then the solution was stirred continually for 12 h at 70 °C. The mixture was cooled and washed with water (3 × 100 mL). The organic layer was dried over anhydrous MgSO₄. After removing chloroform, (S)-2,2'-di(2''-chloroethoxy)-1,1'-binaphthyl was obtained as a pale yellow powder. Yield: 3.571 g (65%). M.p.: 101-103 °C. Anal. Calcd for C₂₄H₂₀O₂Cl₂: C, 70.08; H, 4.90%. Found: C, 70.37; H, 5.14%. ¹H NMR (400 MHz, CDCl₃): δ 7.97 (d, *J* = 8.8 Hz, 2H, Ar*H*), 7.88 (d, *J* = 8.4 Hz, 2H, Ar*H*), 7.43 (d, *J* = 8.8 Hz, 2H, Ar*H*), 7.37 (t, *J* =

7.4 Hz, 2H, Ar*H*), 7.25 (t, J = 7.6 Hz, 2H, Ar*H*), 7.13 (d, J = 8.4 Hz, 2H, Ar*H*), 4.22(m, 4H, C*H*₂), 3.40 (m, 4H, C*H*₂). ¹³C NMR (100 MHz, CDCl₃): δ 153.6 (PhC), 147.9 (PhC), 134.0 (PhC), 129.6 (PhC), 127.9 (PhC), 126.5 (PhC), 125.4 (PhC), 124.1 (PhC), 121.0 (PhC), 116.3 (PhC), 70.1 (OCH₂CH₂), 41.7 (OCH₂CH₂).

Preparation $[(S)-LH_2] \cdot (PF_6)_2$. of А toluene (35)mL) solution of N-picolvl-benzimidazole (1.274)6.1 mmol) and g, (S)-2,2'-di(2"-chloroethoxy)-1,1'-binaphthyl (1.000 g, 2.4 mmol) was stirred for 7 days under refluxing, brown oil of and а (S)-2,2'-bis[2''-(N-picolyl-benzimidazoliumyl)ethoxy]-1,1'-binaphthyl chloride was obtained after the solvent was removed via a rotary evaporator. Then NH_4PF_6 (0.938 g, 5.7 (100)mmol) added methanol solution ml) of was to a (S)-2,2'-bis[2''-(N-picolyl-benzimidazoliumyl)ethoxy]-1,1'-binaphthyl chloride (2.016 g, 2.4 mmol) with stirring for 3 days, and a white precipitate was formed. The precipitate was collected by filtration and washed with a small portion of methanol to give [(S)-LH₂]·(PF₆)₂. Yield: 1.734 g (68%). M.p.: 148-150 °C. Anal. Calcd for C₅₀H₄₂F₁₂N₆O₂P₂: C, 57.25; H, 4.03; N, 8.01%. Found: C, 57.62; H, 4.41; N, 8.35%.¹H NMR (400 MHz, DMSO- d_6): δ 9.43 (s, 2H, 2-bimiH), 8.43 (d, J = 4.4 Hz, 2H, ArH), 7.98 (d, J = 8.8 Hz, 2H, ArH), 7.90 (q, J = 2.6 Hz, 4H, ArH), 7.77 (d, J = 8.4 Hz, 2H, ArH), 7.63 (d, J = 8.0 Hz, 2H, ArH), 7.50 (t, J = 7.8 Hz, 2H, ArH), 7.43 (d, J = 9.6 Hz, 2H, ArH), 7.37 (q, J = 4.0 Hz, 2H, ArH), 7.28 (t, J = 8.4 Hz, 4H, ArH),7.21 (t, J = 7.8 Hz, 2H, ArH), 6.99 (m, J = 5.0 Hz, 2H, ArH), 6.58 (d, J = 8.4 Hz, 2H, ArH), 5.68 (m, 3H, CH₂), 4,63 (t, J = 4.6 Hz, 3H, CH₂), 4.38 (t, J = 4.4 Hz, 3H, CH₂), 4.15 (q, J = 5.2 Hz, 3H, CH₂). ¹³C NMR (100 MHz, DMSO- d_6): δ 153.1 (ArC), 149.9 (ArC), 142.9 (ArC), 138.0 (ArC), 132.9 (bimi-NCN), 131.1 (ArC), 130.9 (ArC), 130.0 (ArC), 129.3 (ArC), 128.4 (ArC), 126.8 (ArC), 126.7 (ArC), 126.6 (ArC), 126.4 (ArC), 125.5 (ArC), 125.0 (ArC), 124.2 (ArC), 123.2 (ArC), 119.6 (ArC), 116.3 (ArC), 114.0 (ArC), 113.0 (ArC), 67.8 (OCH₂CH₂), 51.2 (OCH₂CH₂), 49.0 (CH₂), $47.2 (CH_2) (bimi = benzimidazole).$

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41

Prepartion of [(S)-LPd](PF₆)₂ (1). An acetonitrile/ethylene glycol (20 mL, v:v = 1:1) suspension of K₂CO₃ (0.179 g, 1.3 mmol), **[(S)-LH₂]·(PF₆)₂** (0.209 g, 0.2 mmol) and Pd(OAc)₂ (0.042 g, 0.2 mmol) was stirred for 24 h under 60 °C. The resulting solution was filtered and concentrated to 10 mL, and Et₂O (5 mL) was added to precipitate **1** as a yellow powder. Yield: 0.057 g (25%). M.p.: 280-282 °C. Anal. Calcd for C₅₀H₄₀F₁₂N₆O₂P₂Pd: C, 52.07; H, 3.49; N, 7.29%. Found: C, 52.34; H, 3.63; N, 7.52%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.24 (m, 7H, Ph*H*), 7.98 (m, 4H, Ph*H*), 7.80 (d, *J* = 8.4 Hz, 2H, Ph*H*), 7.65 (t, *J* = 6.6 Hz, 2H, Ph*H*), 7.44 (m, 7H, Ph*H*), 4.90 (m, 2H, CH₂), 4.52 (m, 2H, CH₂), 4.24 (m, 2H, CH₂), 2.89 (s, 3H, CH₂), 2.73 (s, 3H, CH₂). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 179.2 (bimi-NCN), 154.5 (PhC), 153.8 (PhC), 153.2 (PhC), 147.5 (PhC), 135.2 (PhC), 133.6 (PhC), 133.0 (PhC), 129.6 (PhC), 124.6 (CH₃), 112.4 (PhC), 111.7 (PhC), 68.9 (CH₂), 63.3 (CH₂), 62.9 (CH₂).

General procedure for the Suzuki-Miyaura reaction. In a typical reaction, aryl halide (0.5 mmol), arylboronic acid (0.6 mmol), base (1.2 mmol), NHC-Pd(II) complex 1 (0.1 mol%) and TBAB (10 mol%) in solvent (6 mL) were stirred at 40 °C in air. After the desired reaction time, the reaction was stopped, and water (20 mL) was added to the reaction mixture. The mixture was extracted by n-hexane (10 mL × 3), and the organic layer was washed with water (10 mL \times 3) and dried over anhydrous MgSO₄. For the determination of GC-MS yields, the above solution was filtered through a thin pad of silica gel (elution with CH₂Cl₂ (100 mL)) and concentrated to 5 mL. n-Tridecane (92 mg) was added as internal standard. The isolated products were purified by column chromatography on silica gel using a mixture of n-hexane and dichloromethane (from pure n-hexane to n-hexane/dichloromethane = 10/1) as eluant. The isolated yields were determined using aryl halides as standards.

General procedure for the Heck-Mizoroki reaction. A mixture of aryl halide (0.5 mmol), styrene (0.6 mmol), base (1.2 mmol), complex **1** (0.1 mol%) and PEG-400 (10

mol%) in solvent (6 mL) was stirred at 100 °C or 120 °C in air. After the desired reaction time, the reaction was stopped, and water (20 mL) was added to the reaction mixture. The mixture was extracted by ethyl ether (10 mL × 3), and the organic layer was washed with water (10 mL × 3) and dried over anhydrous MgSO₄. Then the solution was filtered and concentrated to about 2 mL. The isolated products were purified by column chromatography on silica gel using a mixture of n-hexane and dichloromethane (from pure n-hexane to n-hexane/dichloromethane = 10/1) as eluant. The isolated yields were determined using aryl halides as standards.

General procedure for the Sonogashira reaction. A mixture of aryl halide (0.5 mmol), phenylacetylene (0.6 mmol), base (1.2 mmol), complex **1** (0.1 mol%) and TBAB (10 mol%) in solvent (6 mL) was stirred at 100 °C in air. After the desired reaction time, the reaction was stopped, and water (20 mL) was added to the reaction mixture. The mixture was extracted by diethyl ether (10 mL × 3), and the organic layer was washed with water (10 mL × 3) and dried over anhydrous MgSO₄. Then the solution was filtered and concentrated to about 2 mL. The isolated products were purified by column chromatography on silica gel using a mixture of n-hexane and dichloromethane (from pure n-hexane to n-hexane/dichloromethane = 10/1) as eluant. The isolated yields were determined using aryl halides as standards

X-ray data collection and structure determinations. X-ray single-crystal diffraction data for complexs were collected by using a Bruker Apex II CCD diffractometer at 173(2) K for **1** with Mo-Ka radiation ($\lambda = 0.71073$ Å) by ω scan mode. There was no evidence of crystal decay during data collection in all cases. Semiempirical absorption corrections were applied by using SADABS and the program SAINT was used for integration of the diffraction profiles.³⁰ All structures were solved by direct methods by using the SHELXS program of the SHELXTL package and refined with SHELXL³¹ by the full-matrix least-squares methods with anisotropic thermal parameters for all non-hydrogen atoms on F^2 . Hydrogen atoms bonded to C atoms were placed geometrically and presumably solvent H atoms were

first located in difference Fourier maps and then fixed in the calculated sites. Further details for crystallographic data and structural analysis are listed in Table 4. Figures were generated by using Crystal-Maker.³²

 Table 4 Summary of crystallographic data complex 1

$1 \cdot 0.5 C_2 H_6 O_2$			
Chemical	$C_{50}H_{40}F_{12}N_6O_2P_2Pd$	<i>F</i> (000)	1198
formula	$\cdot 0.5C_2H_6O_2$		
Formula weight	1184.25	Cryst size, mm	$0.18\times0.17\times0.15$
Cryst syst	Triclinic	$\theta_{\min}, \theta_{\max}, \deg$	1.36, 25.01
Space group	Pī	<i>Т</i> , К	173(2)
<i>a</i> , Å	12.526(1)	No. of data collected	12770
<i>b</i> , Å	14.488(1)	No. of unique data	8625
<i>c</i> , Å	15.330(1)	No. of refined params	677
α, deg	98.5(1)	Goodness-of-fit on F^{2a}	1.016
β , deg	95.0(1)	Final <i>R</i> indices ^b $[I > 2\sigma(I)]$	
γ, deg	115.0(1)	R_1	0.0397
$V, \text{\AA}^3$	2456.2(3)	wR_2	0.1012
Ζ	2	R indices (all data)	
$D_{\rm calcd}$, Mg m ⁻³	1.601	R_1	0.0469
Abs coeff, mm ⁻¹	0.540	wR_2	0.1077

^{*a*}GOF = $[\Sigma w(F_o^2 - F_c^2)^2/(n-p)]^{1/2}$, where *n* is the number of reflection and *p* is the number of parameters refined. ^{*b*}R₁ = $\Sigma(||F_o| - |F_c||)/\Sigma|F_o|$; $wR_2 = [\Sigma[w(F_o^2 - F_c^2)^2]/\Sigma w(F_o^2)^2]^{1/2}$.

Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (No.21172172), Tianjin Natural Science Foundation (No.11JCZDJC22000) and The Program for Innovative Research Team in University of Tianjin (TD12-5038).

Supplementary information

Tables and figures for complex **1**, and ¹H NMR and ¹³C NMR spectra for precursor, complex and all coupling products in Suzuki-Miyaura, Heck-Mizoroki and Sonogashira reactions. CCDC 1023636 contains the supplementary crystallographic data for complex **1**. These data can be obtained free of charge via

http//www.ccdc.cam.ac.uk/conts/retrieving.html, or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK; fax: (+44) 1223-336-033; or e-mail: deposit@ccdc.cam.ac.uk.

References

- 1 (a) M. Melaimi, M. Soleilhavoup and G. Bertrand, Angew. Chem., Int. Ed., 2010, 49, 8810; (b) M. Poyatos, J. A. Mata and E. Peris, Chem. Rev., 2009, 109, 3677; (c) F. E. Hahn and M. C. Jahnke, Angew. Chem., Int. Ed., 2008, 47, 3122; (d) W. A. Herrmann, Angew. Chem., Int. Ed., 2002, 41, 1290; (e) L. Mercs and M. Albrecht, Chem. Soc. Rev., 2010, 39, 1903; (f) S. Budagumpi and S. Endud, Organometallics, 2013, 32, 1537; (g) W. A. Herrmann, Angew. Chem., 2002, 114, 1342; (h) S. P. Nolan, Acc. Chem. Res., 2011, 44, 91; (i) S. Gaillard, C. S. J. Cazin and S. P. Nolan, Acc. Chem. Res., 2012, 45, 778; (j) F. Wang, L. J. Liu, W. Wang, S. Li and M. Shi, Coord. Chem. Rev., 2012, 256, 804.
- 2 (a) C. Boehme and G. Frenking, J. Am. Chem. Soc., 1996, 118, 2039; (b) M. S.
 Viciu, O. Navarro, R. F. Germaneau, R. A. Kelly, W. Sommer, N. Marion, E. D.
 Stevens, L. Cavallo and S. P. Nolan, Organometallics, 2004, 23, 1629.
- 3 (a) W. N. O. Wylie, A. J. Lough and R. H. Morris, *Organometallics*, 2010, 29, 570;
 (b) A. R. Chianese, P. T. Bremer, C. Wong and R. J. Reynes, *Organometallics*, 2009, 28, 5244;
 (c) X. Q. Zhang, Y. P. Qiu, B. Rao and M. M. Luo, *Organometallics*, 2009, 28, 3093;
 (d) N. D. Clement, K. J. Cavell and L. L. Ooi, *Organometallics*, 2006, 25, 4155.
- 4 (a) N. Marion and S. P. Nolan, *Acc. Chem. Res.*, 2008, **41**, 1440; (b) J. Houghton, G. Dyson, R. E. Douthwaite, A. C. Whitwood and B. M. Kariuki, *Dalton Trans.*, 2007, **28**, 3065; (c) W. A. Herrmann, M. Elison, J. Fischer, C. Koecher and G. R. J. Artus, *Angew. Chem.*, *Int. Ed.*, 1995, **34**, 2371.
- 5 (a) N. Stylianides, A. A. Danopoulos, D. Pugh, F. Hancock and A. Zanotti-Gerosa, Organometallics, 2007, 26, 5627; (b) S. Ahrens, A. Zeller, M. Taige and T. Strassner, Organometallics, 2006, 25, 5409; (c) E. A. B. Kantchev, C. J. O'Brien and M. G. Organ, Angew. Chem., Int. Ed., 2007, 46, 2768; (d) L. G. Guang, G. Pei,

20

RSC Advances

H. Pan, Q. Cheng, C. S. Cao, Y. Peng, Z. Shi, G. S. Pang and Y. H. Shi, *Dalton Trans.*, 2012, 41, 5020; (e) A. Rit, T. Pape, A. Hepp and F. E. Hahn, *Organometallics*, 2011, 30, 334; (f) A. Suzuki, *J. Organomet. Chem.*, 2002, 653, 83; (g) A. F. Littke and G. C. Fu, *Angew. Chem.*, 2002, 41, 4176; (h) Q. X. Liu, W. Zhang, X. J. Zhao, Z. X. Zhao, M. C. Shi and X. G. Wang, *Eur. J. Org. Chem.*, 2013, 7, 1253; (i) Q. X. Liu, L. X. Zhao, X. J. Zhao, Z. X. Zhao, Z. Q. Wang, A. H. Chen and X. G. Wang, *J. Organomet. Chem.*, 2013, 731, 35.

- 6 (a) S. P. Nolan, *N-Heterocyclic Carbenes in Synthesis*, Wiley-VCH Verlag GmbH & Co., KGaA, Weinheim, 2006. (b) N. Marion, O. Navarro, J. G. Mei, E. D. Stevens, N. M. Scott and S. P. Nolan, *J. Am. Chem. Soc.*, 2006, **128**, 4101; (c) G. Altenhoff, R. Goddard, C. W. Lehmann and F. Glorius, *J. Am. Chem. Soc.*, 2004, **126**, 15195; (d) V. César, S. Bellemin-Laponnaz and L. H. Gade, *Organometallics*, 2002, **21**, 5204; (e) B. Karimi and D. Enders, *Org. Lett.*, 2006, **8**, 1237.
- 7 F. W. Li, S. Q. Bai and T. S. A. Hor, Organometallics, 2008, 27, 672.
- 8 H. L. Qiu, S. M. Sarkar, D. H. Lee and M. J. Jin, Green Chem., 2008, 10, 37.
- 9 (a) F. Glorius (Ed.), N-heterocyclic Carbenes in Transition Metal Catalysis, in: Topics in Organometallic Chemistry, Springer, Berlin/Heidelberg, Germany, 2007, vol. 21; (b) S. K. Yen, L. L. Koh, F. E. Hahn, H. V. Huynh and T. S. A. Hor, Organometallics, 2006, 25, 5105.
- 10 (a) A. A. Danopoulos, N. Tsoureas, S. A. Macgregor and C. Smith, Organometallics, 2007, 26, 253; (b) C. C. Lee, W. C. Ke, K. T. Chan, C. L. Lai, C. H. Hu and H. M. Lee, Chem. Eur. J., 2007, 13, 582; (c) J. Wolf, A. Labande, J. C. Daran and R. Poli, J. Organomet. Chem., 2006, 691, 433; (d) J. Y. Zeng, M. H. Hsieh and H. M. Lee, J. Organomet. Chem., 2005, 690, 5662; (e) L. D. Field, B. A. Messerle, K. Q. Vuong and P. Turner, Organometallics, 2005, 24, 4241; (f) P. L. Chiu and H. M. Lee, Organometallics, 2005, 24, 1692; (g) H. M. Lee, J. Y. Zeng, C. H. Hu and M. T. Lee, Inorg. Chem., 2004, 43, 6822; (h) C. L. Yang, H. M. Lee and S. P. Nolan, Org. Lett., 2001, 3, 1511; (i) N. Tsoureas, A. A. Danopoulos, A. A. D. Tulloch and M. E. Light, Organometallics, 2003, 22, 4750.
- 11 (a) M. Y. Chiang, Y. X. Li, D. Krishnan, P. Sumod, K. H. Ng and P. H. Leung, Eur.

J. Inorg. Chem., 2010, 1413; (b) N. Stylianides, A. A. Danopoulos and N. Tsoureas, J. Organomet. Chem., 2005, 690, 5948; (c) D. S. McGuinness and K. J. Cavell, Organometallics, 2000, 19, 741.

- 12 (a) N. A. Jones, S. T. Liddle, C. Wilson and P. L. Arnold, Organometallics, 2007,
 26, 755; (b) P. L. Arnold, M. Rodden and C. Wilson, Chem. Commun., 2005, 1743;
 (c) P. L. Arnold, M. Rodden, K. M. Davis, A. C. Scarisbrick, A. J. Blake and C. Wilson, Chem. Commun., 2004, 1612; (d) H. Clavier, L. Coutable, L. Toupet, J. C. Guillemin and M. Mauduit, J. Organomet. Chem., 2005, 690, 5237; (e) A. W. Waltman and R. H. Grubbs, Organometallics, 2004, 23, 3105; (f) P. L. Arnold, A. C. Scarisbrick, A. J. Blake and C. Wilson, Chem. Commun., 2001, 2340.
- 13 (a) A. T. Normand and K. J. Cavell, *Eur. J. Inorg. Chem.*, 2008, 2781; (b) E. A. B.
 Kantchev, C. J. O'Brien and M. G. Organ, *Angew. Chem.*, 2007, **119**, 2824.

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41

- 14 (a) P. L. Arnold and I. J. Casely, *Chem. Rev.*, 2009, **109**, 3599; (b) X. L. Liu and W. Z. Chen, *Organometallics*, 2013, **31**, 6614; (c) Q. Li, X. Li, J. Yang, H. B. Song and L. F. Tang, *Polyhedron*, 2013, **59**, 29; (d) J. H. Chen, X. Q. Zhang, Q. Feng and M. M. Luo, *J. Organomet. Chem.*, 2006, **691**, 470; (e) J. W. Wang, F. H. Meng and L. F. Zhang, *Organometallics*, 2009, **28**, 2334; (f) C. Yan, X. M. Zeng, W. F. Zhang, M. Yan and M. M. Luo, *J. Organomet. Chem.*, 2006, **691**, 3391.
- 15 (a) J. J. Van Veldhuizen, J. E. Campbell, R. E. Giudici and A. H. Hoveyda, J. Am. Chem. Soc., 2005, 127, 6877; (b) J. C. Y. Lin, R. T. W. Huang, C. S. Lee, A. Bhattacharyya, W. S. Hwang and I. J. B. Lin, Chem. Rev., 2009, 109, 3561; (c) C. M. Crudden and D. P. Allen, Coord. Chem. Rev., 2004, 248, 2247; (d) Y. S. Jiang, W. Z. Chen and W. M. Lu, Tetrahedron, 2013, 69, 3669; (e) C. H. Cheng, D. F. Chen, H. B. Song and L. F. Tang, J. Organomet. Chem., 2013, 726, 1; (f) X. Wang, S. Liu, L. H. Weng and G. X. Jin, Organometallics, 2006, 25, 3565; (g) L. H. Gade and S. Bellemin-Laponnaz, Coord. Chem. Rev., 2007, 251, 718; (h) B. Liu, C. Y. Chen, Y. J. Zhang, X. L. Liu and W. Z. Chen, Organometallics, 2013, 32, 5451; (i) D. W. Zhao, Y. F. Xie, H. B. Song and L. F. Tang, J. Organomet. Chem., 2013, 718, 89; (j) J. W. Wang, Q. S. Li, F. B. Xu, H. B. Song and Z. Z. Zhang, Eur. J. Org. Chem., 2006, 1310.

- 16 (a) Q. Xu, W. L. Duan, Z. Y. Lei, Z. B. Zhu and M. Shi, *Tetrahedron*, 2005, 61, 11225; (b) C. S. Cao, Y. Z. Zhuang, J. F. Zhao, Y. Peng, X. L. Li, Z. Shi, G. S. Pang and Y. H. Shi, *Inorg. Chim. Acta*, 2010, 363, 3914; (c) C. S. Cao, L. L. Wang, Z. Y. Cai, L. Q. Zhang, J. Guo, G. S. Pang and Y. H. Shi, *Eur. J. Org. Chem.*, 2011, 1570.
- 17 (a) W. Wei, M. Y. Wu, Y. G. Huang, Q. Gao, F. L. Jiang and M. C. Hong, Z. Anorg. Allg. Chem., 2008, 634, 2623; (b) A. L. Pickering, G. Seeber and D. L. Long, CrystEngComm, 2005, 7, 504.
- 18 A. Suzuki, J. Organomet. Chem., 1999, 576, 147.
- 19 (a) K. Selvakumar, A. Zapf and M. Beller, Org. Lett., 2002, 4, 3031; (b) R. B. Bedford, M. E. Blake, C. P. Butts and D. Holder, Chem. Commun., 2003, 466; (c) D. Yang, Y. C. Chen and N. Y. Zhu, Org. Lett., 2004, 6, 1557; (d) H. V. Huynh, Y. Han, J. H. Ho and G. K. Tan, Organometallics, 2006, 25, 3267; (e) S. H. Li, Y. J. Lin, J. G. Cao and S. B. Zhang, J. Org. Chem., 2007, 72, 4067.
- 20 (a) H. A. Dieck and R. F. Heck, J. Am. Chem. Soc., 1974, 96, 1133; (b) Y. Ben-David, M. Portnoy, M. Gozin and D. Milstein, Organometallics, 1992, 11, 1995; (c) A. Jutand and A. Mosleh, Organometallics, 1995, 14, 1810; (d) X. Fu, S. Zhang, J. Yin, T. L. Mcallister, S. A. Jiang, C. H. Tann, T. K. Thiruvengadam and F. Zhang, Tetrahedron Lett., 2002, 43, 573; (e) A. L. Hansen and T. Skrydstrup, Org. Lett., 2005, 7, 5585.
- 21 (a) J. H. Li, X. C. Hu, Y. Liang and Y. X. Xie, *Tetrahedron Lett.*, 2006, 62, 31; (b)
 J. H. Li, W. J. Liu and Y. X. Xie, *J. Org. Chem.*, 2005, 70, 5409.
- 22 (a) L. G. Yang, Y. F. Li, C. Qian, Y. F. Du and C. S. Cao, *Tetrahedron*, 2013, 69, 5178; (b) F. T. Luo and H. K. Lo, J. Organomet. Chem., 2011, 696, 1262.
- 23 (a) N. Debono, A. Labande, E. Manoury, J. C. Daran and R. Poli, *Organometallics*, 2010, 29, 1879; (b) M. Charbonneau, G. Addoumieh, P. Oguadinma and A. R. Schmitzer, *Organometallics*, 2014, 33, 6544; (c) L. Benhamou, C. Besnard and E. P. Kündig, *Organometallics*, 2014, 33, 260; (d) L. Wu, A. Salvador, A. Ou, M. W. Shi, B. W. Skelton and R. Dorta, *Synlett.*, 2013, 24, 1215; (e) S. Meiries, G. L. Duc, A. Chartoire, A. Collado, K. Speck, K. S. A. Arachchige, A. M. Z. Slawin and S. P.

Nolan, *Chem. Eur. J.*, 2013, **19**, 17358; (f) G. Grach, G. Pieters, A. Dinut, V. Terrasson, R. Medimagh, A. Bridoux, V. Razafimahaleo, A. Gaucher, S. Marque, J. Marrot, D. Prim, R. Gil, J. G. Planas, C. Viñas, I. Thomas, J. P. Roblin and Y. Troin, *Organometallics*, 2011, **30**, 4074; (g) G. W. Shigeng, J. K. Tang, D. Zhang, Q. R. Wang, Z. X. Chen and L. H. Weng, *J. Organomet. Chem.*, 2012, **700**, 223; (h) Y. L. Li, J. K. Tang, J. Gu, Q. R. Wang, P. P. Sun and D. Zhang, *Organometallics*, 2014, **33**, 876; (i) Y. Q. Tang, J. M. Lu and L. X. Shao, *J. Organomet. Chem.*, 2011, **696**, 3741; (j) M. M. T. Strassner, *Eur. J. Inorg. Chem.*, 2012, 5872; (k) H. Lebel, M. K. Janes, A. B. Charette and S. P. Nolan, *J. Am. Chem. Soc.*, 2004, **126**, 5046; (l) J. D. Blakemore, M. J. Chalkley, J. H. Farnaby, L. M. Guard, N. Hazari, C. D. Incarvito, E. D. Luzik Jr. and H. W. Suh, *Organometallics*, 2011, **30**, 1818; (m) C. E. Ellul, G. Reed, M. F. Mahon, S. I. Pascu and M. K. Whittlesey, *Organometallics*, 2010, **29**, 4097.

Published on 05 October 2015. Downloaded by University of Pennsylvania Libraries on 05/10/2015 14:52:41

- 24 (a) Y. F. Li, L. G. Yang, Q. Chen, C. S. Cao, P. Guan, G. S. Pang and Y. H. Shi, Z. Anorg. Allg. Chem., 2013, 639, 575; (b) K. Q. Ge, C. S. Cao, G. Liu, Y. L. Li, J. Zhang, G. S. Pang and Y. H. Shi, Z. Anorg. Allg. Chem., 2014, 640, 444; (c) U. I. Tessin, X. Bantreil, O. Songis and C. S. J. Cazin, Eur. J. Inorg. Chem., 2013, 2007; (d) C. S. Lee, Y. B. Lai, W. J. Lin, R. R. Zhuang and W. S. Hwang, J. Organomet. Chem., 2013, 724, 235; (e) H. V. Huynh and R. Jothibasu, J. Organomet. Chem., 2011, 696, 3369; (f) T. Yagyu, A. Tsubouchi and Y. Nishimura, J. Organomet. Chem., 2014, 767, 1; (g) H. Türkmen, T. Pape, F. E. Hahn and B. Çetinkaya, Eur. J. Inorg. Chem., 2009, 285; (h) H. B. Metzner, P. T. Körzdörfer, A. Kelling and H. J. Holdt, Eur. J. Inorg. Chem., 2014, 2952; (i) A. L. Gottumukkala, J. G de Vries and A. J. Minnaard, Chem. Eur. J., 2011, 17, 3091; (j) Q. Teng, D. Upmann, S. A. Z. N. Wijaya and H. V. Huynh, Organometallics, 2014, 33, 3373; (k) A. Z. Wilczewska and I. Misztalewska, Organometallics, 2014, 33, 5203.
- 25 (a) L. G. Yang, Y. F. Li, Q. Chen, Y. F. Du, C. S. Cao, Y. H. Shi and G. S. Pang, *Tetrahedron*, 2013, **69**, 5178; (b) S. C. Holm, F. Rominger and B. F. Straub, J. *Organomet. Chem.*, 2012, **719**, 54; (c) A. John, S. Modak, M. Madasu, M. Katari and P. Ghosh, *Polyhedron*, 2013, **64**, 20; (d) L. G. Yang, Y. F. Li, Q. Chen, J. F.

RSC Advances

Zhao, C. S. Cao, Y. H. Shi and G. S. Pang, *Transition Met. Chem.*, 2013, 38, 367; (e)
C. Dash, M. M. Shaikh and P. Ghosh, *Eur. J. Inorg. Chem.*, 2009, 1608; (f) C. Chen,
H. Qiu and W. Z. Chen, *Inorg. Chem.*, 2011, 50, 8671; (g) J. C. Bernhammer, N. X.
Chong, R. Jothibasu, B. Zhou and H. V. Huynh, *Organometallics*, 2014, 33, 3607;
(h) L. Ray, S. Barman, M. M. Shaikh and P. Ghosh, *Chem. Eur. J.*, 2008, 14, 6646.

- 26 (a) Y. H. Chang, Z. Y. Liu, Y. H. Liu, S. M. Peng, J. T. Chen and S. T. Liu, *Dalton Trans.*, 2011, 40, 489; (b) T. Zhang, S. J. Liu, M. Shi and M. X. Zhao, *Synthesis*, 2008, 17, 2819; (c) M. T. Ma and J. M. Lu, *Appl. Organometal. Chem.*, 2012, 26, 175.
- 27 (a) A. Zanardi, J. A. Mata and E. Peris, Organometallics, 2009, 28, 4335; (b) A. Zanardi, J. A. Mata and E. Peris, Organometallics, 2009, 28, 1480; (c) S. K. Yen, L. L. Koh, H. V. Huynh and T. S. A. Hor, Chem. Asian J., 2008, 3, 1649.
- 28 (a) E. Mas-Marzà, J. A. Mata and E. Peris, *Angew. Chem.*, *Int. Ed.*, 2007, 46, 3729;
 (b) A. Zanardi, R. Corberan, J. A. Mata and E. Peris, *Organometallics*, 2008, 27, 3570; (c) M. Viciano, M. Sanaú and E. Peris, *Organometallics*, 2007, 26, 6050.
- 29 M. L. Cole and P. C. Junk, CrystEngComm, 2004, 6, 173.

30 Bruker AXS, SAINT Software Reference Manual, Madison, WI, 1998.

31 G. M. Sheldrick, SHELXTL NT (Version 5.1), Program for Solution and Refinement of Crystal Structures, University of Göttingen, Germany, 1997.

32 D. C. Palmer, CrystalMaker 7.1.5, CrystalMaker Software, Yarnton, UK, 2006.

A table of contents entry:

NHC-Pd(II) complex 1 has been prepared. The catalytic activity of 1 in C-C coupling reactions was studied.



Software of Graphics:

Scheme 1: Chem Draw 8.0 Fig. 1: Diamand 3.0