

Scheme 3. Transformation of **7** to the vinyl iodide **11** and the following coupling reaction.

easily available in quantity, and the reaction operation is easy.

## **Experimental Section**

Typical procedure for the site-selective C–C bond formation of **2** with organocuprates (preparation of **3aa**, Table 1, entry 1): To a stirred solution of CuI (1.99 g, 10.4 mmol) in Et<sub>2</sub>O (10 mL) was added a 1.04 m solution of MeLi in ether (20.0 mL, 20.8 mmol) at 0°C, under argon, to give a homogeneous solution. The solution was cooled to -50 °C, then **2a** (1.90 g, 3.48 mmol) was added to the solution. After stirring at -50 °C for 3 h, the reaction mixture was quenched by the addition of aqueous NH<sub>4</sub>Cl and the organic products were extracted with diethyl ether. The combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo to give a crude oil. Chromatography on silica gel (hexane) afforded compound **3aa** (1.35 g, 90%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 0.16 (s, 9H), 0.85–0.90 (m, 6H), 1.25–1.60 (m, 16H), 1.65 (s, 3H), 2.18 (t, *J* = 8.0 Hz, 2H), 2.51 (t, *J* = 7.1 Hz, 2H), 6.19 ppm (s, 1H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  = 0.41, 14.18, 14.21, 19.67, 22.73, 22.76, 27.94, 28.84, 29.45, 29.85, 31.63, 31.91, 36.61, 45.06, 110.59, 131.35, 136.40, 149.32 ppm.

Typical procedure for conversion of **4** into **5** (synthesis of **11**; Scheme 3): To a stirred solution of **7** (0.126 g, 0.345 mmol) in absolute ethanol (6 mL) was added AgNO<sub>3</sub> (0.088 g, 0.518 mmol) at 0 °C. The mixture was stirred for 10 min to ensure complete dissolution. Iodine (0.114 g, 0.449 mmol) was added in one portion and stirred at 0 °C for 20 min. The reaction mixture was diluted with diethyl ether, filtering through a celite pad. The filtrate was washed with sodium thiosulfate solution, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo to give a crude oil. Chromatography on silica gel (hexane) afforded compound **11** (0.118 g, 82%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 0.86–0.91 (m, 9H), 1.21–1.42 (m, 20H), 1.89 (t, *J* = 7.4 Hz, 2H), 2.05 (t, *J* = 7.1 Hz, 2H), 2.24 (t, *J* = 7.1 Hz, 2H), 2.42 (s, 3 H), 5.47 ppm (s, 1H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  = 14.16, 14.20, 14.23, 22.74, 22.76, 23.13, 27.94, 28.04, 29.20, 29.27, 29.61, 30.75, 31.67, 31.82, 31.86, 35.47, 43.36, 96.24, 121.60, 142.69, 143.15 ppm.

Received: May 2, 2002 [Z19197]

- [1] T. Hamada, D. Suzuki, H. Urabe, F. Sato, J. Am. Chem. Soc. 1999, 121, 7342-7344.
- [2] Recent reviews of synthetic reactions mediated by a Ti(O-*i*Pr)₄/ 2*i*PrMgX reagent: a) F. Sato, H. Urabe, S. Okamoto, *Chem. Rev.* 2000, 100, 2835–2886; b) F. Sato, S. Okamoto, *Adv. Synth. Catal.* 2001, 343, 759–784.
- [3] S. Yamaguchi, R. Jin, K. Tamao, F. Sato, J. Org. Chem. 1998, 63, 10060-10062.
- [4] R. B. Miller, G. McGarvey, J. Org. Chem. 1979, 44, 4623-4633.
- [5] a) G. H. Posner, Org. React. 1975, 22, 253-400; b) B. H. Lipshutz, S. Sengupta, Org. React. 1992, 41, 135-631; c) B. H. Lipshutz in Organometallics in Synthesis (Ed.: M. Schlosser), Wiley, Chichester, 1994, pp. 283-382.
- [6] N. Miyaura, A. Suzuki, Chem. Rev. 1995, 95, 2457-2483.

## COMMUNICATIONS

- [7] a) K. Sonogashira, Y. Tohda, N. Hagihara, *Tetrahedron Lett.* 1975, 4467–4470; b) K. Sonogashira in *Comprehensive Organic Synthesis*, *Vol. 3* (Eds.: B. M. Trost, I. Fleming, G. Pattenden), Pergamon, Oxford, 1991, pp. 521–549.
- [8] T. H. Chan, I. Fleming, Synthesis 1979, 761-786.
- [9] a) S. Rousset, J. Thibonnet, M. Abarbri, A. Duchêne, J.-C. Parrain, *Synlett* **2000**, 260–262; b) S. R. Wilson, L. A. Jacob, *J. Org. Chem.* **1986**, *51*, 4833–4836.
- [10] The stereochemistry of **11** was determined by NOESY NMR spectroscopy.

## Reaction of "[Mn<sup>II</sup>(CH<sub>2</sub>*t*Bu)<sub>2</sub>]" with Bidentate Diimine Ligands: From Simple Base Adducts to C-C Activation of the Ligand\*\*

Virginie Riollet, Christophe Copéret,\* Jean-Marie Basset, Laurence Rousset, Denis Bouchu, Laurent Grosvalet, and Monique Perrin

Base adducts of dialkyl  $Mn^{II}$  complexes are typically monomeric tetrahedral high-spin systems;<sup>[1]</sup> our synthetic target is to extend these adducts to bidentate diimine ligands. Gambarotta and co-workers<sup>[2]</sup> recently reported an approach to  $Mn^{II}$  analogues of Brookhart–Gibson complexes<sup>[3]</sup> and compared their performance in olefin polymerization with those of Fe<sup>II</sup> and Ni<sup>II</sup> catalysts. In this approach, they showed that the reaction of [MnCl<sub>2</sub>L<sub>2</sub>] complexes with organolithium reagents gave the corresponding Mn<sup>I</sup> and Mn<sup>0</sup> alkyl complexes, that is, the lithium reagents alkylate and reduce the Mn<sup>II</sup> dichloro complex. These results have led us to disclose our investigation on the reaction of a bisneopentyl Mn<sup>II</sup> complex with bidentate Shiff base ligands that were designed to generate complexes with structure **1** (Scheme 1). The major difference in our approach is the reaction of diimine ligands

- [\*\*] We are indebted to the CNRS and ESCPE Lyon for financial support. V.R. is grateful to the Région Rhône-Alpes for a predoctoral fellowship. We also thank Prof. R. A. Andersen (U. C. Berkeley), Prof. Dr. Koelher (TUM), Dr. R. Gauvin (LCOMS), and Dr. C. Boisson (LCPP) for helpful discussions.
- Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.

0044-8249/02/4116-3025 \$ 20.00+.50/0

Angew. Chem. Int. Ed. 2002, 41, No. 16 © 2002 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

3025

<sup>[\*]</sup> Dr. C. Copéret, V. Riollet, Dr. J.-M. Basset Laboratoire de Chimie Organométallique de Surface UMR-9986 CNRS-ESCPE Lyon
43 bd du 11 Novembre 1918, 69616 Villeurbanne Cedex (France) Fax: (+ 33)472-431-795
E-mail: coperet@cpe.fr
L. Rousset, Dr. D. Bouchu Centre de Spectrométrie de Masse Université Claude Bernard Lyon I
43 bd du 11 Novembre 1918, 69622 Villeurbanne Cedex (France)
L. Grosvalet, Prof. Dr. M. Perrin Laboratoire de Cristallographie UMR-5078 Université Claude Bernard Lyon I
43 bd du 11 Novembre 1918, 69622 Villeurbanne Cedex (France)
Kaboratoire de Cristallographie UMR-5078
Université Claude Bernard Lyon I
43 bd du 11 Novembre 1918, 69622 Villeurbanne Cedex (France)

## COMMUNICATIONS

with manganese(II) dialkyl complexes, which proceeds without reduction. Depending on the structure of the ligand it is possible to isolate either the desired base adduct or compounds in which the imine functional group is alkylated.

A brown solution of  $[{Mn(CH_2tBu)_2}_4]$ (1 equiv) in toluene was added to a colorless solution of trans-N,N'-bis(benzylidene)-1,2-cyclohexanediamine in toluene, and the solution turned dark-purple. After 12 h, removal of toluene and addition of pentane led to precipitation of unconsumed ligand (excess ligand: vide infra). After complete removal of the ligand and concentration to dryness, a concentrated solution of the solid in toluene was cooled to -30 °C to give single crystals as brown rectangular parallelepipeds (20%). However, X-ray analysis revealed the presence of 2 (Figure 1) with a structure that was dramatically different from that of the expected 1a.<sup>[4]</sup> Complex 2 is dinuclear, and the Mn-Mn distance of 2.822 Å implies that there is no metal-metal bond. Mn(1) is bound to one neopentyl ligand as well as to an imino and an amido ligand, whereas

Mn(2) is bonded to the electron lone pair of the nitrogen atom of this amido group, but retains the two neopentyl ligands. Complex **2** is probably obtained via the expected **1a** (Scheme 1), which then undergoes migration of the neopentyl ligand onto the imine group<sup>[5]</sup> to generate a mononeopentyl Mn<sup>II</sup> complex with one imine and one amido ligand, which further reacts with the putative "[Mn(CH<sub>2</sub>*t*Bu)<sub>2</sub>]" to give **2**. Thus, two equivalents of "[Mn(CH<sub>2</sub>*t*Bu)<sub>2</sub>]" are required for each equivalent of ligand. This suggested reaction pathway is, however, difficult to substantiate, since starting material and product(s) contain high-spin Mn<sup>II</sup>. Moreover, **2** was isolated as a single stereoisomer. The stereochemistry at the carbon center corresponds to the attack of the neopentyl fragment



Figure 1. Thermal-ellipsoid plot of **2**. Selected bond lengths [Å] and angles [°]: Mn(1)-C(24) 2.121(5), Mn(1)-C(26) 2.419(5), Mn(1)-N(1) 2.132(4), Mn(1)-N(2) 2.242(4), Mn(2)-C(26) 2.223(5), Mn(2)-C(231) 2.122(5), Mn(2)-N(1) 2.129(4); N(1)-Mn(1)-N(2) 79.35(2), N(1)-Mn(1)-C(24) 142.63(2), N(2)-Mn(1)-C(24) 118.48(2), N(1)-Mn(2)-C(26) 104.05(2), N(1)-Mn(2)-C(231) 129.99(2), C(26)-Mn(2)-C(231) 123.8(2).

3026 © 2002 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim



Scheme 1. Generation of **2** via **1**, through migration of a neopentyl group and further reaction with the putative " $[Mn(CH_2/Bu)_2]$ ".

from the *Si* face of the imine moiety, that is, the face with the methine hydrogen atom rather than the C–C framework of the cyclohexane ring.

Reaction of a more highly substituted bisbenzylidene ligand (Ar = mesityl (Mes)) with " $[Mn(CH_2tBu)_2]$ " gave the desired base adduct **1b** in 36% yield of isolated product. X-ray analysis shows that the geometry of the complex is a distorted tetrahedron with a wide C-Mn-C bond angle of 146.4° (Figure 2).<sup>[4]</sup> The methyl substituents of the aromatic ring probably shield the imino group (see Scheme 1 and Figure 2) and thus prevent the migration of the neopentyl ligand to the imino group.

To understand the scope of this chemistry further, we also investigated the reaction of the Brookhart imino ligand with



Figure 2. Thermal-ellipsoid plot of **1b**. Selected bond lengths [Å] and angles [°]: Mn–C(40) 2.141(6), Mn–C(50) 2.165(6), Mn–N(1) 2.344(4), Mn–N(2) 2.279(4); C(40)-Mn-C(50) 146.4(2), C(40)-Mn-N(1) 102.1(2), C(40)-Mn-N(2) 110.3(2), C(50)-Mn-N(1) 105.62(2), C(50)-Mn-N(2) 95.71(2), N(1)-Mn-N(2) 73.12(2).

0044-8249/02/4116-3026 \$ 20.00+.50/0

Angew. Chem. Int. Ed. 2002, 41, No. 16

" $[Mn(CH_2tBu)_2]$ ". In this case the reaction yield is higher (51% of single crystals), but structural analysis again showed major reorganization of the putative base adduct by carbon–carbon migration within the ligand (Figure 3).<sup>[4]</sup> The reaction probably involves adduct formation followed by the events outlined in Scheme 2. Migration of one neopentyl ligand in **4** 



Figure 3. Thermal ellipsoid plot of 6. Selected bond lengths [Å] and angles [°]: Mn-C(13) 2.105(3), Mn-N(1) 1.960(2), Mn-N(2) 2.185(3); C(13)-Mn-N(1) 150.86(12), C(13)-Mn-N(2) 129.68(11), N(1)-Mn-N(2) 79.11(10).



Scheme 2. Proposed mechanism for the migration reaction.

onto the imino group to give **5** is apparently followed by a 1,2migration to the vicinal imino group. This rearrangement could also be a two-step process involving successive reversible 1,3-migration of the Me group from the C atom to the Mn atom and then to the other C atom to give an apparent overall 1,2-migration. Complex **6** has an Mn<sup>II</sup> center in a trigonal environment with an imino, an amido, and a neopentyl ligand, like in **2**, but does not undergo further reaction with "[Mn(CH<sub>2</sub>tBu)<sub>2</sub>]", probably because the 2,6diisopropylphenyl substituent prevents access to the lone pair of the amido ligand.

In conclusion, we have shown that diimine Mn<sup>II</sup> dialkyl complexes are accessible without reduction of the Mn center, which demonstrates the importance to access perhydrocarbyl metal complexes with the appropriate oxidation state. We have also demonstrated the necessity to protect the imino group with bulky substituents to avoid their reaction with

metal–carbon bonds. Most interestingly, we have uncovered pathways in which the C=N bonds undergo alkylation to amido groups. These reactions might be important deactivation pathways in some olefin polymerization reactions. <sup>[6]</sup>

Received: April 12, 2002 [Z19085]

- a) R. A. Andersen, E. Carmona-Guzman, J. F. Gibson, G. Wilkinson, J. Chem. Soc. Dalton Trans. 1976, 2204; b) C. G. Howard, G. S. Girolami, G. Wilkinson, M. Thornton-Pett, M. B. Hursthouse, J. Chem. Soc. Dalton Trans. 1983, 2631; c) C. G. Howard, G. Wilkinson, M. Thornton-Pett, M. B. Hursthouse, J. Chem. Soc. Dalton Trans. 1983, 2025; d) R. A. Andersen, A. Haaland, K. Rypdal, H. V. Volden, J. Chem. Soc. Chem. Commun. 1985, 1807; e) G. S. Girolami, C. G. Howard, G. Wilkinson, H. M. Dawes, M. Thornton-Pett, M. Motevalli, M. B. Hursthouse, J. Chem. Soc. Dalton Trans. 1985, 921; f) J. Morris, G. S. Girolami, Organometallics 1989, 8, 1478; g) P. B. Hitchcock, M. F. Lappert, W.-P. Leung, J. Organometal. Chem. 1990, 394, 57.
- [2] D. Reardon, G. Aharonian, S. Gambarotta, G. P. A. Yap, Organometallics 2002, 21, 786.
- [3] a) L. K. Johnson, C. M. Killian, M. Brookhart, J. Am. Chem. Soc. 1995, 117, 6414; b) L. K. Johnson, S. Mecking, M. Brookhart, J. Am. Chem. Soc. 1996, 118, 267; c) C. M. Killian, D. J. Tempel, L. K. Johnson, M. Brookhart, J. Am. Chem. Soc. 1996, 118, 11664; d) G. J. P. Britovsek, V. C. Gibson, B. S. Kimberley, P. J. Maddox, S. J. Mctavish, G. A. Solan, A. J. P. White, D. J. Williams, Chem. Commun. 1998, 849; e) B. L. Small, M. Brookhart, A. M. A. Bennett, J. Am. Chem. Soc. 1998, 120, 4049; f) G. J. P. Britovsek, M. Bruce, V. C. Gibson, B. S. Kimberley, P. J. Maddox, S. Mastroianni, S. J. McTavish, C. Redshaw, G. A. Solan, S. Strömberg, A. J. P. White, D. J. Williams, J. Am. Chem. Soc. 1999, 121, 8728; g) G. J. P. Brotovsek, S. Mastroianni, G. A. Solan, S. P. D. Baugh,
  - Sek, S. Mastrolanni, G. A. Solan, S. P. D. Baugn, C. Redshaw, V. C. Gibson, A. J. P. White, D. J. Willimas, M. R. J. Elsegood, *Chem. Eur. J.* **2000**, *6*, 2221; h) E. L. Dias, M. Brookhart, P. S. White, *Chem. Commun.* **2001**, 423; i) V. C. Gibson, M. J. Humphries, K. P. Tellmann, D. F. Wass, A. J. P. White, D. J. Williams, *Chem. Commun.* **2001**, 2252; j) T. M. Kooistra, Q. Knijnenburg, J. M. M. Smits, A. D. Horton, P. H. M. Budzelaar, A. W. Gal, *Angew. Chem.* **2001**, *113*, 4855; *Angew. Chem. Int. Ed.* **2001**, *40*, 4719.
  - [4] CCDC-183 390, CCDC-183 147, and CCDC-182 472 contain the supplementary crystallographic data for this paper (2, 1b, and 6, respectively). These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; fax: (+44) 1223-336-033; or deposit@ccdc.cam.ac.uk).
- [5] a) J. M. Klerks, D. J. Stufkens, G. van Koten, K. Vrieze, J. Organometal. Chem. 1979, 181, 271; b) C. Floriani, S. Ciurli, A. Chiesi-Villa, C. Guastini, Angew. Chem. 1987, 99, 86; Angew. Chem. Int. Ed. Engl. 1987, 26, 70; c) L. Giannini, E. Solari, S. D. Angelis, T. R. Ward, C. Floriani, A. Chiesi-Villa, C. Rizzoli, J. Am. Chem. Soc. 1995, 117, 5801; d) M. Bruce, V. C. Gibson, C. Redshaw, G. A. Solan, A. J. P. White, D. J. Williams, J. Organomet. Chem. 1998, 550, 453; f) A. Martin, R. Uhrhammer, T. G. Gardner, R. F. Jordan, Organometalis, 1998, 17, 382; g) D. Reardon, F. Conan, S. Gambarotta, G. Yap, Q. Wang, J. Am. Chem. Soc. 1999, 121, 9318; h) P. D. Knight, A. J. Clarke, B. S. Kimberley, R. A. Jackson, P. Scott, Chem. Commun. 2002, 352; i) R. Gauvin, PhD thesis, University of Strasbourg, 1999
- [6] No ethylene polymerization activities were found in either case using MAO or  $B(C_6F_5)_3$  as co-catalysts.