

# Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: L. Candish, M. Freitag, T. Gensch and F. Glorius, *Chem. Sci.*, 2017, DOI: 10.1039/C6SC05533H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

## COMMUNICATION

## Mild, visible light-mediated decarboxylation of aryl carboxylic acids to access aryl radicals


Received 00th January 20xx,  
Accepted 00th January 20xxL. Candish,<sup>a</sup> M. Freitag,<sup>a</sup> T. Gensch<sup>a</sup> and F. Glorius<sup>a</sup>

DOI: 10.1039/x0xx00000x


www.rsc.org/

Herein we present the first example of aryl radical formation via the visible light-mediated decarboxylation of aryl carboxylic acids using photoredox catalysis. This method constitutes a mild protocol for the decarboxylation of cheap and abundant aryl carboxylic acids and tolerates both electron-rich substrates and those lacking *ortho*-substitution. The *in situ* formation of an acyl hypobromite is proposed to prevent unproductive hydrogen atom abstraction and trapping of the intermediate aryloxy radical, enabling mild decarboxylation.


Since seminal publications by Lei,<sup>1a</sup> Nishibayashi<sup>1b</sup> and MacMillan,<sup>1c</sup> the decarboxylation of alkyl carboxylic acids via photoredox catalysis<sup>2</sup> has become a commanding method to generate alkyl radicals<sup>1d</sup> which can undergo either direct functionalisation or participate in dual-catalytic processes.<sup>1e</sup> Despite strong interest in this area, the decarboxylation of aryl carboxylic acids using photoredox catalysis is yet to be realised. While aryl radicals can be accessed via photoredox catalysis from electron-poor arenes, such as diazonium and iodonium salts, aryl halides,<sup>3</sup> and arylborates,<sup>4</sup> many of these reagents are unstable, expensive and often not commercially available. In contrast, aryl carboxylic acids are abundant and cheap, making them appealing starting materials. Considering the wealth of methods for the functionalisation of alkyl carboxylic acids using photoredox catalysis,<sup>1</sup> the development of a mild protocol for the decarboxylative functionalisation of aryl carboxylic acids is of high importance. The decarboxylation of aryloxy radicals (**I**) is known to be challenging,<sup>5a</sup> in fact, Barton categorised **I** as a “non-decarboxylating acyloxy radical”, noting that **I** does not decarboxylate at temperatures below 120–130 °C.<sup>5b–c</sup> Though generation of **I** via photoredox catalysis has been described, either via single electron reduction of benzoyl peroxide<sup>6a</sup> or

a. Photoredox mediated generation of aryl acyloxy radical **I**

## b. Preparation of benzoyl hypohalites



## c. This strategy



Scheme 1. Challenges of photoredox-catalysed decarboxylation of aryl carboxylic acids and this approach to achieve a mild decarboxylation protocol.

single electron oxidation of benzoates,<sup>6b,7</sup> decarboxylation was not reported in these cases. Instead, **I** has been reported to add to arenes to provide benzoate esters (Scheme 1a).<sup>6</sup> Additionally, our laboratory recently disclosed the use of **I** as a hydrogen atom transfer (HAT) catalyst for the functionalisation of unactivated C(sp<sup>3</sup>)-H bonds.<sup>7</sup> While the activation energy for the decarboxylation of **I** is only 8–9 kcal/mol,<sup>8a</sup> the rate of decarboxylation ( $k = 1.4 \times 10^6 \text{ s}^{-1}$ ) is not competitive with nondecarboxylative pathways including addition to arenes ( $k = 2.2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ )<sup>8a</sup> and HAT ( $k = 1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>8a,9</sup> As such, while the decarboxylation of **I** is theoretically possible its realisation is practically challenging. By comparison, the more facile decarboxylation of alkyl carboxylic acids, which readily occurs at ambient temperatures, proceeds at least one thousand times faster than the decarboxylation of **I**.<sup>8c</sup> Recently, Greaney<sup>10a,b</sup> and Su<sup>10c</sup> disclosed their work describing the catalytic oxidative radical decarboxylation of aryl

<sup>a</sup> Organisch-Chemisches Institut  
Westfälische Wilhelms-Universität Münster.  
Corrensstrasse 40, 48149, Münster, Germany.  
glorius@uni-muenster.de


Electronic Supplementary Information (ESI) available: See  
DOI: 10.1039/x0xx00000x



## COMMUNICATION

carboxylic acids using Ag(I) and a stoichiometric oxidant. This work is complementary to the aryl decarboxylation protocols developed by Gooßen and Larossa involving CO<sub>2</sub> extrusion/arene metalation,<sup>11</sup> however, characteristic of all methods for aryl decarboxylation is the requirement of forcing reaction conditions. Given photoredox catalysis has become a commanding method to generate radicals under mild reaction conditions<sup>2</sup> we believed we could harness this potential in order to develop a mild process for the decarboxylation of aryl carboxylic acids. To this end, we were inspired by reports from Hammond<sup>12a</sup> and da Silva Corrêa<sup>12b</sup> that aryloxy radicals, formed via the thermal decomposition of benzoyl peroxide, react rapidly with iodination reagents to yield benzoyl hypoiodites, such as **II** (X = I) (Scheme 1b). Acyl hypoalites are also proposed intermediates in the Hunsdiecker reaction, formed via the halogenation of stoichiometric metal carboxylates (M = Ag, Hg, Tl), with heating of the benzoyl hypoalite initiating a radical chain that furnishes the halogenated product.<sup>13,14</sup> We proposed that generation of an aryloxy radical via photoredox catalysis in the presence of a suitable bromination reagent may enable the *in situ* generation of benzoyl hypobromite **II** (X = Br), thus inhibiting undesired reactions of **I** including HAT and addition to arenes (Scheme 2c). It was hoped that a subsequent single electron reduction of **II** would facilitate decarboxylation, thus yielding an aryl radical, which could be trapped with arenes to furnish biaryl products.

As the success of this process hinges on the ability of **II** to participate in single electron reduction, and for the resultant radical anion to undergo decarboxylation at temperatures unprecedented for aryl carboxylic acids, hypobromite **1** was initially synthesised and subjected to photoredox conditions. Iridium-centred photocatalyst [Ir(dF(CF<sub>3</sub>)ppy)<sub>2</sub>(dtbbpy)]PF<sub>6</sub> (dF(CF<sub>3</sub>)ppy = 2-(2,4-difluorophenyl)-5-(trifluoromethyl)pyridine, dtbbpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine) (**PC**) was selected, as luminescence quenching studies suggested it capable of oxidising benzoates, while also being significantly reducing ( $E_{1/2}^{IV/III*} = -0.89$  V or  $E_{1/2}^{III/II} = -1.37$  V vs SCE). Hypobromite **1** was prepared at low temperature and then heated to 55 °C in chlorobenzene in the absence of **PC**.<sup>15</sup> As predicted, both under irradiation with blue LEDs ( $\lambda_{max} = 455$  nm) and in the dark, only benzoate ester **2a** was formed (Eq. 1). Gratifyingly, when hypobromite **1** was irradiated at 55 °C in the presence of **PC** the desired biaryl product **3a** was afforded, along with ester **2a** (Eq. 2). This result suggests that oxidative quenching of **PC** ( $E_{1/2}^{IV/III*} = -0.89$  V vs SCE) by **1** is possible, thus providing radical anion **III** and facilitating decarboxylation. Addition of the aryl radical to the arene solvent and oxidation of the intermediary cyclohexadienyl radical by the Ir(IV) should, after deprotonation, afford biaryl **3a** with regeneration of the ground-state **PC**.



With evidence that decarboxylation of a hypobromite could be facilitated by the photocatalyst and light, we commenced studies into the photoredox catalysed decarboxylation of aryl carboxylic acid **4a**. The irradiation of **4a** under visible light at 55 °C with **PC** (3 mol%) and Cs<sub>2</sub>CO<sub>3</sub> (2 equiv) in the absence of bromine resulted in only 1% yield of biaryl **3b** (Chart 1). However, the addition of bromine (3.5 equiv) to the reaction enabled the product **3b** to form in 51% yield (Chart 1), along with approximately 10% of corresponding benzoate ester **2b**. When this reaction was performed in the absence of **PC**, a mixture of mono-, di-, tri- and tetra-bromobenzene was observed,<sup>16</sup> along with 14% **2b**. As bromine is a toxic, volatile liquid, incompatible with numerous functional groups, we decided to explore alternative brominating agents to determine whether they could also affect the reaction. Pleasingly, we found many brominating reagents capable of facilitating product formation (see SI for a list of the halogenation reagents screened). In line with our previous studies, bromomalonate derivatives were found to be excellent reagents,<sup>1h,17</sup> with **5** providing the product in 77% yield (with no benzoate ester observed), while *N*-bromosuccinimide (NBS) and *N*-bromophthalimide (NBP) provided the product in lower yields (Chart 1).

To gain a deeper understanding of the decarboxylation, the influence of the sterics and electronics of the aryl carboxylate and the electronics of the trapping arene were studied (Scheme 2). The decarboxylation of electron-deficient acids with *ortho*-substituents proceeded smoothly at 55 °C to provide biaryls (**3b-f**). The reaction of *ortho*-bromobenzoic acid (**4b**) was performed on 1.5 mmol scale, though an extended reaction time (60 h) was required.





Chart 1. Screen of different bromination reagents. <sup>a</sup>Reactions were performed using **4a** (0.1 mmol), **PC** (3 mol%), Cs<sub>2</sub>CO<sub>3</sub> (2 equiv), [**Br**] (3.5 equiv), arene (2.2 mL), blue LED ( $\lambda_{max} = 455$  nm), 55 °C, 22 h. Yields determined by GC-MS using decane as standard.





Scheme 2. Scope of the biaryl synthesis. <sup>a</sup>Reactions were performed using benzoic acid (0.2 mmol), **PC** (3 mol%),  $\text{Cs}_2\text{CO}_3$  (2 equiv), **5** (3.5 equiv), arene (4.5 mL), blue LED ( $\lambda_{\text{max}} = 455 \text{ nm}$ ), 55 °C, 22 h. Isolated yields (average of two reactions) following column chromatography. <sup>b</sup>Reaction were performed with arene (150 equiv) in  $\text{CH}_3\text{CN}$  (1:1 vol/vol). <sup>c</sup>Reaction were performed on 1.5 mmol scale, reaction time = 60 h. <sup>d</sup>Reaction were performed with blue LED ( $\lambda_{\text{max}} = 415 \text{ nm}$ ) at 80 °C. <sup>e</sup>2 equiv **5** used. <sup>f</sup>Reaction were performed using 23 W CFL. <sup>g</sup>Ratio of isomers determined by GC-MS. <sup>h</sup>Ratio of isomers determined by  $^{19}\text{F}$  NMR spectroscopic analysis of the crude mixture.

The reaction of electron-deficient acids containing *meta*- or *para*-substituents required heating the reaction to 80 °C to achieve acceptable yields of the products (**3g-h**). Acids containing electron-donating substituents are generally not tolerated in the catalytic oxidative radical decarboxylation reactions described by Greaney and Su,<sup>10b,c</sup> presumably because of their slower rate of decarboxylation.<sup>18</sup> Interestingly, electron-rich acids were found to react smoothly under our reaction conditions to afford products **3j-n**. A small amount of decarboxylative bromination was observed for electron-rich benzoic acids, although this could be suppressed by reducing the equivalents of **5**, though lower yields of the

biaryl products were observed. Pleasingly, the decarboxylation of *para-tert*-butylbenzoic acid proceeded smoothly, proving the product (**3k**) in similar yield (54% c.f. 57%) using a 23W CFL (compact fluorescent lamp) instead of blue LEDs. Substituted nicotinic and isonicotinic acids also underwent decarboxylative arylation in reasonable yields to provide phenylpyridines (**3r-u**).

The scope of the trapping arene was also explored, with 1,4-disubstituted arenes affording single biaryl products (**3v-w**), while electron-rich and electron-deficient monosubstituted arenes act as efficient radical traps to provide the coupled products as a mixture of isomers (**3a, 3x-y**). Additionally, electron-deficient heteroarenes also provided satisfactory yields of phenylpyridine products (**3z, 3aa-ab**).<sup>19</sup> In all cases, the arene was used in excess. However, filtration of the crude reaction mixture through a short silica gel plug followed by distillation generally allowed greater than 80% of the remaining arene to be reisolated in high purity.


In order to gain insight into the reaction mechanism we studied the kinetics of the excited **PC** via Stern-Volmer luminescence quenching. On the basis of these results we propose that reductive quenching of **PC** ( $\text{Ir(III)}^*/\text{Ir(II)}$ ) by the benzoate anion provides aryloxy radical **I** (Scheme 3). Two alternative pathways were then considered for decarboxylation of **I** to provide the aryl radical. Pathway A involves the direct decarboxylation of **I**, a process that is known to be energetically feasible but is unprecedented at these reaction temperatures.<sup>5a,b,8a,20</sup> Alternatively, pathway B is proposed to involve bromination of **I** to furnish hypobromite **II**, thus circumventing unproductive reactions of **I** with the solvent. Based on the results of the stoichiometric experiments of **1** (*vide supra*), we propose a single electron reduction of **II** by the sufficiently reducing  $\text{Ir(II)}$  ( $E_{1/2}^{\text{III/II}} = -1.37 \text{ V vs SCE}$ ) provides radical anion **III**, which undergoes decarboxylation to afford the aryl radical. Theoretical analysis suggests that upon reduction to the radical anion **III**, stepwise bond dissociation does not take place. Instead, concerted C–C and O–Br bond cleavage occurs to yield  $\text{CO}_2$ , bromide anion and the phenyl radical (see the SI for details of computational analysis). Subsequent trapping of the aryl radical (formed via either pathway) with a (hetero)arene should generate the cyclohexadienyl radical **IV**, which, following oxidation to the aryl cation and deprotonation, affords the biaryl product **3**.

Our mechanistic hypothesis was supported by a number of experiments. The reaction was undertaken using biphenyl-2-carboxylic acid (**4b**), thus enabling trapping of the aryloxy radical **I** via 6-*endo-trig* cyclisation to form lactone **6** in 89% yield (Scheme 4a).<sup>6b</sup> Additionally, to confirm the reaction was proceeding via the intermediacy of an aryl radical, the selectivity of the reaction of 4-methoxybenzoic acid with fluorobenzene was analysed and compared to the reported selectivity for the addition of 4-methoxyphenyl radical to fluorobenzene (Scheme 4b).<sup>21</sup> The observation of identical regioselectivity for the biaryl product **3ac** afforded to that reported suggests the reaction is proceeding via the same reactive species.




## COMMUNICATION

Journal Name



Scheme 3. Proposed reaction mechanism. Counterions and ligand sphere of Ir are omitted for clarity.

We next attempted to determine whether the reaction proceeds via direct decarboxylation of **I** (pathway A), or if pathway B is more likely. Though the decarboxylation of aryloxy radicals are unprecedented at 55 °C, we were curious whether the bromination reagents were merely acting as oxidants for **PC**. Thus, the reaction of **4a** was performed under the standard reaction conditions, in the absence of **5**, which was replaced with a number of reagents known to be capable of oxidising **PC**. Under these conditions, biaryl **3b** was either not formed or only observed in trace amounts (<5%) (See SI for details). Additionally, to test Barton's theory, that aryloxy radicals require >120–130 °C to undergo decarboxylation (*vide supra*), the reaction of **4a** was undertaken at 130 °C in the presence of ammonium persulfate. Biaryl **3b** was afforded in 27%, cf. 1% when the reaction was performed at 55 °C (Scheme 4c). Finally, when the reaction of biphenyl-2-carboxylic acid (**4b**) was undertaken with ammonium persulfate, instead of **5**, lactone **6** was isolated in 34% yield.<sup>6b</sup> This confirms that ammonium persulfate is a competent oxidant under the reaction conditions at 55 °C. Taken together, we believe that this provides strong experimental evidence, which is in-line with literature precedence, that the reaction is not proceeding via the direct decarboxylation of aryloxy radical **I** (pathway A). Based on the results of these mechanistic studies and the stoichiometric experiments of hypobromite **1** (Eq. 1 and 2) we tentatively propose that pathway B is operative under our reaction conditions. Experimental studies and detailed computational analysis are currently on going to gain deeper insight into the reaction mechanism.



Scheme 4. a) Trapping of the intermediate aryloxy radical; b) Comparison of selectivities for biaryl formation, ratio determined by <sup>19</sup>F NMR spectroscopic analysis; c) Visible light-mediated decarboxylation of **4a** at elevated temperatures.

## Conclusions

In summary, the first decarboxylation of aryl carboxylic acids to provide aryl radicals via photoredox catalysis is disclosed. Moreover, we have developed a method for the relatively mild decarboxylation of aryl carboxylic acids, proceeding at much lower temperatures than methods previously reported. The reaction does not require *ortho*-substituents on the aryl carboxylic acid, and, unlike previously reported catalytic oxidative decarboxylative protocols, tolerates electron-rich acids. Key to this achievement is the *in situ* formation of benzoyl hypobromite is proposed. We strongly believe that this work paves the way for the development of mild protocols for the decarboxylative functionalisation of cheap, abundant aryl carboxylic acids.

## Acknowledgements

We thank M. Teders, Dr. A. Gómez-Suárez and Dr. E. A. Standley (WWU Münster) for experimental assistance and helpful discussions. Generous financial support by the Alexander von Humboldt Foundation (L.C.) and the Deutsche Forschungsgemeinschaft (Leibniz Award) are gratefully acknowledged.

## Notes and references

- (a) J. Liu, Q. Liu, H. Yi, C. Qin, R. Bai, X. Qi, Y. Lan and A. Lei, *Angew. Chem. Int. Ed.* **2014**, *53*, 502; (b) Y. Miyake, K. Nakajima and Y. Nishibayashi, *Chem. Commun.* **2013**, *49*, 7854; (c) Z. Zuo and D. W. C. MacMillan, *J. Am. Chem. Soc.* **2014**, *136*, 5257. For recent reviews on the visible light-promoted decarboxylation of alkyl carboxylic acids see: (d) J. Xuan, Z.-G. Zhang and W.-J. Xiao, *Angew. Chem. Int. Ed.*





- 2015, 54, 15632; (e) H. Huang, K. Jia and Y. Chen, *ACS Catalysis* **2016**, 6, 4983 and references therein. For recent publications see: (f) J. D. Griffin, M. A. Zeller and D. A. Nicewicz, *J. Am. Chem. Soc.* **2015**, 137, 11340; (g) L. Candish, L. Pitzer, A. Gómez-Suárez and F. Glorius, *Chem. Eur. J.* **2016**, 22, 4753; (h) L. Candish, E. A. Standley, A. Gómez-Suárez, S. Mukherjee and F. Glorius, *Chem. Eur. J.* **2016**, 22, 9971; (i) A. Millet, Q. Lefebvre and M. Rueping, *Chem. Eur. J.* **2016**, 22, 13464.
- 2 For selected reviews on visible light photoredox catalysis, see: (a) K. Zeitler, *Angew. Chem. Int. Ed.* **2009**, 48, 9785; (b) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.* **2011**, 40, 102; (c) J. W. Tucker and C. R. J. Stephenson, *J. Org. Chem.* **2012**, 77, 1617; (d) J. Xuan and W.-J. Xiao, *Angew. Chem. Int. Ed.* **2012**, 51, 6828; (e) L. Shi and W. Xia, *Chem. Soc. Rev.* **2012**, 41, 7687; (f) S. Fukuzumi and K. Ohkubo, *Chem. Sci.* **2013**, 4, 561; (g) D. P. Hari and B. König, *Angew. Chem. Int. Ed.* **2013**, 52, 4734; (h) Y. Xi, H. Yi and A. Lei, *Org. Biomol. Chem.* **2013**, 11, 2387; (i) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.* **2013**, 113, 5322; (j) J. Xuan, L.-Q. Lu, J.-R. Chen and W.-J. Xiao, *Eur. J. Org. Chem.* **2013**, 6755; (k) M. Reckenthäler and A. G. Griesbeck, *Adv. Synth. Catal.* **2013**, 355, 2727; (l) M. N. Hopkinson, B. Sahoo, J.-L. Li and F. Glorius, *Chem. Eur. J.* **2014**, 20, 3874; (m) T. Koike and M. Akita, *Top. Catal.* **2014**, 57, 967; (n) D. A. Nicewicz and T. M. Nguyen, *ACS Catal.* **2014**, 4, 355; (o) E. Meggers, *Chem. Commun.* **2015**, 51, 3290; (p) N. A. Romero and D. A. Nicewicz, *Chem. Rev.* **2016**, 116, 10075; (q) K. L. Skubi, T. R. Blum and T. P. Yoon, *Chem. Rev.* **2016**, 116, 10035.
- 3 I. Ghosh, L. Marzo, A. Das, R. Shaikh and B. König, *Acc. Chem. Res.* **2016**, 49, 1566.
- 4 Y. Yasu, T. Koike and M. Akita, *Adv. Synth. Catal.* **2012**, 354, 3414.
- 5 (a) I. B. Seiple, S. Su, R. A. Rodriguez, R. Gianatassio, Y. Fujiwara, A. L. Sobel and P. S. Baran, *J. Am. Chem. Soc.* **2010**, 132, 13194; (b) D. H. R. Barton and M. Ramesh, *Tetrahedron Lett.* **1990**, 31, 949; (c) D. H. R. Barton, B. Lacher and S. Z. Zard, *Tetrahedron*, **1987**, 43, 4321.
- 6 (a) H. Rao, P. Wang and C.-J. Li, *Eur. J. Org. Chem.* **2012**, 6503; (b) N. P. Ramirez, I. Bosque and J. C. Gonzalez-Gomez, *Org. Lett.* **2015**, 17, 4550.
- 7 S. Mukherjee, B. Maji, A. Tlahuext-Aca and F. Glorius, *J. Am. Chem. Soc.* DOI: 10.1021/jacs.6b09970.
- 8 Rate constants for the decarboxylation ( $k$  in  $s^{-1}$ ): (a) aryloxy radical  $4\text{-ClC}_6\text{H}_4\text{C(O)O}^\bullet$ :  $1.4 \pm 0.3 \times 10^6$  ( $\text{CCl}_4$ ): J. Chateaufeuf, J. Luszytyk and K. U. Ingold, *J. Am. Chem. Soc.* **1988**, 110, 2886; (b) J. Chateaufeuf, J. Luszytyk and K. U. Ingold, *J. Am. Chem. Soc.* **1988**, 110, 2877; (c) alkyloxy radical  $= 2.2 \times 10^9$  ( $\text{CCl}_4$ ): B. Giese, In Houben-Weyl, *Methoden der Organischen Chemie, Band E 19a, C-Radikale*; M. Regitz and B. Giese, Eds.; Georg Thieme Verlag: Stuttgart, New York, 1989; p 140.
- 9 For selected example of HAT with I, see: (a) A. P. Krasutsky, C. J. Kuehl and V. V. Zhdankin, *Synlett* **1995**, 1081; (b) V. V. Zhdankin, A. P. Krasutsky, C. J. Kuehl, A. J. Simonsen, J. K. Woodward, B. Mismash and J. T. Bolz, *J. Am. Chem. Soc.* **1996**, 118, 5192; (c) S. A. Moteki, A. Usui, S. Selvakumar, T. Zhang and K. Maruoka, *Angew. Chem. Int. Ed.* **2014**, 53, 11060.
- 10 (a) S. Seo, J. B. Taylor and M. F. Greaney, *Chem. Commun.* **2012**, 48, 8270; (b) S. Seo, M. Slater and M. F. Greaney, *Org. Lett.* **2012**, 14, 2650; (c) J. Kan, S. Huang, J. Lin, M. Zhang and W. Su, *Angew. Chem. Int. Ed.* **2015**, 54, 2199.
- 11 (a) N. Rodriguez and L. J. Goossen, *Chem. Soc. Rev.* **2011**, 40, 5030; (b) D. Katayev, D. Hackenberger and L. J. Goossen, *The Catalyst Review* **2015**, 28, 6; (c) J. Cornella and I. Larrosa, *Synthesis* **2012**, 653; (d) R. Grainger, J. Cornella, D. C. Blakemore, I. Larrosa and J. M. Campanera, *Chem. Eur. J.* **2014**, 20, 16680.
- 12 (a) G. S. Hammond, *J. Am. Chem. Soc.* **1950**, 72, 3737; (b) C. M. M. da Silva Corrêa and M. A. B. C. S. Oliveira, *J. Chem. Soc. Perkin Trans. II*, **1983**, 711.
- 13 (a) R. G. Johnson and R. K. Ingham, *Chem. Rev.* **1956**, 56, 219; (b) The Hunsdiecker and Related Reactions in *Comprehensive Organic Synthesis* (Ed.: B. M. Trost, I. Fleming), vol. 7, Pergamon Press, Oxford, **1991**, p.717.
- 14 Heating (130 °C) of benzoyl hypoiodites (formed *in situ* from silver benzoate and  $\text{I}_2$ ) in chlorobenzene results in the formation of a mixture of biaryl, benzoate ester and benzoic acid products. Similar reactions, but with benzoyl hypobromites, affords only benzoate ester products, see: D. Bryce-Smith, *Nature*, **1953**, 172, 863.
- 15 See supporting information for details of the synthesis and characterisation of **1**.
- 16 The reaction of 2-chlorobenzoic acid under these conditions (no PC,  $\text{Br}_2$  (3.5 equiv)) afforded phenyl 2-chlorobenzoate (12%). No chlorinated arene by-products were formed, only bromobenzene mixtures.
- 17 M. N. Hopkinson, A. Gómez-Suárez, M. Teders, B. Sahoo and F. Glorius, *Angew. Chem. Int. Ed.* **2016**, 55, 4361.
- 18 Rate constants for the decarboxylation of  $4\text{-CH}_3\text{OC}_6\text{H}_4\text{C(O)O}^\bullet$ :  $k = 3.4 \pm 0.3 \times 10^5 s^{-1}$  ( $\text{CCl}_4$ ), ref 8a.
- 19 (a) F. V. Minisci and F. E. Fontane, *Heterocycles* **1989**, 28, 489; (b) F. Minisci, E. Vismara, F. Fontana, G. Morini, M. Serravalle and C. Giordano, *J. Org. Chem.* **1986**, 51, 4411; (c) F. Minisci, F. Fontana and E. Vismara, *J. Heterocycl. Chem.* **1990**, 27, 79.
- 20 G.-X. Li, C. A. Morales-Rivera, Y. Wang, F. Gao, G. He, P. Liu and G. Chen, *Chem. Sci.* **2016**, 7, 6407.
- 21 A. Dewanji, S. Murarka, D. P. Curran and A. Studer, *Org. Lett.* **2013**, 15, 6102.