Tetrahedron 66 (2010) 8623-8628

Contents lists available at ScienceDirect

Tetrahedron

journal homepage: www.elsevier.com/locate/tet

Indium triflate-catalyzed allylation reactions of *N*-sulfonyl aldimines or *N*-alkoxycarbonylamino *p*-tolylsulfones with allyltrimethylsilane: synthesis of protected homoallylic amines

Ponnaboina Thirupathi, Sung Soo Kim*

Department of Chemistry, Inha University, Incheons 402-751, South Korea

ARTICLE INFO

Article history: Received 13 August 2010 Received in revised form 10 September 2010 Accepted 11 September 2010 Available online 17 September 2010

Keywords: N-Sulfonyl aldimines N-Alkoxycarbonylamino p-tolylsulfone Allyltrimethylsilane Allylation Protected homoallylic amines

ABSTRACT

Indium triflate-catalyzed allylation reactions of *N*-sulfonyl aldimines or *N*-alkoxyloxycarbonylamino *p*-tolylsulfone with allyltrimethylsilane have been successfully developed to produce protected homo-allylic amines.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

The allylation of carbonyl derivatives, such as aldehydes, ketones, and imines is one of the important carbon–carbon bond forming reactions because of the versatility of homoallylic alcohols and amines as synthetic intermediates.¹ In particular, Lewis acidcatalyzed allylation of imines has attracted increasing attention because of the significance for the preparation of nitrogen containing molecules.^{1,2} The homoallylic amines are useful fundamental building blocks for the synthesis of many nitrogen containing natural products and bioactive molecules.^{3,4} The development of new methods for the synthesis of homoallylic amines is a significant task for synthetic chemist. A number of methods have been reported for the preparation of protected homoallylic amines using various substrates.^{5,6}

In continuation of the development of useful synthetic methodologies for C–C bond forming reactions⁷ we report herein the Indium triflate-catalyzed allylation reactions of *N*-sulfonyl aldimines or *N*-alkoxycarbonylamino *p*-tolylsulfone with allyltrimethylsilane. In recent years the utility of indium salts as Lewis acid in organic synthesis has received a great deal of attention due to the relatively low toxicity, stability in air and water, strong tolerance to oxygen- and nitrogen-containing substrates.⁸

2. Results and discussions

2.1. Allylation of N-sulfonyl aldimines

Recently, Bateyand Li reported the synthesis of protected homoallylic amines by palladium pincer complex.^{6f} Szabo and co-workers have demonstrated the preparation of protected homoallylic amines with the reaction of *N*-sulfonyl aldimines with potassium trifluoro (allyl)borate.^{6g} The *N*-sulfonyl aldimines (**3**) were prepared from the condensation of aldehydes (**1**), sulfonamides, and sodium salt of arenesulfinic acid to produce the sulfonamidesulfone (**2**) followed by the treatment with saturated aq NaHCO₃ (Scheme 1).⁹

The reactions of *N*-tosyl(phenylbenzylidene)-4-methylbenzenesulfonamide (**3a**) with allyltrimethylsilane (**4**) were examined under reaction conditions employing various catalysts (Table 1). The reaction of *N*-tosyl(phenylbenzylidene)-4-methylbenzenesulfonamide (**3a**) and allyltrimethylsilane (**4**) in the absence of the catalyst did not take place at all (entry 1). The reactions have been screened in the presence of various Lewis acids. Among the various iron and indium catalysts tested, In(OTf)₃ was shown to be the most effective one (entry 11). In(NO₃)₃·XH₂O and In (acac)₃·XH₂O did not allow the reaction to proceed at all (entries 6 and 7). Other metal triflates La(OTf)₃, Ce(OTf)₃, heteropoly acids H₃PO₄·12WO₃·XH₂O, H₃PMo₁₂O₄₀ (PMA), and heterogeneous catalysts Amberlyst-15, PMA·SiO₂ also catalyzed the reaction with only decreased yields (entries 12–17).





^{*} Corresponding author. Fax: +82 32 867 5604; e-mail address: sungsoo@inha. ac.kr (S.S. Kim).

^{0040-4020/\$ -} see front matter @ 2010 Elsevier Ltd. All rights reserved. doi:10.1016/j.tet.2010.09.038



Scheme 1. General preparation method for the synthesis of N-sulfonyl aldimines and synthesis of protected homoallyl amines.

 Table 1

 The reaction conditions for allylation of N-tosyl(phenylbenzylidene)-4-methylbenzenesulfonamide with allyltrimethylsilane^a



Entry	Catalyst (mol %)	Reaction time (h)	Isolated yield (%) ^b
1	_	16	NR
2	$FeCl_3 \cdot 6H_2O(10)$	10	21
3	FeBr ₃ (10)	10	49
4	FeCl ₃ (10)	10	57
5	InCl ₃ (10)	8	80
6	$In(NO_3)_3 \cdot XH_2O(10)$	10	NR
7	$In(acac)_3 \cdot XH_2O(10)$	10	NR
8	InBr ₃ (10)	6	81
9	$In(ClO_4)_3 \cdot XH_2O(10)$	8	46
10	In(OTf) ₃ (10)	6	85
11	$In(OTf)_3(5)$	6	84
12	La(OTf) ₂ (10)	16	76
13	Ce(OTf) ₃ (10)	8	82
14	$H_{3}PO_{4} \cdot 12WO_{3} \cdot XH_{2}O(1)$	8	76
15	H ₃ PMo ₁₂ O ₄₀ (PMA) (1)	6	80
16	Amberlyst-15 (100 mg)	16	56
17	PMA.SiO ₂ (100 mg)	16	74

^a Reaction condition: *N*-Tosyl-(phenylbenzylidene)-4-methylbenzenesulfonamide (1.0 mmol) and allyltrimethylsilane (1.5 mmol) are used.

^b Yield of isolated product after column chromatography.

The $In(OTf)_3$ -catalyzed reactions of various *N*-sulfonyl aldimine with allyltrimethylsilane that proceeded smoothly to give the corresponding protected homoallylic amines (Table 2). The reaction of *N*tosyl(phenylbenzylidene)-4-methylbenzenesulfonamide (**3a**) with allyltrimethylsilane (**4**) produces the protected homoallylic amine (5**a**) in 84% yield (entry 1). The reaction of *N*-sulfonyl aldimines

Table 2

Allylation of various N-sulfonyl aldimines with allyltrimethylsilane catalyzed by In $(\mathrm{OTf})_3{}^\mathrm{a}$

NSO₂Tc ∐	ol-p	SiMe,	$In(OTf)_3 (5 mol\%)$	NHSO ₂ Tol- <i>p</i>
$R \sim H$	+	// ~ 3	CH ₂ Cl ₂	R
3a-n		4	r. t.	5a-n

Entry	R in 3	Reaction time (h)	Product 5	Isolated yield ^b (%)
1	3a , C ₆ H ₄	6.0	5a	84
2	3b , 4-ClC ₆ H ₄	5.0	5b	88
3	3c , 3-ClC ₆ H ₄	6.0	5c	85
4	3d, 2-ClC ₆ H ₄	6.0	5d	82
5	3e , 4-BrC ₆ H ₄	6.0	5e	89
6	3f , 4-MeC ₆ H ₄	5.0	5f	87
7	3g , 4-MeOC ₆ H ₄	4.0	5g	88
8	3h , 3,4,5-(MeO) ₃ C ₆ H ₂	3.5	5h	91
9	3i , 4-NO ₂ C ₆ H ₄	6.0	5i	78
10	3j , 4-CNC ₆ H ₄	6.0	5j	82
11	3k , 3-CNC ₆ H ₄	6.0	5k	76

^a Reaction condition: *N*-sulfonyl aldimine (1.0 mmol), trimethylsilane (1.5 mmol), and In(OTf)3 (5 mol % with respective to *N*-sulfonyl aldimine) are used.

^b Yield of isolated product after column chromatography.

containing halogen groups, such as chlorine and bromine on the phenyl ring proceeded smoothly to give the corresponding products in good yields (entries 2–5). *N*-Tosyl(4-methylbenzylidene) (**3f**), *N*-tosyl(4-methoxybenzylidene) (**3g**), and *N*-tosyl(3,4,5-trimethoxybenzylidene)-4-methylbenz-enesulfonamide (**3h**) react with allyl-trimethylsilane (**4**) to give **5f**, **5g**, and **5h** in 87, 88, and 91%, respectively (entries 6–8). The reactions of *N*-sulfonyl aldimines bearing an electron-withdrawing group on the phenyl ring took place in slightly lower yields (entries 9–11).

2.2. Allylation of N-alkoxycarbonylamino p-tolylsulfone

The acylated homoallylic amines are important synthons for many synthetic applications, notably for acyliminum chemistry.¹⁰The varieties of synthetic methods have been reported for the synthesis of homoallylic amines. The in situ generations of imines are highly attractive, among which the three-component allylation reaction have been introduced by Panek^{5a} and developed by Veenstra.^{5b}Yet, many Lewis acids cannot be used in this reaction because they decompose or are deactivated in the presence of the amine or water produced during imine formation. Thus the Sakurai reaction of *N*-alkoxy-carbonylamino *p*-tolylsulfone (**6**) with allyltrimethylsilane (**4**) in the presence of a suitable Lewis acid constitutes an important method for the synthesis of *N*-alkoxycarbonyl-protected homoallylic amines. This reaction proceeds through a very reactive *N*-acyliminium species or *N*-alkoxycarbonyliminium derivative (**7**, Scheme 2).

Recently, very few Lewis acid catalysts, such as $TiCl_4$, $SnCl_4$, $GaCl_3$, $InCl_3$, or $Bi(OTf)_3$ have been employed for the preparation of alkoxycarbonyl-protected homoallylic amines.¹¹ However, $TiCl_4$ or $SnCl_4$ is moisture sensitive and required in stoichiometric quantity or in large excess. Some of these methods show longer reaction time and lower yield. To overcome these limitations, we report herein In (OTf)₃-catalyzed Sakurai reaction of *N*-alkoxycarbonylamino *p*-tolylsulfone with allyltrimethylsilane. Under optimized reaction conditions as in Table 1, the scope of allylation reactions have been studied using *N*-alkoxycarbonylamino *p*-tolylsulfone (**6**) and allyltrimethylsilane (**4**). The results are summarized in Table 3.

N-Alkoxycarbonylamino p-tolylsulfones (6) with electron-donating group attached to the benzene ring react with allyltrimethylsilane (**4**) to produce the Cbz-protected homoallylic amines in good yield (entries 2-6). The reactions of p-, m-, and ohalogen containing N-alkoxycarbonylamino p-tolylsulfone gave corresponding products in 90, 83, 89, 87, 84, and 83% yield, respectively (entries 7-12). This shows no serious electron-withdrawing effects of halogen atoms. o-Isomers exhibit a slight steric and electron effect, that is responsible for the less yield and longer reaction time (entries 8 and 11). The electron-withdrawing p-CN group also (**6k**) proceeds smoothly to give the allylation product in good yield (entry 13). 2-Furyl (6n) and 2-naphthyl N-alkoxycarbonylamino *p*-tolylsulfone (**60**) react with allyltrimethylsilane (4) to afford the corresponding products (entries 14 and 15). The reactions of acyclic aliphatic N-alkoxycarbonylamino p-tolylsulfone give the Cbz-protected homoallylic amines in comparatively low yield (entries 16 and 17). Cyclohexyl and 2-cyclohexenyl N-alkoxycarbonylamino p-tolylsulfone bring about the corresponding allylation derivatives (entry 18 and 19).



Scheme 2. General equation for Lewis acid-catalyzed allylation of N-benzyloxycarbonylamino p-tolylsulfone.

Table 3

Allylation of various N-alkoxycarbonylamino p-tolylsulfone with allyltrimethylsilane catalyzed by $In(OTf)_3^a$



Entry	R in 6	Reaction time (h)	Product 8	Isolated yield (%) ^b
1	6a , C ₆ H ₄	8.0	8a	87
2	6b , 4-MeC ₆ H ₄	9.0	8b	89
3	6c , 2-MeC ₆ H ₄	9.0	8c	85
4	6d , 4-MeOC ₆ H ₄	8.0	8d	91
5	6e , 3,4-(MeO) ₃ C ₆ H ₂	8.0	8e	91
6	6f , 3-PhOC ₆ H ₄	9.0	8f	87
7	6g , 4-BrC ₆ H ₄	9.0	8g	92
8	6h , 2-BrC ₆ H ₄	9.0	8h	83
9	6i , 4-ClC ₆ H ₄	8.0	8i	89
10	6j , 3-ClC ₆ H ₄	9.0	8j	87
11	6k , 2-ClC ₆ H ₄	9.0	8k	84
12	61 , 3-FC ₆ H ₄	9.0	81	83
13	6m , 4-CNC ₆ H ₄	9.0	8m	80
14	6 n,	9.0	8n	76
15	60,	9.0	80	84
16	60 $CH_2(CH_2)_2$	10.0	8n	70
17	6a . $CH_2(CH_2)_c$	10.0	8a	76
18	6r, 6r,	10.0	8r	75
19	6s,	10.0	8s	78

^a Reaction condition: *N*-alkoxycarbonylamino *p*-tolylsulfone (**6**) (1.0 mmol), trimethylsilane (1.5 mmol), and In(OTf)3 (5 mol % with respective to *N*-sulfonyl aldimine) are used. ^b Yield of isolated product after column chromatography.

3. Conclusion

An effective Indium triflate-catalyzed allylation reaction of *N*-sulfonyl aldimines or *N*-alkoxycarbonylamino *p*-tolylsulfone with allyltrimethylsilane is described. The present method provides a facile and convenient route for the synthesis of protected homoallylic amines. The advantages of the present method include: (1) low catalyst loading and (2) mild reaction conditions.

4. Experimental section

4.1. Experimental procedure for the allylation of *N*-sulfonyl aldimines with allyltrimethylsilane

To a stirred solution of *N*-sulfonyl aldimines (1 mmol) and In $(OTf)_3$ (5 mol %) in CH₂Cl₂ (3 mL) under nitrogen atmosphere at rt, allyltrimethylsilane (1.5 mmol) is added. The mixture is stirred at rt

and monitored by TLC. After the completion, the reaction mixture is filtered, and the residue is washed with CH_2Cl_2 (2×5 mL). The filtrate is dried over anhydrous Na_2SO_4 and concentrated under vacuum. The crude product is subject to flash column chromatography (silica gel, hexane–EtOAc, 9:1) to obtain the pure product.

4.2. Experimental procedure for the allylation of *N*-alkoxycarbonylamino *p*-tolylsulfone with allyltrimethylsilane

Allyltrimethylsilane (1.5 mmol) was added drop-wise to a stirred solution of *N*-alkoxycarbonylamino *p*-tolylsulfone (1 mmol) and In(OTf)₃ (5 mol %) in CH₂Cl₂ (3 mL) under nitrogen atmosphere at rt, The mixture is stirred at rt and monitored by TLC. After the completion, the reaction mixture is filtered, and the residue is washed with CH₂Cl₂ (2×5 mL). The filtrate is dried over anhydrous Na₂SO₄ and concentrated under vacuum. The crude product is subject to flash column chromatography (silica gel, hexane–EtOAc, 90:10 to 95:5) to obtain the pure product. The products were characterized by ¹H and ¹³C NMR data that are consistent with literature values. Melting point, HRMS-FAB, ¹H, and ¹³C NMR values for new products are given below.

4.2.1. Compound **5a**. White solid, mp 75–76 °C; IR (KBr): ν_{max} 3261, 1458, 1314, 1159, 1088 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.51 (d, *J*=8.4 Hz, 2H), 7.14–7.01 (m, 7H), 5.50–5.43 (m, 1H), 5.11 (br s, 1H), 4.99 (d, *J*=14.0 Hz, 2H), 4.33 (q, *J*=7.0 Hz, 1H), 2.46–2.36 (m, 2H), 2.32 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 143.0, 140.3, 137.4, 133.1, 129.2, 128.3, 127.3, 127.1, 126.5, 119.1, 57.2, 41.8, 21.4.^{5c,6g,1}

4.2.2. Compound **5b**. White solid, mp 159–160 °C; IR (KBr): ν_{max} 3261, 1456, 1314, 1159, 1076 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.53 (d, *J*=8.4 Hz, 2H), 7.13–7.07 (m, 4H), 6.98 (d, *J*=8.0 Hz, 2H), 5.51–5.44 (m, 2H), 5.03 (s, 1H), 5.00–4.98 (m, 1H), 4.32 (q, *J*=7.0 Hz, 1H), 2.43–2.36 (m, 2H), 2.38 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 143.2, 138.8, 137.2, 132.9, 132.7, 129.2, 128.3, 128.0, 127.0, 119.2, 56.7, 41.5, 21.3.^{5c,6l}

4.2.3. Compound **5c**. Viscous mass; IR (KBr): ν_{max} 3273, 1589, 1422, 1291, 1148, 1088 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.54 (d, *J*=8.4 Hz, 2H), 7.14 (d, *J*=8.4 Hz, 2H), 7.09 (d, *J*=6.0 Hz, 2H), 7.02–6.98 (m, 1H), 6.94 (s, 1H), 5.54–5.44 (m, 2H), 5.06 (s, 1H), 5.02 (d, *J*=5.8 Hz, 1H), 4.34 (q, *J*=7.0 Hz, 1H), 2.44–2.38 (m, 2H), 2.36 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 143.2, 142.3, 137.1, 134.0, 132.6, 129.5, 129.2, 127.3, 127.0, 126.8, 124.8, 119.4, 56.8, 41.5, 21.3; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₇H₁₉ClNO₂S 336.0825, found 336.0825.

4.2.4. Compound **5d**. White solid, mp 113–114 °C; IR (KBr): ν_{max} 3261, 1446, 1327, 1159, 1086 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.60 (d, *J*=8.4 Hz, 2H), 7.21–7.17 (m, 2H), 7.12 (d, *J*=7.0 Hz, 2H), 7.08–7.05 (m, 2H), 5.58–5.44 (m, 2H), 5.03 (d, *J*=15 Hz, 2H), 4.83 (q, *J*=7.0 Hz, 1H), 2.48–2.38 (m, 2H), 2.34 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 143.1, 137.8, 136.8, 132.7, 131.9, 129.4, 129.2, 128.3, 128.2, 127.1, 126.7, 119.3, 54.0, 40.3, 21.4; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₇H₁₉ClNO₂S 336.0825, found 336.0826.

4.2.5. *Compound* **5***e*. White solid, mp 129–130 °C; IR (KBr): ν_{max} 3249, 1446, 1327, 1159, 1064 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.52 (d, *J*=8.4 Hz, 2H), 7.25 (d, *J*=8.4 Hz, 2H), 7.13 (d, *J*=8.4 Hz, 2H), 6.93 (d, *J*=8.4 Hz, 2H), 5.54–5.45 (m, 1H), 5.34 (d, *J*=6.6 Hz, 1H), 5.05 (s, 1H), 5.02 (d, *J*=8.4 Hz, 1H), 4.32 (q, *J*=7.0 Hz, 1H), 2.43–2.38 (m, 2H), 2.39 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 143.3, 139.3, 137.2, 132.6, 131.3, 129.3, 128.3, 127.0, 121.1, 119.4, 56.6, 41.5, 21.4; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₇H₁₉BrNO₂S 380.0143, found 380.0137.

4.2.6. Compound **5***f*. White solid, mp 118–119 °C; IR (KBr): ν_{max} 3261, 1446, 1314, 1159, 1088 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.55 (d, *J*=8.4 Hz, 2H), 7.14 (d, *J*=8.7 Hz, 2H), 7.01–6.93 (m, 4H), 5.55–5.45 (m, 1H), 5.02 (dd, *J*=12.0, 2.0 Hz, 3H), 4.31 (q, *J*=7.0 Hz, 1H), 2.49–2.39 (m, 2H), 2.37 (s, 3H), 2.26 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 142.9, 137.5, 137.3, 137.0, 129.6, 129.2, 128.9, 127.1, 126.4, 119.0, 56.9, 41.7, 21.4, 20.9.^{5c,6g,1}

4.2.7. Compound **5g**. White solid, mp 86–87 °C; IR (KBr): ν_{max} 3261, 1530, 1314, 1159, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.52 (d, *J*=7.8 Hz, 2H), 7.09 (d, *J*=8.0 Hz, 2H), 6.94 (d, *J*=8.8 Hz, 2H), 6.64 (d, *J*=8.4 Hz, 2H), 5.49–5.43 (m, 1H), 5.24 (br s, 1H), 4.99 (d, *J*=6.6 Hz, 2H), 4.27 (q, *J*=7.0 Hz, 1H), 3.69 (s, 3H), 2.44–2.36 (m, 2H), 2.32 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 158.7, 142.8, 137.5, 133.3, 132.4, 131.9, 129.2, 127.7, 127.0, 118.7, 113.6, 56.8, 55.1, 41.7, 21.3.^{5c}

4.2.8. Compound **5h**. White solid, mp 67–68 °C; IR (KBr): ν_{max} 3257, 1458, 1314, 1159, 1111 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.50 (d, *J*=7.8 Hz, 2H), 7.08 (d, *J*=7.8 Hz, 2H), 6.21 (s, 2H), 5.58–5.52 (m, 2H), 5.05 (d, *J*=8.4 Hz, 1H), 5.02 (s, 1H), 4.30 (q, *J*=7.0 Hz, 1H), 3.72 (s,

3H), 3.67 (s, 6H), 2.45–2.39 (m, 2H), 2.32 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 152.8, 142.9, 137.6, 136.8, 135.8, 133.2, 129.5, 129.0, 127.1, 126.3, 118.8, 103.5, 60.4, 57.7, 55.8, 41.7, 21.3; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₂₀H₂₆NO₅S 391.1453, found 391.1453.

4.2.9. Compound **5i**. White solid, mp 127 °C; IR (KBr): ν_{max} 3285, 1518, 1362, 1148, 1064 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.00 (d, *J*=8.8 Hz, 2H), 7.55 (d, *J*=8.4 Hz, 2H), 7.28 (d, *J*=8.8 Hz, 2H), 7.14 (d, *J*=8.0 Hz, 2H), 5.60 (d, *J*=6.2 Hz, 1H), 5.50–5.48 (m, 1H), 5.04 (d, *J*=5.5 Hz, 1H), 5.02 (d, *J*=12.5 Hz, 1H), 4.44 (q, *J*=7.0 Hz, 1H), 2.44–2.39 (m, 2H), 2.35 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 147.9, 146.9, 143.8, 131.9, 129.4, 127.5, 127.0, 123.4, 120.1, 56.6, 41.4, 21.3.^{6g}

4.2.10. Compound **5***j*. White solid, mp 148–149 °C; IR (KBr): ν_{max} 3297, 2210, 1338, 1135, 1064 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.51 (d, *J*=8.4 Hz, 2H), 7.39 (d, *J*=7.8 Hz, 2H), 7.17 (d, *J*=8.0 Hz, 2H), 7.10 (d, *J*=8.0 Hz, 2H), 5.65 (d, *J*=6.6 Hz, 1H), 5.42–5.39 (m, 1H), 4.99 (d, *J*=4.7 Hz, 1H), 4.97 (d, *J*=11.7 Hz, 1H), 4.35 (q, *J*=7.0 Hz, 1H), 2.37–2.33 (m, 2H), 2.34 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.9, 143.5, 136.9, 132.0, 129.3, 127.4, 127.0, 119.8, 118.5, 110.9, 56.8, 41.3, 21.3.

4.2.11. Compound **5k**. White solid, mp 84–85 °C; IR (KBr): ν_{max} 3249, 2210, 1327, 1159, 1052 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.49 (d, *J*=8.4 Hz, 2H), 7.39–7.34 (m, 2H), 7.27–7.22 (m, 2H), 7.10 (d, *J*=8.4 Hz, 2H), 5.60 (br s, 1H), 5.47–5.39 (m, 1H), 5.03–4.96 (m, 2H), 4.35 (q, *J*=6.7 Hz, 1H), 2.40–2.35 (m, 2H), 2.36 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 143.5, 141.9, 137.0, 132.1, 131.2, 130.7, 130.2, 129.4, 129.0, 126.9, 119.8, 118.3, 112.1, 56.6, 41.4, 21.3; HRMS-FAB (*m*/*z*): [M+H]⁺ calcd for C₁₈H₁₉N₂O₂S: 327.1167, found: 327.1165.

4.2.12. Compound **8a**. White solid, mp 67 °C; IR (KBr): ν_{max} 3367, 1685, 1530, 1257, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.31–7.22 (m, 10H), 5.70–5.59 (m, 1H), 5.20 (br s, 1H), 5.10–5.01 (m, 4H), 4.79 (br s, 1H), 2.51 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 155.7, 142.0, 136.5, 133.8, 128.6, 128.5, 128.1, 127.3, 126.3, 118.4, 66.8, 54.6, 41.1.^{5g}

4.2.13. Compound **8b**. White solid, mp 63–64 °C; IR (KBr): ν_{max} 3357, 1697, 1518, 1267, 1040 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.27 (s, 5H), 7.11–7.07 (m, 5H), 5.68–5.58 (m, 1H), 5.32 (br s, 1H), 5.07–4.98 (m, 4H), 4.74 (br s, 1H), 2.46 (s, 2H), 2.28 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 155.6, 138.9, 136.7, 136.4, 133.8, 129.1, 128.3, 127.9, 126.1, 118.1, 66.5, 54.2, 40.9, 20.9.^{11f}

4.2.14. Compound **8c**. White solid, mp 51–52 °C; IR (KBr): ν_{max} 3344, 1673, 1541, 1267, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.35 (s, 5H), 7.25–7.18 (m, 5H), 5.80–5.70 (m, 1H), 5.23 (br s, 1H), 5.17–5.04 (m, 5H), 2.52 (s, 2H), 2.43 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 155.7, 140.3, 136.5, 135.4, 134.0, 130.7, 128.5, 128.1, 127.2, 126.3, 125.1, 118.3, 66.8, 50.8, 40.3, 19.3.^{11h}

4.2.15. Compound **8d**. White solid, mp 69–70 °C; IR (KBr): ν_{max} 3309, 1684, 1541, 1255, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.30 (s, 5H), 7.16 (d, *J*=7.8 Hz, 2H), 6.83 (d, *J*=8.4 Hz, 2H), 5.70–5.59 (m, 1H), 5.14 (br s, 1H), 5.10–5.00 (m, 4H), 4.72 (br s, 1H), 3.76 (s, 3H), 2.49 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 158.7, 155.6, 136.4, 133.8, 128.4, 128.0, 127.3, 118.2, 113.8, 66.6, 55.1, 54.0, 40.9.^{5g}

4.2.16. Compound **8e**. White solid, mp 70 °C; IR (KBr): ν_{max} 3344, 1697, 1530, 1243, 1124 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.35–7.25 (m, 5H), 6.46 (s, 2H), 5.73–5.62 (m, 1H), 5.18–5.08 (m, 6H), 4.75 (br s, 1H), 3.81 (s, 9H), 2.51 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 156.3, 153.8, 138.3, 137.6, 136.9, 134.3, 129.0, 128.6, 119.0, 103.7, 67.4, 61.3, 56.6, 55.3, 41.7.^{11f}

4.2.17. Compound **8f**. White solid, mp 45–46 °C; IR (KBr): v_{max} 3311, 1679, 1534, 1255, 1056 cm⁻¹; ¹H NMR (400 MHz, CDCl₃)

 δ 7.36–7.28 (m, 7H), 7.14–7.10 (m, 1H), 7.03–6.97 (m, 4H), 6.90 (d, $J{=}8.0$ Hz, 2H), 5.73–5.64 (m, 1H), 5.19 (br s, 1H), 5.14–5.05 (m, 4H), 4.82 (br s, 1H), 2.48 (s, 2H); 13 C NMR (100 MHz, CDCl₃) δ 157.3, 157.0, 155.6, 144.1, 136.3, 133.4, 129.8, 129.7, 128.4, 128.0, 123.2, 121.0, 118.8, 118.5, 117.4, 116.6, 66.7, 54.1, 40.9; HRMS-FAB (m/z) [M+H]⁺ calcd for C₂₄H₂₄NO₃ 373.1756, found 374.1755.

4.2.18. Compound **8g**. White solid, mp 87–88 °C; IR (KBr): ν_{max} 3344, 1684, 1541, 1267, 1040 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.45 (d, *J*=8.4 Hz, 2H), 7.33 (s, 5H), 7.14 (d, *J*=7.0 Hz, 2H), 5.68–5.60 (m, 1H), 5.20 (br s, 1H), 5.13–5.06 (m, 4H), 4.75 (br s, 1H), 2.49 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 155.6, 141.0, 136.2, 133.2, 131.6, 128.5, 128.1, 127.9, 121.0, 118.8, 66.8, 53.9, 40.8; HRMS-FAB (*m/z*) [M+H]⁺ calcd for C₁₈H₁₉BrNO₂ 360.0599, found 360.0597.

4.2.19. Compound **8h**. White solid, mp 73 °C; IR (KBr): ν_{max} 3311, 1693, 1547, 1268, 1030 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.55 (d, *J*=8.0 Hz, 2H), 7.34–7.24 (m, 6H), 7.14–7.09 (m, 1H), 5.75–5.64 (m, 1H), 5.31 (br s, 1H), 5.16–5.03 (m, 5H), 2.56 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 155.5, 141.0, 136.3, 133.5, 133.2, 128.7, 128.5, 128.2, 127.6, 127.3, 122.7, 118.7, 66.8, 54.0, 39.4; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₈H₁₉BrNO₂ 360.0599, found 360.0597.

4.2.20. Compound **8i**. White solid, mp 63 °C; IR (KBr): ν_{max} 3347, 1675, 1543, 1267, 1040 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.27 (m, 6H), 7.17 (d, *J*=7.0 Hz, 2H), 5.68–5.60 (m, 1H), 5.23 (br s, 1H), 5.17–5.01 (m, 4H), 4.78 (br s, 1H), 2.51 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 156.2, 141.3, 136.9, 133.8, 133.6, 129.2, 129.1, 121.0, 128.9, 128.7, 128.2, 128.1, 119.4, 67.4, 54.5, 41.4.^{11h}

4.2.21. Compound **8***j*. White solid, mp 61–62 °C; IR (KBr): ν_{max} 3344, 1673, 1530, 1255, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.34–7.24 (m, 8H), 7.15 (d, *J*=2.0 Hz, 1H), 5.73–5.60 (m, 1H), 5.26 (br s, 1H), 5.14–5.04 (m, 4H), 4.89 (br s, 1H), 2.50 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 155.6, 144.2, 136.2, 134.3, 133.1, 129.7, 128.4, 128.1, 127.4, 126.3, 124.4, 118.8, 66.8, 54.0, 40.8; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₈H₁₉ClNO₂ 316.1104, found 316.1105.

4.2.22. Compound **8**k. White solid, mp 65–66 °C; IR (KBr): v_{max} 3309, 1673, 1541, 1267, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.37–7.17 (m, 8H), 5.75–5.63 (m, 1H), 5.33 (br s, 1H), 5.21–5.03 (m, 5H), 2.54 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 156.1, 140.1, 136.9, 134.0, 133.7, 130.5, 129.0, 128.7, 127.9, 127.5, 119.4, 67.4, 52.6, 39.8.^{5g,h}

4.2.23. Compound **8**I. Viscous mass; IR (KBr): ν_{max} 3320, 1684, 1530, 1243, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.31–7.22 (m, 6H), 7.03 (d, *J*=7.0 Hz, 1H), 6.97–6.90 (m, 2H), 5.68–5.90 (m, 1H), 5.26 (d, *J*=5.4 Hz, 1H), 5.11–5.02 (m, 4H), 4.79 (br s, 1H), 2.48 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 164.1, 161.6, 155.6, 144.7, 136.2, 133.2, 130.0, 129.9, 128.4, 128.1, 121.8, 118.7, 114.2, 113.9, 113.2, 113.0, 66.8, 54.0, 40.8; HRMS-FAB (*m/z*) [M+H]⁺ calcd for C₁₈H₁₉FNO₂ 300.1400, found 300.1398.

4.2.24. Compound **8m**. Viscous mass; IR (KBr): ν_{max} 3366, 2208, 1689, 1540, 1255, 1040 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.59 (d, *J*=7.8 Hz, 2H), 7.37–7.32 (m, 1H), 5.66–5.56 (m, 1H), 5.41 (d, *J*=5.8 Hz, 1H), 5.13–5.05 (m, 4H), 4.82 (br s, 1H), 2.48 (d, *J*=4.8 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 155.5, 147.5, 136.0, 132.6, 132.2, 128.4, 128.1, 128.0, 126.9, 119.2, 118.6, 110.9, 66.8, 54.2, 40.5.^{11h}

4.2.25. Compound **8n**. Viscous mass; IR (KBr): ν_{max} 3333, 1697, 1506, 1219, 1040 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.30 (s, 6H), 6.26 (dd, *J*=3.0, 1.0 Hz, 1H), 6.13 (s, 1H), 5.71–5.61 (m, 1H), 5.09–5.03 (m, 5H), 4.89 (br s, 1H), 2.55 (t, *J*=6.7 Hz, 2H); ¹³C NMR

(100 MHz, CDCl₃) δ 154.9, 153.4, 141.2, 135.7, 132.6, 127.8, 127.4, 117.9, 109.5, 105.5, 66.2, 48.0, 37.8. $^{\rm 5g}$

4.2.26. Compound **80**. White solid, mp 66–67 °C; IR (KBr): ν_{max} 3368, 1673, 1518, 1243, 1028 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.75–7.66 (m, 3H), 7.42–7.28 (m, 6H), 5.68–5.59 (m, 1H), 5.41 (br s, 1H), 5.09–4.95 (m, 5H), 2.54 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 155.7, 139.3, 136.3, 133.6, 133.2, 132.6, 128.4, 128.3, 128.0, 127.8, 127.5, 126.1, 125.7, 124.8, 124.4, 118.4, 66.7, 54.6, 40.8.^{11f}

4.2.27. Compound **8p**. Viscous mass; IR (KBr): ν_{max} 3320, 1697, 1530, 1219, 1088 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.32 (s, 5H), 5.82–5.74 (m, 1H), 5.17–5.06 (m, 4H), 4.78 (br s, 1H), 3.64–3.59 (m, 1H), 2.36–2.23 (m, 1H), 2.18–2.13 (m, 1H), 1.42–1.38 (m, 2H), 0.99–0.93 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 156.3, 136.8, 134.8, 128.5, 128.0, 117.4, 66.5, 55.2, 41.4, 36.7, 29.7, 28.3, 26.4, 26.2, 26.1.^{5d}

4.2.28. Compound **8***q*. Viscous mass; IR (KBr): ν_{max} 3320, 1684, 1541, 1267, 1064 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.33 (s, 5H), 5.82–5.74 (m, 1H), 5.66 (s, 2H), 5.14–5.05 (m, 4H), 4.67 (br s, 1H), 3.68–3.64 (m, 1H), 2.43–2.30 (m, 1H), 2.22–2.14 (m, 1H), 2.08–2.00 (m, 3H), 1.85–1.75 (m, 2H), 1.67–1.63 (m, 1H), 1.48–1.20 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 156.2, 136.5, 134.3, 128.4, 127.9, 126.9, 125.9, 117.7, 66.5, 54.4, 37.4, 36.7, 28.6, 26.9, 25.6, 25.2, 24.3; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₈H₂₄NO₂ 290.2120, found 290.2120.

4.2.29. Compound **8r**. White solid, mp 64–65 °C; IR (KBr): ν_{max} 3309, 1673, 1541, 1255 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.39–7.28 (s, 5H), 5.82–5.76 (m, 1H), 5.09–5.05 (m, 4H), 4.67 (br s, 1H), 3.68–3.58 (m, 1H), 2.31–2.13 (m, 3H), 1.80–1.76 (m, 1H), 0.87–0.77 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 156.0, 136.7, 134.8, 134.3, 128.5, 128.0, 117.8, 66.5, 50.4, 39.5, 36.8, 19.1, 13.9.^{3e,5h}

4.2.30. Compound **8s**. Viscous mass; IR (KBr): ν_{max} 3320, 1684, 1518, 1243, 1040 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.36–7.28 (m, 5H), 5.82–5.73 (m, 1H), 5.10–5.04 (m, 4H), 4.74 (br s, 1H), 3.71 (br s, 1H), 2.28–2.16 (m, 2H), 1.47 (br s, 1H), 1.44–1.26 (m, 9H), 0.89 (t, *J*=7.0 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 155.9, 136.6, 134.2, 128.3, 128.0, 127.8, 117.6, 66.3, 50.6, 39.3, 34.5, 31.6, 29.0, 25.7, 22.4, 13.9; HRMS-FAB (*m*/*z*) [M+H]⁺ calcd for C₁₈H₂₈NO₂ 286.1807, found 286.1806.

Acknowledgements

Authors thank the Centre for Biological Modulators of the KRICT for the financial support. We also acknowledge the assistance by BK21 provided by Korea Research Council.

References and notes

- 1. Yamamoto, Y.; Asao, N. Chem. Rev. 1993, 93, 2207-2293.
- 2. Kobayashi, S.; Ishitani, H. Chem. Rev. 1999, 99, 1069-1094.
- (a) Yamaguchi, R.; Moriyasu, M.; Kawanisi, M. J. Org. Chem. **1985**, 50, 287–288;
 (b) Yamamoto, Y.; Schmid, M. J. Chem. Soc., Chem. Commun. **1989**, 1310–1312;
 (c) Ovaa, H.; Stragies, R.; Van der Marel, G. A.; Van Boom, J. H.; Blechert, S. Chem. Commun. **2000**, 1501–1502;
 (d) Hunt, J. C. A.; Laurent, P.; Moody, C. J. Chem. Commun. **2000**, 1771–1772;
 (e) Felpin, F.-X.; Girard, S.; Vo-Thanh, G.; Robins, R. J.; Villieras, J.; Lebreton, J. J. Org. Chem. **2001**, 66, 6305–6312.
- (a) Enders, D.; Reinhold, U. Tetrahedron: Asymmetry 1997, 8, 1895–1946; (b) Bloch, R. Chem. Rev. 1998, 98, 1407–1438; (c) Wright, D. L.; Schulte, J. P., II; Page, M. A. Org. Lett. 2000, 2, 1847–1850; (d) Puentes, C. O.; Kouznetsov, V. J. Heterocycl. Chem. 2002, 39, 595–614; (e) Jain, R. P.; Williams, R. M. J. Org. Chem. 2002, 67, 6361–6365; (f) Ramachandran, P. V.; Burghardt, T. E.; Bland-Berry, L.J. Org. Chem. 2005, 70, 7911–7918; (g) Besada, P.; Mamedova, L.; Thomas, C. J.; Costanzi, S.; Jacobson, K. A. Org. Biomol. Chem. 2005, 3, 2016–2025.
- (a) Panek, J. S.; Jain, N. F. J. Org. Chem. **1994**, 59, 2674–2675; (b) Veenstra, S. J.; Schmid, P. Tetrahedron Lett. **1997**, 38, 997–1000; (c) Masuyama, Y.; Tosa, J.; Kurusu, Y. Chem. Commun. **1999**, 1075–1076; (d) Hunt, J. C. A.; Laurent, P.; Moody, C. J. J. Chem. Soc., Perkin Trans. 1 **2002**, 2378–2389; (e) Ollevier, T.; Ba, T.

Tetrahedron Lett. **2003**, *44*, 9003–9005; (f) Smitha, G.; Miriyala, B.; Williamson, J. S. *Synlett* **2005**, 839–841; (g) Phukan, P. *J. Org. Chem.* **2004**, 69, 4005–4006; (h) Song, Q.-Y.; Yang, B.-L.; Tian, S.-K. *J. Org. Chem.* **2007**, *72*, 5407–5410.

- (a) Killinger, T. A.; Boughton, N. A.; Runge, T. A.; Wolinsky, J.J. Organomet. Chem. 1977, 124, 131–134; (b) Sun, X.-W.; Xu, M.-H.; Lin, G.-Q. Org. Lett. 2006, 8, 4979–4982; (c) Vilaivan, T.; Winotapan, C.; Banphavichit, V.; Shinada, T.; Ohfune, Y. J. Org. Chem. 2005, 70, 3464–3471; (d) Vilaivan, T.; Winotapan, C.; Shinada, T.; Ohfune, Y. Tetrahedron Lett. 2001, 42, 9073–9076; (e) Legros, J.; Meyer, F.; Coliboeuf, M.; Crousse, B.; Bonnet-Delpon, D.; Begue, J.-P.J. Org. Chem. 2003, 68, 6444–6446; (f) Li, S.-W.; Batey, A. R. Chem. Commun. 2004, 1382–1383; (g) Solin, N.; Wallner, O. A.; Szabo, K. J. Org. Lett. 2005, 7689–691; (h) Fan, R.; Pu, D.; Wen, F.; Wu, J.J. Org. Chem. 2008, 73, 3623–3625; (j) Kallstrom, S.; Saloranta, T.; Minnaard, A. J.; Leino, R. Tetrahedron Lett. 2007, 48, 6958–6961; (k) Li, J.; Minnaard, A. J.; Klein Gebbink, R. J. M.; Van Koten, G. Tetrahedron Lett. 2009, 50, 2232–2235; (l) Fan, R.; Li, W.; Pu, D.; Zhang, L. Org. Lett. 2009, 11, 1425–1428.
- (a) Thirupathi, P.; Kim, S. S. Tetrahedron 2010, 66, 2995–3003; (b) Thirupathi, P.; Kim, S. S. J. Org. Chem. 2009, 74, 7755–7761; (c) Thirupathi, P.; Kim, S. S. Eur. J. Org. Chem. 2010, 1798–1808; (d) Kadam, S. T.; Thirupathi, P.; Kim, S. S. Tetrahedron 2010, 66, 1684–1688; (e) Kadam, S. T.; Thirupathi, P.; Kim, S. S. Tetrahedron 2009, 65, 10383–10389; (f) Kadam, S. T.; Kim, S. S. Appl. Organomet. Chem. 2009, 23, 119–123; (g) Rajagopal, G.; Kim, S. S. Tetrahedron

2009, 65, 4351–4355; (h) Thirupathi, P.; Kim, S. S. Tetrahedron **2009**, 65, 5168–5173.

- (a) Prajapati, D.; Laskar, D. D.; Sadhu, J. S. Tetrahedron Lett. 2000, 41 8639–8643;
 (b) Frost, C. G.; Chauhan, K. K. J. Chem. Soc., Perkin Trans. 1 2000, 3015–3019; (c) Loh, T.-P.; Hu, Q.-Y.; Tan, K.-T.; Cheng, H.-S. Org. Lett. 2001, 3, 2669–2672; (d) Fringuelli, F.; Piermatti, O.; Pizzo, F.; Vaccaro, L. Curr. Org. Chem. 2003, 7, 1661–1689; (e) Cosh, R.; Maiti, S.; Chakraborty, A.; Maiti, D. K. J. Mol. Catal. A: Chem. 2004, 210, 53–57; (f) Frost, C. G.; Hartley, J. P. Mini-Rev. Org. Chem. 2004, 1, 1–7; (g) Zhang, Z.-H. Synlett 2005, 711–712; (h) Yahada, R.; Obika, S.; Kono, H.; Takemoto, Y. Angew. Chem., Int. Ed. 2006, 45, 3822–3825; (i) Takahashi, K.; Midori, M.; Kawano, K.; Ishihara, J.; Hatekeyama, S. Angew. Chem., Int. Ed. 2008, 47, 6244–6246; (j) Yadav, J. S.; Antony, A.; Goerge, J.; Subba Reddy, B. V. Eur. J. Org. Chem. 2010, 595–605.
- 9. Chemla, F.; Hebbe, V.; Normant, J.-F. Synthesis 2000, 75-77.
- Hieman, H.; Speckamp, W. N. In *Comprehensive Organic Synthesis*; Fleming, I., Ed.; Pergamon: Oxford, 1991; Vol. 2, p 1047.
- (a) Giardina, A.; Mecozzi, T.; Petrini, M. J. Org. Chem. 2000, 65, 8277–8282; (b) Schunk, S.; Enders, D. Org. Lett. 2001, 3, 3177–3180; (c) Enders, D.; Oberbcrsch, S. Synlett 2002, 471–473; (d) Petrini, M.; Torregiani, E. Tetrahedron Lett. 2005, 46, 5999–6003; (e) Ollevier, T.; Li, Z. Org. Biomol. Chem. 2006, 4, 4440–4443; (f) Das, B.; Damodar, K.; Saritha, D.; Chowdhury, N.; Krishnaiah, M. Tetrahedron Lett. 2007, 48, 7930–7933; (g) Kumar, R. S. C.; Reddy, G. V.; Babu, K. S.; Rao, J. M. Chem. Lett. 2009, 38, 564–565; (h) Ollevier, T.; Li, Z. Adv. Synth. Catal. 2009, 351, 3251–3259.