


# RSC Advances



This article can be cited before page numbers have been issued, to do this please use: S. Samanta and P. Biswas, *RSC Adv.*, 2015, DOI: 10.1039/C5RA18151H.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

## COMMUNICATION

# Metal Free Visible Light Driven Oxidation of Alcohols to Carbonyl Derivatives Using 3, 6-Di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz) as Catalyst<sup>†</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Suvendu Samanta and Papu Biswas\*

www.rsc.org/

**3, 6-di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz) catalyzed oxidation of alcohols to corresponding carbonyl compounds under visible light irradiation is described. This reaction occurs smoothly at room temperature and shows good tolerance of functional groups. It provides an alternative approach for the synthesis of alkyl and aryl aldehydes and ketones.**


## Introduction

Recently, organic reactions driven by visible light at ambient temperature in presence of dyes has attracted much interest of synthetic chemists. Usually, polypyridine complexes of ruthenium and iridium have been utilized as dyes in most of these reactions.<sup>1</sup> More recently, few organic dyes have also been reported as visible light photoredox catalysts.<sup>2</sup>

Among organic transformations, the selective oxidation of alcohols to the corresponding carbonyl groups is one of the most important and widely utilized in the manufacture of fine chemicals and intermediates.<sup>3</sup> Consequently, several effective reagents and methods have been developed, including stoichiometric oxidizing agents such as hypochlorite,<sup>4</sup> chromium based reagents,<sup>5</sup> active manganese oxides and permanganate,<sup>6,7</sup> Dess-Martin periodinane,<sup>8</sup> *o*-iodoxybenzoic acid (IBX),<sup>9</sup> ruthenium based reagents,<sup>10</sup> osmium(VIII) oxide<sup>11</sup> and activated DMSO in Swern oxidation.<sup>12</sup> However, most of these reagents are toxic, hazardous and sometimes generates harmful wastes. Numerous homogenous as well as heterogeneous transition metal<sup>13-19</sup> and TEMPO<sup>20</sup> based catalysts have also been reported to be active for aerobic oxidation of alcohols, as they are particularly advantageous from environmental and economic perspective. However, they require relatively expensive metal catalysts, a large amount of additives, and higher temperature to obtain satisfactory results. Few reports also appeared on photocatalytic conversion of alcohols to aldehydes and ketones using visible


light.<sup>21,22</sup> Due to the importance of this reaction, the development of efficient, mild, and metal free safe oxidation processes are still highly desirable.

Recently, we have reported the visible light absorption and reversible one-electron reduction properties of the 3, 6-di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz) molecule (Figure 1)<sup>2b</sup> and its application to the photocatalytic synthesis of 2-substituted benzimidazoles and benzothiazoles from different aldehydes. *s*-Tetrazines (Tz) are known to have a strong electron-deficient character and accept electrons and protons



**Fig. 1** 3, 6-di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz).

simultaneously to form dihydrotetrazenes (H<sub>2</sub>Tz) via anion radical (Scheme 1).<sup>22</sup> This reducing nature of the *s*-tetrazines is even more prominent for the first excited state, which therefore has a relatively strong oxidizing power.<sup>22</sup> Consequently, *s*-tetrazines interact with various electron donor substrates at an excited state.



**Scheme 1** Reduction of *s*-Tetrazine (Tz) to Dihydrotetrazine (H<sub>2</sub>Tz) via Anion Radical


In our efforts to further explore the photocatalytic activity of pytz and expand its synthetic utility, we report here metal-free efficient oxidation of alcohols to the corresponding carbonyl compounds under visible light irradiation using pytz as catalyst (Scheme 2). During oxidation of alcohols, pytz is converted to H<sub>2</sub>pytz. In order to reoxidize the H<sub>2</sub>pytz, an efficient co-oxidant is required. We have utilized O<sub>2</sub> or *tert*-

Department of Chemistry, Indian Institute of Engineering Science and Technology, Shibpur, Howrah, India, 711 103.

E-mail: papubiswas\_besus@yahoo.com.

<sup>†</sup>Electronic Supplementary Information (ESI) available: Experimental procedures, <sup>1</sup>H NMR data and spectra. See DOI: 10.1039/x0xx00000x

butylnitrite/AcOH mixture as sacrificial co-oxidant to establish two alternative methods of oxidation.




**Scheme 2** Oxidation of Alcohols to Carbonyls by Pytz Under Visible Light Irradiation

## Results and discussion

Our preliminary studies were carried out with benzyl alcohol **1a** as a model substrate in presence of pytz in CH<sub>3</sub>CN. In a typical experiment, benzyl alcohol **1a** in CH<sub>3</sub>CN was irradiated under visible light in presence of 5 mol% of pytz catalyst at am-

**Table 1** Optimization of the Reaction Conditions<sup>a</sup>




entry	catalyst	solvent	time (h)	yield (%) <sup>b</sup>
1 <sup>c</sup>	pytz	CH <sub>3</sub> CN	12	traces
2 <sup>d</sup>	pytz	CH <sub>3</sub> CN	12	13
3	pytz	CH <sub>3</sub> CN	10	85
4	pytz	H <sub>2</sub> O	12	26
5	pytz	dioxane	12	30
6	pytz	THF	12	42
7	pytz	CH <sub>3</sub> Cl	12	55
8	pytz	CH <sub>2</sub> Cl <sub>2</sub>	12	48
9	pytz	DMF	12	38
10	phtz	CH <sub>3</sub> CN	10	35
11 <sup>e</sup>	pytz	CH <sub>3</sub> CN	12	traces
12 <sup>f</sup>	none	CH <sub>3</sub> CN	12	none
13 <sup>g</sup>	pytz	CH <sub>3</sub> CN	4	92
14 <sup>g</sup>	none	CH <sub>3</sub> CN	5	none

<sup>a</sup>Reaction Condition (until otherwise specified): 2 mmol benzyl alcohol, solvent (5 mL solvents used for all cases), pytz (5 mol%), O<sub>2</sub>-atmosphere, room temperature, visible light. <sup>b</sup>Isolated yield.

<sup>c</sup>Reaction done in inert atmosphere. <sup>d</sup>Reaction done in presence of air. <sup>e</sup>Reaction carried out in absence of visible-light (dark). <sup>f</sup>Reaction carried out without any catalyst but in presence of visible-light. <sup>g</sup>Reaction carried out in presence of *tert*-butylnitrite (1.1 equiv), AcOH (1.1 equiv).

bient temperature. Use of 5 mol% of pytz catalyst was found to be optimum and any other substrate to catalyst ratio furnished inferior yields. A Xenon lamp with a power of 300 W equipped with a cut-off filter ( $\lambda > 420$  nm) was used as a visible light source (for schematic diagram of reaction set-up, see Scheme S1<sup>†</sup>). Under nitrogen, only trace of the desired product **2a** was obtained (Table 1, entry 1). Changing from oxygen-free environment to aerobic condition resulted in a substantial improvement in conversion (entry 2). Then we have considered the use of pure molecular oxygen. Accordingly, the conversion increased dramatically when reaction was carried out under O<sub>2</sub>-atmosphere (entry 3). No noticeable over oxidation to carboxylic acid was isolated or identified. Performing the reaction in other solvents including water, dioxane, THF, chloroform, or dichloromethane gave much lower yields of **2a** (entries 4–9). We have also investigated the catalytic activity of the 3,6-diphenyl-1,2,4,5-tetrazine (phtz), under identical conditions (entry 10). Phtz shows much inferior activity. To establish that visible light is essential for the reaction to proceed, a control reaction was carried out in the dark in presence of pytz. No catalytic activity was observed in the absence of visible light, which clearly demonstrates the role of visible light in this reaction (entry 11). Similarly, a control reaction in the absence of pytz did not proceed under identical conditions (entry 12). Encouraged by these results, we further carried out the reaction in presence of *tert*-butylnitrite/acetic acid (AcOH) mixture as oxidant. Oxidation of H<sub>2</sub>Tz by NaNO<sub>2</sub>/acid mixture is long known.<sup>22</sup> In the presence of air, these generate nitrogen dioxide, a sufficiently strong oxidant to efficiently recycle the H<sub>2</sub>Tz. The effect of reaction time and solvent on the reaction was again evaluated thoroughly. Interestingly, the reaction proceeds within much lower time in CH<sub>3</sub>CN (entry 13). No product was obtained when a control reaction was done in presence of *tert*-butylnitrite/AcOH mixture but without catalyst in presence of visible light (entry 14). So, the optimum reactions were found to involve **1a**, pytz (5 mol%), and O<sub>2</sub> in CH<sub>3</sub>CN at ambient temperature under visible light irradiation (Method A, Scheme 3), and **1a**, pytz (5 mol%), *tert*-butylnitrite (1.1 equiv), and AcOH (1.1 equiv) in CH<sub>3</sub>CN at ambient temperature under visible light irradiation (Method B, Scheme S2<sup>†</sup>).

Having the optimum conditions in hands, the scope for the oxidation was subsequently investigated with various primary alcohols; reactions under oxygen environment (method A) are shown for comparison. As shown in Scheme 3, high yields were obtained for a wide range of alcohols. Under these reaction conditions, various benzylic alcohols with different substituents on the phenyl ring containing both electron-withdrawing and electron-donating can be efficiently oxidized to the corresponding aldehydes in high yields (Scheme 3, **2a–e**). Anthracen-9-ylmethanol (**1f**) and heterocyclic alcohols (**1g** and **1h**) were smoothly oxidized to give the aldehydes in good yields (Scheme 3, **2f–h**). In addition, primary allylic alcohol (**1i**) and  $\beta$ -O-4 lignin model, 4-(hydroxymethyl)-2-methoxyphenol (**1j**), were also effective substrates, affording the corresponding aldehydes in good yields (Scheme 3, **2i** and **2j**). The present catalytic system is not efficient for aliphatic prima-




**Scheme 3** Oxidation of Primary Alcohols. Method A: 2 mmol alcohol, acetonitrile (5 mL), pytz (5 mol%), O<sub>2</sub>-atmosphere, room temperature, visible light; Method B: 2 mmol alcohol, acetonitrile (5 mL), pytz (5 mol%), room temperature, visible light, *t*-BuONO<sub>2</sub> (2.2 mmol), AcOH (2.2 mmol). Isolated yields are shown.

ry alcohols and shows relatively poor yields (< 40%).

Secondary alcohols were also found to be effective substrates and generally more reactive than primary alcohols. Various secondary aryl alcohols were effectively oxidized to ketones (Scheme 4, entries **4a-g**). A number of functionalities, such as bromine, methyl, nitro, and diphenylmethanol are compatible with the reaction conditions and furnished the corresponding ketones in excellent yields (Scheme 4, **4b-e**). 1-(Anthracen-9-yl)ethan-1-ol (**3f**) and 9,10-dihydroanthracen-9-ol (**3g**) underwent smooth oxidation to provide the ketones in good yield (Scheme 4, **4f** and **4g**). Furthermore, heterocyclic secondary alcohol, 1-(thiophen-3-yl)ethan-1-ol (**3h**) and alkyl secondary alcohol, cyclohexanol (**3i**), were also effectively oxidised to corresponding ketones (Scheme 4, **4h** and **4i**) with 71 and 79% yields, respectively by method B.

The visible light driven mild pytz-catalysed oxidation of alcohols to carbonyls also proved easy to scale up, as demonstrated by a gram scale experiment in which 20 mmol of benzyl alcohol in 50 mL of acetonitrile (**1a**) was irradiated under visible light in presence of *t*-BuONO/AcOH. The reaction resulted in 2.06 g (19.4 mmol, 97% yield) of analytically pure benzaldehyde in 7 h.

To understand this visible light driven oxidation of alcohols to carbonyls catalyzed by pytz, a detailed investigation on the mechanism was undertaken. Due to their electron deficient character, tetrazine molecules can be reversibly reduced in organic solvents. Almost all Tetrazine molecules accept one




**Scheme 4** Oxidation of Secondary Alcohols. Method A: 2 mmol alcohol, acetonitrile (5 mL), pytz (5 mol%), O<sub>2</sub>-atmosphere, room temperature, visible light; Method B: 2 mmol alcohol, acetonitrile (5 mL), pytz (5 mol%), room temperature, visible light, *t*-BuONO (2.2 mmol), AcOH (2.2 mmol). Isolated yields are shown.

electron to produce an anion radical which is very stable in absence of any proton donor.<sup>22c,23</sup> Most of the tetrazine derivatives can accept a second electron though this process is not electrochemically reversible in standard conditions. The unstable dianion formed after transfer of second electron reacts with proton donor to give dihydrotetrazine<sup>22c</sup> as shown in Scheme 1. Fukuzumi et al. reported the reduction of 3,6-diphenyl-*s*-tetrazine (PhTz) by 10-methyl-9,10-dihydroacridine (AcrH<sub>2</sub>) promoted by scandium ion.<sup>24</sup> In presence of Sc<sup>3+</sup>, hydride transfer from AcrH<sub>2</sub> or 10,10'-dimethyl-9,9'-biacridine [(AcrH<sub>2</sub>)<sub>2</sub>] to PhTz occurs efficiently at room temperature to yield 10-methylacridinium ion (AcrH<sup>+</sup>). So, tetrazines can be utilized as two electron oxidising agent under suitable conditions.

In order to elucidate the mechanism of the present direct alcohol oxidation, the optical and electrochemical properties of the pytz molecule were studied by UV-vis absorption spectroscopy and cyclic voltammetry in acetonitrile. Pytz exhibits maximum absorption peaks at 532 nm ( $\epsilon = 190 \text{ M}^{-1} \text{ cm}^{-1}$ , Fig. 2a) and two quasi-reversible reduction peak at +0.31 and -0.91 V in acetonitrile (Fig. 2b and c). Pytz molecule successively accepts two electrons quasi-reversibly in acetonitrile and can be conveniently converted into H<sub>2</sub>pytz in presence of proton donor. Occurrence of first reduction potential at +ve potential indicates that pytz molecule is highly susceptible towards reduction and forms stable anion radical.

While a precise understanding of the reaction mechanism awaits further study, a plausible catalytic pathway is proposed





**Fig. 2** (a) UV-vis absorption spectra and cyclic voltammogram (CV) of 3,6-di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz) in acetonitrile in the range (b) +1.0 to −0.5 V and (c) −0.5 to −1.0 V. Scan rate for cyclic voltametric measurements were 50 mVs<sup>−1</sup>.

in Scheme 2. According to the previous studies,<sup>1a,b,21c,d</sup> we proposed that the reaction may proceed through the formation of radical cation, ROH<sup>•+</sup>. In presence of visible light, pytz is converted to pytz<sup>•+</sup>, which is transformed into pytz radical anion through single electron transfer (SET) from alcohol to form radical cation, ROH<sup>•+</sup>. The poor conversion of aliphatic primary alcohols supports the formation of radical cation as aliphatic primary radical cations are not stable. Then, hydrogen atom abstraction by stable pytz radical anion from the radical cation ROH<sup>•+</sup> followed by deprotonation gives the carbonyl compound and pytz is subsequently converted to H<sub>2</sub>pytz. Oxidation of H<sub>2</sub>pytz to regenerate catalyst, pytz, can easily be carried out in presence of molecular oxygen or *tert*-butylnitrite/AcOH mixture. The overall reaction is as follows (eqn 1):



To prove the radical pathway, this reaction was also done in the presence of a radical scavenger, TEMPO (TEMPO = 2,2,6,6-tetramethylpiperidinoxyl). Addition of TEMPO in a 1:1 molar proportion (with respect to benzyl alcohol) to the reaction mixture, only trace yield can be identified by GC. To check the possibility of pytz as the sensitizer of O<sub>2</sub> and the activated molecular O<sub>2</sub> is the real oxidant, we performed the oxidation reaction of benzyl alcohol in the presence of one equivalent of pytz under inert atmosphere and the reaction yield ~ 80% benzaldehyde within 8 h. These results did not support the involvement of activated molecular O<sub>2</sub> as the real oxidant. Moreover, molecular oxygen accepts one electron

from activated photocatalysts to produce superoxide anion, O<sub>2</sub><sup>•−</sup>. As pytz molecule itself accepts electron in its excited state to act as an oxidising agent, formation of superoxide is not feasible under present reaction protocol. Hence, the involvement of activated molecular O<sub>2</sub> as the real oxidant is ruled out.

## Conclusion

In summary, we have developed a highly efficient metal free oxidation of alcohols under visible light irradiation using pytz as catalyst and molecular oxygen or *tert*-butylnitrite as co-oxidant. Various primary and secondary alcohols can be efficiently oxidized under mild conditions to the corresponding aldehydes and ketones in high yields. The present procedure shows the following favourable features: (1) the reaction does not require any metal based oxidant; (2) the reaction proceeds under neutral and mild conditions (neither acid nor base required) and thus making it suitable with acid- or base sensitive substrates; (3) the present method is effective for a wide range of substrates bearing various functional groups; (4) the reaction can also be carried out in presence of molecular oxygen further making it environmentally benign procedure; (5) our developed catalytic system should be suitable for large-scale synthesis of carbonyls from alcohols. Further development of other reaction processes with pytz and related tetrazines is currently underway.

## Acknowledgements

S. S. is indebted to CSIR, India for his Senior Research Fellowship [08/003(0083)/2011-EMR-1]. P.B. acknowledges CSIR-India for the Project (Sanction letter no. 01(2459)/11/EMR-II dated 16/05/2011). The authors also acknowledge the Sophisticated Analytical Instruments Facility at North Eastern Hill University (SAIF-NEHU) for <sup>1</sup>H NMR analysis.

## References


- For leading references on visible-light photoredox catalysis, see: (a) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (b) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102; (c) J. W. Tucker and C. R. J. Stephenson, *J. Org. Chem.*, 2012, **77**, 1617; (d) T. P. Yoon, M. A. Ischay and J. Du, *Nature Chem.*, 2010, **2**, 527 and references therein.
- (a) J. H. Park, K. C. Ko, E. Kim, N. Park, J. H. Ko, D. H. Ryu, T. K. Ahn, J. Y. Lee and S. U. Son, *Org. Lett.*, 2012, **14**, 5502; (b) S. Samanta, S. Das and P. Biswas, *J. Org. Chem.*, 2013, **78**, 11184; (c) X. Gu, X. Li, Y. Chai, Q. Yang, P. Li and Y. Yao, *Green Chem.*, 2013, **15**, 357; (d) J. Luo, X. Zhang and J. Zhang, *ACS Catal.*, 2015, **5**, 2250; (e) X. Wu, C. Meng, X. Yuan, X. Jia, X. Qian and J. Ye, *Chem. Commun.*, 2015, **51**, 11864; (f) D. A. Nicewicz and T. M. Nguyen, *ACS Catal.*, 2014, **4**, 355–360; (g) D. Ravelli and M. Fagnoni, *ChemCatChem*, 2012, **4**, 169; (h) C. Vila, J. Lau and M. Rueping, *Beilstein J. Org. Chem.*, 2014, **10**, 1233; (i) J. Li, H. Wang, L. Liu and J. Sun, *RSC Adv.*, 2014, **4**, 49974; (j) S. Fukuzumi and K. Ohkubo, *Org. Biomol. Chem.*, 2014, **12**, 6059.
- (a) M. Hudlucky, *Oxidations in Organic Chemistry*; ACS Monograph Series; American Chemical Society: Washington, DC, 1990; (b) T. Mallat and A. Baiker, *Chem. Rev.*, 2004, **104**,

- 3037; (c) R. C. Larock, *Comprehensive Organic Transformations*, 2nd ed.; Wiley-VCH: New York, 1999; (c) I. W. C. E. Arends and R. A. Sheldon, In *Modern Oxidation Methods*, 2nd ed. J.-E. Bäckvall, Ed. Wiley-VCH: Weinheim, 2010.
- 4 R. V. Stevens, K. T. Chapman and H. N. Weller, *J. Org. Chem.*, 1980, **45**, 2030.
  - 5 (a) E. J. Corey and J. W. Suggs, *Tetrahedron Lett.*, 1975, **16**, 2647; (b) F. A. Luzzio and F. S. Jr. Guziec, *Org. Prep. Proced. Int.*, 1988, **20**, 533.
  - 6 (a) Harfenist, M.; Bavley, A.; Lazier, W. A. *J. Org. Chem.* **1954**, **19**, 1608. (b) R. J. Highet and W. C. Wildman, *J. Am. Chem. Soc.*, 1955, **77**, 4399; (c) R. J. K. Taylor, M. Reid, J. Foot and S. A. Raw, *Acc. Chem. Res.*, 2005, **38**, 851.
  - 7 F. M. Menger and C. Lee, *Tetrahedron Lett.*, 1981, **22**, 1655.
  - 8 (a) D. B. Dess and J. C. Martin, *J. Org. Chem.*, 1983, **48**, 4155; (b) D. B. Dess and J. C. Martin, *J. Am. Chem. Soc.*, 1991, **113**, 7277; (c) R. E. Ireland and L. Liu, *J. Org. Chem.*, 1993, **58**, 2899; (d) S. D. Meyer and S. L. Schreiber, *J. Org. Chem.*, 1994, **59**, 7549; (e) K. C. Nicolaou, Y.-L. Zhong and P. S. Baran, *J. Am. Chem. Soc.*, 2000, **122**, 7596; (f) M. Uyanik and K. Ishihara, *Chem. Commun.*, 2009, 2086.
  - 9 (a) M. Mulbaier and A. Giannis, *Angew. Chem. Int. Ed.*, 2001, **40**, 4393; (b) G. Sorg, A. Mengel, G. Jung and J. Rademann, *Angew. Chem., Int. Ed.*, 2001, **40**, 4395; (c) K. C. Nicolaou, T. Montagnon, P. S. Baran and Y.-L. Zhong, *J. Am. Chem. Soc.*, 2002, **124**, 2245; (d) J. D. More and N. S. Finney, *Org. Lett.*, 2002, **4**, 3001; (e) A. P. Thottumkara, M. S. Bowsheer and T. K. Vinod, *Org. Lett.*, 2005, **7**, 2933.
  - 10 (a) L. M. Berkowitz and P. N. Rylander, *J. Am. Chem. Soc.* 1958, **80**, 6682; (b) H. Tomioka, K. Takai, K. Oshima and H. Nozaki, *Tetrahedron Lett.*, 1981, **22**, 1605.
  - 11 A. M. Maione and A. Romeo, *Synthesis*, 1984, 955.
  - 12 (a) K. Omura and D. Swern, *Tetrahedron*, 1978, **34**, 1651; (b) T. T. Tidwell, *Synthesis*, 1990, 857; (c) L. De Luca, G. Giacomelli and A. Porcheddu, *J. Org. Chem.*, 2001, **66**, 7907.
  - 13 For leading references on vanadium-catalyzed oxidation of alcohols, see: (a) C. Li, P. Zheng, J. Li, H. Zhang, Y. Cui, Q. Shao, X. Ji, J. Zhang, P. Zhao and Y. Xu, *Angew. Chem. Int. Ed.*, 2003, **42**, 5063; (b) V. D. Pawar, S. Bettigeri, S.-S. Weng, J.-Q. Kao and C.-T. Chen, *J. Am. Chem. Soc.*, 2006, **128**, 6308. (c) K. Alagiri and K. R. Prabhu, *Tetrahedron*, 2011, **67**, 8544.
  - 14 For leading references on manganese-catalyzed oxidation of alcohols, see: (a) Y.-C. Son, V. D. Makwana, A. R. Howell and S. L. Suib, *Angew. Chem. Int. Ed.*, 2001, **40**, 4280; (b) H.-K. Kwong, P.-K. Lo, K.-C. Lau and T.-C. Lau, *Chem. Commun.*, 2011, 4273.
  - 15 For leading references on iron-catalyzed oxidation of alcohols, see: (a) F. Shi, M. K. Tse, M.-M. Pohl, A. Brückner, S. Zhang and M. Beller, *Angew. Chem., Int. Ed.*, 2007, **46**, 8866; (b) T. Kunisu, T. Oguma and T. Katsuki, *J. Am. Chem. Soc.*, 2011, **133**, 12937.
  - 16 For leading references on cobalt-catalyzed oxidation of alcohols, see: (a) B. S. Tovrog, S. E. Diamond, F. Mares and A. Szalkiewicz, *J. Am. Chem. Soc.*, 1981, **103**, 3522; (b) T. Iwahama, Y. Yoshino, T. Keitoku, S. Sakaguchi and Y. Ishii, *J. Org. Chem.*, 2000, **65**, 6502; (c) N. Gunasekaran, P. Jerome, S. W. Ng, E. R. T. Tiekink and R. Karvembu, *J. Mol. Catal. A: Chem.*, 2012, **156**, 353.
  - 17 For leading references on copper-catalyzed oxidation of alcohols, see: (a) M. F. Semmelhack, C. R. Schmid, D. A. Cortés and C. S. Chou, *J. Am. Chem. Soc.*, 1984, **106**, 3374; (b) E. Markó, P. R. Giles, M. Tsukazaki, S. M. Brown and C. J. Urch, *Science*, 1996, **274**, 2044; (c) I. A. Ansari and R. Gree, *Org. Lett.*, 2002, **4**, 1507; (d) P. Gamez, I. W. C. E. Arends, J. Reedijk and R. A. Sheldon, *Chem. Commun.*, 2003, 2414; (d) I. E. Markó, A. Gautier, R. Dumeunier, K. Doda, F. Philippart, S. M. Brown and C. J. Urch, *Angew. Chem. Int. Ed.*, 2004, **43**, 1588; (e) N. Jiang and A. J. Ragauskas, *J. Org. Chem.*, 2006, **71**, 7087; (f) J. M. Hoover and S. S. Stahl, *J. Am. Chem. Soc.*, 2011, **133**, 16901. (g) Y. Zhu, B. Zhao and Y. Shi, *Org. Lett.*, 2013, **15**, 992.
  - 18 For leading references on palladium and gold-catalyzed oxidation of alcohols, see: (b) G.-J. ten Brink, I. W. C. E. Arends and R. A. Sheldon, *Science*, 2000, **287**, 1636; (a) D. R. Jensen, J. S. Pugsley and M. S. Sigman, *J. Am. Chem. Soc.*, 2001, **123**, 7475; (b) E. M. Ferreira and B. M. Stoltz, *J. Am. Chem. Soc.*, 2001, **123**, 7725; (c) S. S. Stahl, J. L. Thorman, R. C. Nelson and M. A. Kozee, *J. Am. Chem. Soc.*, 2001, **123**, 7188; (d) H. Tsunoyama, H. Sakurai, Y. Negishi and T. Tsukuda, *J. Am. Chem. Soc.*, 2005, **127**, 9374; (e) B. Guan, D. Xing, G. Cai, X. Wan, N. Yu, Z. Fang, L. Yang and Z. Shi, *J. Am. Chem. Soc.*, 2005, **127**, 18004; (f) J. Ni, W.-J. Yu, L. He, H. Sun, Y. Cao, H.-Y. He and K.-N. Fan, *Green Chem.*, 2009, **11**, 756; (g) R. Dun, X. Wang, M. Tan, Z. Huang, X. Huang, W. Ding and X. Lu, *ACS Catal.*, 2013, **3**, 3063.
  - 19 For leading references on ruthenium-catalyzed oxidation of alcohols, see: (a) W. P. Griffith, S. V. Ley, G. P. Whitcombe and A. D. White, *J. Chem. Soc., Chem. Commun.*, 1987, 1625; (b) P. E. Morris and D. E. Kiely, *J. Org. Chem.*, 1987, **52**, 1149; (d) W.-H. Fung, W.-Y. Yu and C.-M. Che, *J. Org. Chem.*, 1998, **63**, 2873; (e) G. Csajnyik, A. H. Éll, L. Fadini, B. Pugin and J.-E. Bäckvall, *J. Org. Chem.*, 2002, **67**, 1657; (f) K. Yamaguchi and N. Mizuno, *Angew. Chem., Int. Ed.*, 2002, **41**, 4538; (g) L. Gonsalvi, I. W. C. E. Arends and R. A. Sheldon, *Org. Lett.*, 2002, **4**, 1659; (h) B.-Z. Zhan, M. A. White, T.-K. Sham, J. A. Pincock, R. J. Doucet, K. V. R. Rao, K. N. Robertson and T. S. Cameron, *J. Am. Chem. Soc.*, 2003, **125**, 2195.
  - 20 For leading references on TEMPO-catalyzed oxidation of alcohols, see: (b) J. Einhorn, C. Einhorn, F. Ratajczak and J.-L. Pierre, *J. Org. Chem.*, 1996, **61**, 7452; (c) S. D. Rychnovsky and R. Vaidyanathan, *J. Org. Chem.*, 1999, **64**, 310; (d) C. Bolm, A. S. Magnus and J. P. Hildebrand, *Org. Lett.*, 2000, **2**, 1173; (e) R. Liu, X. Liang, C. Dong and X. Hu, *J. Am. Chem. Soc.*, 2004, **126**, 4112; (f) M. Shibuya, Y. Osada, Y. Sasano, M. Tomizawa and Y. Iwabuchi, *J. Am. Chem. Soc.*, 2011, **133**, 6497.
  - 21 (a) M. Rueping, C. Vila, A. Szadkowska, R. M. Koenigs and J. Fronert, *ACS Catal.*, 2012, **2**, 2810; (b) F. Su, S. C. Mathew, G. Lipner, X. Fu, M. Antonietti, S. Blechert and X. Wang, *J. Am. Chem. Soc.*, 2010, **132**, 16299; (c) M. Zhang, C. C. Chen, W. H. Ma and J. C. Zhao, *Angew. Chem., Int. Ed.*, 2008, **47**, 9730; (d) H. Cano-Yelon and A. Deronzier, *Tetrahedron Lett.*, 1984, **25**, 5517.
  - 22 (a) W. Kaim, *Coord. Chem. Rev.*, 2002, **230**, 127; (b) N. Saracoglu, *Tetrahedron*, 2007, **63**, 4199; (c) G. Clavier and P. Audebert, *Chem. Rev.*, 2010, **110**, 3299 and ref. therein.
  - 23 Y.-H. Gong, P. Audebert, J. Tang, F. Miomandre, G. Clavier, S. Badré, R. Méallet-Renault and J. Marrot, *J. Electro. Anal. Chem.*, 2006, **592**, 147.
  - 24 S. Fukuzumi, J. Yuasa and T. Suenobu, *J. Am. Chem. Soc.*, 2002, **124**, 12566.

### Graphical Abstract

#### Metal Free Visible Light Driven Oxidation of Alcohols to Carbonyls Using 3, 6-Di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz) as Catalyst<sup>†</sup>

Suvendu Samanta and Papu Biswas



A metal-free photocatalytic system for the transformation of alcohols to corresponding carbonyls in high yields with visible-light irradiation has been achieved. Di(pyridin-2-yl)-1,2,4,5-tetrazine (pytz) was found to be an efficient catalyst for this oxidation under photo irradiation conditions.