# Bioconjugate Chemistry

# Trihydroxamate Siderophore–Fluoroquinolone Conjugates Are Selective Sideromycin Antibiotics that Target *Staphylococcus aureus*

Timothy A. Wencewicz,<sup>†</sup> Timothy E. Long,<sup>†,#</sup> Ute Möllmann,<sup>‡</sup> and Marvin J. Miller\*<sup>,†</sup>

<sup>†</sup>Department of Chemistry and Biochemistry, 251 Nieuwland Science Hall, University of Notre Dame, Notre Dame, Indiana 46556, United States

<sup>‡</sup>Leibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute, Beutenbergstrasse 11a, 07745 Jena, Germany

**Supporting Information** 

**ABSTRACT:** Siderophores are multidentate iron(III) chelators used by bacteria for iron assimilation. Sideromycins, also called siderophore—antibiotic conjugates, are a unique subset of siderophores that enter bacterial cells via siderophore uptake pathways and deliver the toxic antibiotic in a "Trojan horse" fashion. Sideromycins represent a novel antibiotic delivery technology with untapped potential for developing sophisticated microbe-selective antibacterial agents that limit



the emergence of bacterial resistance. The chemical synthesis of a series of mono-, bis-, and trihydroxamate sideromycins are described here along with their biological evaluation in antibacterial susceptibility assays. The linear hydroxamate sideromycins used for the sideromycins in this study were derived from the ferrioxamine family and inspired by the naturally occurring salmycin sideromycins. The antibacterial agents used were a  $\beta$ -lactam carbacepholosporin, Lorabid, and a fluoroquinolone, ciprofloxacin, chosen for the different locations of their biological targets, the periplasm (extracellular) and the cytoplasm (intracellular). The linear hydroxamate-based sideromycins were selectively toxic toward Gram-positive bacteria, especially *Staphylococcus aureus* SG511 (MIC = 1.0  $\mu$ M for the trihydroxamate–fluoroquinolone sideromycin). Siderophore–sideromycin competition assays demonstrated that only the fluoroquinolone sideromycins required membrane transport to reach their cytoplasmic biological target and that a trihydroxamate siderophore backbone was required for protein-mediated active transport of the sideromycins into *S. aureus* cells via siderophore uptake pathways. This work represents a comprehensive study of linear hydroxamate sideromycins and teaches how to build effective hydroxamate-based sideromycins as Gram-positive selective antibiotic agents.

# INTRODUCTION

Bacterial resistance to antibiotics is developing at an alarming rate.<sup>1</sup> As the current pipeline of useful antibacterial agents diminishes, the crisis worsens and there is now a desperate need for new methods of limiting the emergence of resistance to antibiotics. One promising approach involves the development of more sophisticated, pathogen-targeted antibiotic delivery technologies that bypass or limit exposure of the antibacterial agent to known resistance mechanisms. This approach is attractive because it does not require the discovery of new antibacterial scaffolds or validation of new biological targets, which has proven to be an extremely difficult and timeconsuming task.<sup>2,3</sup> New antibiotic delivery methodologies present the opportunity to recycle old antibiotics rendered useless by resistance that have already been structurally optimized to interact with their biological target, revisit potent antibiotics abandoned due to toxicity associated with the existence of human orthologs to the biological target,<sup>4</sup> bring new life to antibiotic scaffolds that failed due to membrane permeability problems,<sup>5,6</sup> and extend the useful clinical lifetime of new antibiotics coming to market by better managing resistance.<sup>7</sup> Considering the serious financial investment associated with bringing a new drug to market and the limited market lifetime of antibiotics, investing in drug delivery technology is a prudent choice for antibiotic discovery programs.<sup>8</sup>

One of the biggest challenges for targeted antibiotic delivery is finding useful biological pathways in bacteria to exploit for membrane transport.<sup>9</sup> The ideal membrane transport pathway for antibiotic delivery should be specific to bacterial cells (to eliminate toxicity toward eukaryotic cells), essential for virulence (to eliminate resistance development via deletion of uptake pathway), and general enough to accept unnatural substrates (to ensure successful uptake of the delivery vector derivatized to carry the antibiotic). Bacterial iron-acquisition pathways have been identified as suitable pathways for developing such antibiotic delivery systems.<sup>10–12</sup>

The most common pathway for bacterial iron acquisition involves the biosynthesis and excretion of low-molecular-weight multidentate iron(III)-chelators, known as siderophores, used

```
Received:November 17, 2012Revised:January 10, 2013Published:January 28, 2013
```



**Figure 1.** Generic schematic of siderophore-mediated iron uptake and genetic regulation in Gram-negative and Gram-positive bacteria. Iron metabolism in bacteria is under genetic control by the ferric uptake regulator (Fur) transcription factor protein. During times of iron abundance, the Fur protein complexes Fe(II) and adopts a conformation that binds to a region of DNA known as a "Fur box" which represses the expression of siderophore-related proteins, including biosynthetic enzymes. During times of iron deficiency, the apo-Fur protein dissociates from the "Fur box" allowing expression of siderophore-related genes. The siderophore system is now fully functional and begins with siderophore biosynthesis and efflux to the extracellular environment. For Gram-negative bacteria, after the siderophore chelates Fe(III) to form a soluble complex it is transported into the periplasm by a high-affinity outer membrane transport protein (OMT; formerly called OMR). This transport is driven by the proton motive force with energy transfer mediated by the membrane-spanning TonB-ExbB/D protein complex. The Fe(III)-loaded siderophore is then bound by a siderophore periplasmic binding protein (SPBP) and trafficked to an ATP-binding cassette transporter (ABC) that moves the siderophore-Fe(III) complex into the cytoplasm. The iron nutrient is then removed from the siderophore, typically via reduction of Fe(III) to Fe(II) by an iron reductase enzyme (Red), and is distributed to parts of the cell in need or stored in bacterioferritin. In most cases, the siderophore is then recycled for further use. Gram-positive bacteria lack the outer membrane found in Gram-negative bacteria and therefore also lack the OMTs and TonB complex. Instead, extracellular siderophore-Fe(III) complexes are recognized and transported by membrane-anchored SPBPs. The remaining steps of the siderophore pathway in Gram-positive bacteria are analogous to those described for Gram-negative bacteria. This figure was inspired by a variety of siderophore t

for extracellular solubilization of otherwise insoluble iron(III) minerals followed by protein-mediated active transport of the metal complexes into bacterial cells (Figure 1).<sup>13–15</sup> To ensure a competitive growth advantage, bacteria commonly possess the ability to obtain iron(III) using siderophores produced by competing organisms.<sup>16</sup> Along this evolutionary path, some bacteria have learned to exploit this "iron thievery" for antibiotic delivery by covalently attaching a toxic antibacterial agent to the siderophore. This unique subset of siderophores, known as sideromycins (Figure 2a), enter competing bacterial cells via siderophore uptake pathways (Figure 2b) and deliver the toxic agent in a "Trojan horse" fashion.<sup>17</sup>

The albomycins and salmycins (Figure 2a) are two naturally occurring classes of sideromycins. The albomycins are the best-studied of all sideromycins.<sup>17</sup> They were originally reported in 1947 from a variety of *Streptomyces* strains<sup>18</sup> and the structure was correctly elucidated in 1982.<sup>19</sup> The albomycins consist of a ferrichrome-like trihydroxamate siderophore joined through an amide bond to a thioribosyl pyrimidine inhibitor of seryl-*t*-RNA synthetase and display *in vitro* and *in vivo* broad-spectrum antibacterial activity against Gram-positive and Gram-negative

bacteria with extremely potent minimum inhibitory concentrations (MIC) as low as 5 ng/ $\mu$ L.<sup>17</sup> This impressive potency is attributed to active transport into bacteria via the ferrichrome membrane transport proteins FhuA<sup>20</sup> and FhuD.<sup>21</sup> Once internalized, the seryl-*t*-RNA synthetase inhibitor is released by a serine protease, peptidase N, which cleaves the amide bond linkage to the siderophore.<sup>22</sup> The albomycins are highly effective in mouse infection models,<sup>17</sup> and recently, the biosynthetic gene cluster was discovered that makes possible the production of novel analogues and engineering of superproducers.<sup>23</sup>

Far less is known about the salmycins that were isolated from *Streptomyces violaceus* DSM 8286 in 1995 by Vértesy and coworkers.<sup>24</sup> A total synthesis of the salmycins was completed by Miller and co-workers in 2002 which corrected the structure assignment.<sup>25</sup> The salmycins consist of a linear trihydroxamate siderophore from the ferrioxamine family, known as danoxamine, and an aminoglycoside antibiotic joined through a succinoyl linker (Figure 3). They show potent and selective *in vitro* antibacterial activity against Gram-positive bacteria, including highly antibiotic resistant strains, with reported



**Figure 2.** (A) Structures of generic and natural sideromycins (albomycins and salmycins). (B) Generic schematic of sideromycins as "Trojan horse" antibiotic delivery agents. Sideromycins (siderophore–antibiotic conjugates) act as normal siderophores for sideromycin producing microbes, which can self-protect from the antibiotic, by scavenging extracellular Fe(III) via the pathways described in Figure 1. For competing microbes attempting to steal the sideromycin-bound Fe(III) nutrient, active transport of the sideromycin via siderophore uptake pathways exposes the competing microbe to the antibiotic, for which it lacks the inherent resistance found in the producer, and ultimately causes cell death.



Figure 3. Structures of desferrisalmycin B, desferridanoxamine (3a), desferrioxamine B (DFO-B), and synthetic desferridanoxamine-antibiotic conjugates from this study.

MIC values as low as 10 ng/ $\mu$ L.<sup>17</sup> Similar to the albomycins, the incredible potency is attributed to active transport of the salmycins through hydroxamate siderophore membrane transport proteins.<sup>26</sup> Unfortunately, the salmycins only show weak *in vivo* activity in mouse infection models probably due to extracellular hydrolysis of the labile ester linkage.<sup>17</sup> We hypothesized that intracellular hydrolysis of the aminoglycoside

antibiotic was required for activity and that the ferrioxamine siderophore was responsible for the narrow spectrum of antibacterial activity against Gram-positive bacteria.<sup>27</sup>

In this report, we set out to better understand the selective antibiotic activity of the salmycins using a series of synthetic mono-, bis-, and trihydroxamate sideromycins (Figure 3) and exploit this knowledge for building more effective synthetic



Figure 4. Structures of synthetic hydroxamate siderophores (1a-3a) and sideromycins (1b-3b and 1c-3c) used in this work.

Scheme 1. Syntheses of Iron-Free (1a-3a) and Iron-Bound (3a-Fe) Hydroxamate Siderophores<sup>*a*</sup>



<sup>a</sup>Reagents and conditions: (a) 10% Pd–C, H<sub>2</sub> (1 atm), MeOH; 6 h, 88% for 1a; 4.5 h, 90% for 2a; 11 h, 86% for 3a. (b) Fe(acac)<sub>3</sub>, MeOH; 4 h, 99% for 3a-Fe.

sideromycins. Using broad-spectrum antibiotics with different biological targets,  $\beta$ -lactams and fluoroquinolones, we demonstrate that the full trihydroxamate siderophore backbone is required for active membrane transport and that this type of delivery vector is selective for Gram-positive bacteria that possess hydroxamate siderophore transport proteins, as reported for the salmycins and observed here against *S. aureus*. We also establish that sideromycins with intracellular antibiotic targets, which require membrane transport to be reached, should be used to target Gram-positive pathogens.

# RESULTS AND DISCUSSION

Syntheses of Siderophores and Sideromycins. The synthetic sideromycins used in this study were rationally designed to mimic the structure of the naturally occurring salmycins (Figure 3). The linear hydroxamate siderophores (1a–3a) were all derived from the natural siderophore portion of the salmycins containing a succinoyl linker. Sideromycins 1b–3b featured the broad-spectrum  $\beta$ -lactam carbacephalosporin antibiotic Lorabid which has a periplasmic biological target. Sideromycins 1c–3c featured the broad-spectrum fluoroquinolone antibiotic ciprofloxacin, which has a cytoplasmic biological target (Figure 4).<sup>28</sup>

The hydroxamate siderophores and sideromycins were synthesized using a benzyl-protecting group strategy. The bis-O-benzyl (4), tris-O-benzyl (5), and tetra-O-benzyl (6) protected siderophore precursors were synthesized using a literature protocol developed by Roosenberg and Miller.<sup>29</sup> The corresponding deprotected, iron-free siderophores (1a, 2a, and 3a) were prepared and isolated in high yield and purity using Pd-catalyzed hydrogenolysis followed by precipitation from MeOH/Et<sub>2</sub>O. The iron(III)-complexed siderophore, danoxamine (3a-Fe), was isolated in quantitative yield as an orange solid after treatment of desferridanoxamine (3a) with Fe(acac)<sub>3</sub> and precipitation from MeOH/Et<sub>2</sub>O (Scheme 1).<sup>30</sup>

Antibiotic derivatives 8 and 10 were synthesized for use as coupling partners with the O-benzyl-protected siderophores 4, 5, and 6. N-Boc-O-PNB-Lorabid (7; a gift from Eli Lilly and Co.) was treated with anhydrous TFA to give TFA salt 8. Ciprofloxacin (9) was converted to O-benzyl-ciprofloxacin HCl salt (10) by sequential N-Boc protection, carboxylate benzylation, and N-Boc deprotection according to a literature protocol (Scheme 2).<sup>31</sup> The free amines of the protected  $\beta$ lactam and fluoroquinolone antibiotics, 8 and 10, were poised for conjugation to the succinoyl group of benzyl-protected siderophore precursors 4-6 through amide bond formations. Importantly,  $\beta$ -lactam<sup>32</sup> and fluoroquinolone<sup>33</sup> antibiotics are known to tolerate bulky substituents at these points of derivatization and representative siderophore-conjugates of these classes of antibiotics have been shown to retain affinity for their distinct biological targets.<sup>34,35</sup>

The fully protected sideromycins (11-16) were assembled using EDC-mediated amide couplings between the free succinoyl carboxylate of the protected siderophores (4-6)and the antibiotic amine salts (8 and 10) under basic conditions (Scheme 3). The protected Lorabid sideromycins Scheme 2. Synthesis of Protected Antibiotics (8 and 10) Suitable for Siderophore Conjugation<sup>a</sup>



<sup>a</sup>Reagents and conditions: (a) N-Boc-O-PNB-Lorabid (7), TFA,  $CH_2Cl_2$ , 30 min, 99%. (b) ciprofloxacin (9), Boc<sub>2</sub>O, NaHCO<sub>3</sub>, DMF, 2.5 h; then, BnBr, 90 °C, 25.5 h, 98%. (c) conc. HCl,  $CH_2Cl_2$ , 40 min, 92%.

(11, 13, and 15) were universally deprotected using hydrogenolysis conditions optimized to preserve the vinyl chloride functionality of the  $\beta$ -lactam bicycle.<sup>36–39</sup> Purification by preparative HPLC and recrystallization from MeOH/Et<sub>2</sub>O provided analytically pure samples of the  $\beta$ -lactam sideromycins (1b, 2b, and 3b). The O-benzyl protected ciprofloxacin sideromycins (12, 14, and 16) were universally deprotected by Pd-catalyzed hydrogenolysis, and the fluoroquinolone sideromycins (1c, 2c, and 3c) were obtained in pure form after recrystallization from  $MeOH/Et_2O$  with no need for chromatographic purification.

Antibacterial Activity of Siderophores and Sideromycins Against ESKAPEE Pathogens. The siderophores, sideromycins, and control antibiotics (Lorabid and ciprofloxacin) were tested for antibacterial activity against a panel of ESKAPEE bacteria (E. faecium, S. aureus, K. pneumonia, A. baumannii, P. aeruginosa, E. aerogenes, E. coli)<sup>1</sup> by determining the minimum inhibitory concentrations (MIC) using the broth microdilution assay (Table 1).<sup>40</sup> As expected, the siderophores (1a-3a and 3a-Fe) had no antibacterial activity (MIC values >128  $\mu$ M) since these molecules should be growth-promoting factors. The  $\beta$ -lactam-hydroxamate conjugates (1b-3b) had moderate activity against S. aureus (MIC values from 32 to 64  $\mu$ M), but at a reduced potency relative to the parent antibiotic, Lorabid (MIC value of 1  $\mu$ M). Conjugates 1b-3b also had moderate activity against A. baumannii (MIC values from 32 to 128  $\mu$ M), while Lorabid had no measurable activity (MIC value of >128  $\mu$ M). Overall, the  $\beta$ -lactam-hydroxamate conjugates (1b-3b) had a narrowed spectrum of activity and reduced potency (higher MIC values) against the ESKAPEE bacteria relative to Lorabid alone.

The fluoroquinolone-hydroxamate conjugates (1c-3c) also showed a reduced spectrum of activity relative to the broadly active parent antibiotic, ciprofloxacin (Table 1). The monohydroxamate conjugate 1c had activity against *S. aureus* (32  $\mu$ M), *K. pneumonia* (16  $\mu$ M), *A. baumannii* (16  $\mu$ M), *P. aeruginosa* (128  $\mu$ M), *E. aerogenes* (4  $\mu$ M), and *E. coli* (2  $\mu$ M),

Scheme 3. Syntheses of Hydroxamate Sideromycins  $(1b-3b \text{ and } 1c-3c)^a$ 



<sup>a</sup>Reagents and conditions: (a) 8, *i*Pr<sub>2</sub>EtN, DMAP, CH<sub>2</sub>Cl<sub>2</sub>; 18 h, 73% for 11; 16 h, 66% for 13; 20 h, 87% for 15. (b) 10% Pd–C, H<sub>2</sub> (1 atm), HCl (3 equiv), DMF:H<sub>2</sub>O (95:5); 41 h, 38% for 1b; 24 h, 42% for 2b; 24 h, 42% for 3b. (c) 10, *i*Pr<sub>2</sub>EtN, DMAP, CH<sub>2</sub>Cl<sub>2</sub>; 5 h, 92% for 12; 23 h, 66% for 14; 4.5 h, 84% for 16. (d) 10% Pd–C, H<sub>2</sub> (1 atm), MeOH; 12 h, 70% for 1c; 16 h, 86% for 2c; 16 h, 97% for 3c.

		test organisms						
entry	compound	E. faecium NCTC 7171	S. aureus SG 511	K. pneumonia ATCC 700603	A. baumannii ATCC 17961	P. aeruginosa ATCC 27853	E. aerogenes ATCC 35029	E. coli ATCC 25922
1	la	>128	>128	>128	>128	>128	>128	>128
2	2a	>128	>128	>128	>128	>128	>128	>128
3	3a	>128	>128	>128	>128	>128	>128	>128
4	3a-Fe	>128	>128	>128	>128	128	>128	>128
5	1b	>128	16	>128	≥128	>128	>128	>128
6	2b	>128	64	>128	32	>128	>128	≥128
7	3b	>128	64	>128	64	>128	>128	128
8	1c	>128	32	16	16	128	4	2
9	2c	>128	64	128	64	64	4	2
10	3c	64	1	>128	≥128	>128	64	>128
11	Lorabid	32	1	128	>128	>128	>128	2
12	ciprofloxacin	8	0.5	0.25	0.25	0.125	<0.015	<0.015

 $^{a}$ MIC values were determined using the broth microdilution method in Mueller-Hinton broth No. 2 (MHII) using visual end point analysis according to the CLSI guidelines.  $^{40}$  <sup>b</sup>Each compound was tested in triplicate.

but at significantly reduced potencies relative to ciprofloxacin (MIC values of 0.5, 0.25, 0.25, 0.125, <0.015, and <0.015 µM, respectively). The bishydroxamate conjugate 2c had similar activity to 1c against S. aureus (64 µM), K. pneumonia (128  $\mu$ M), A. baumannii (64  $\mu$ M), P. aeruginosa (64  $\mu$ M), E. aerogenes (4  $\mu$ M), and E. coli (2  $\mu$ M), but again at significantly reduced potencies compared to ciprofloxacin. Trihydroxamate conjugate 3c showed selective antibacterial activity toward S. aureus (MIC value of 1  $\mu$ M) at approximately equivalent potency as ciprofloxacin (MIC value of 0.5 uM). Mono- and bishydroxamate conjugates, 1c and 2c, had spectra of antibacterial activity similar to that of ciprofloxacin (broad spectrum with increased potency toward Gram-negative bacteria) at a reduced potency, while the trihydroxamate conjugate 3c had a narrow spectrum of activity primarily against Gram-positive bacteria, especially S. aureus.

Effect of Fe(III) Concentration on Sideromycin Antibacterial Activity. Since the hydroxamate-antibiotic conjugates (1b-3b and 1c-3c) all had activity toward S. aureus SG511, this organism was chosen for further experiments to determine if the synthetic sideromycins were actively transported into the bacterial cells via siderophore uptake pathways. Siderophore transport pathways are known to be upregulated during times of Fe(III)-restriction,41 and consequently, it has been discovered that the potency of antibacterial activity of a sideromycin increases (lower MIC value) under these circumstances.<sup>42</sup> As shown in Table 2, MIC values against S. aureus were determined for all the conjugates using Fe(III)-supplemented (MHII + Fe) and Fe(III)restricted (MHII - Fe) testing media.43 None of the MIC values observed for the  $\beta$ -lactam-hydroxamate conjugates (1b-3b) under varying Fe(III) concentrations changed significantly (a significant change for the broth microdilution assay is >2-fold). Only trihydroxamate conjugate 3c produced a significant (4-fold) decrease in MIC value when going from Fe(III)-supplemented media (4  $\mu$ M) to Fe(III)-restricted media (1  $\mu$ M). The MIC values for the control antibiotics, Lorabid and ciprofloxacin, showed no dependence on the Fe(III) concentration. These data suggest that the fluoroquinolone-trihydroxamate conjugate 3c might be utilizing siderophore-transport pathways.

Effect of Competing Siderophores on Sideromycin Antibacterial Activity. Sideromycin antibacterial activity is Table 2. MIC Values ( $\mu$ M) of Compounds 1b-3b and 1c-3c against *S. aureus* SG511 under Varying Concentrations of Fe(III)<sup>*a,b*</sup>

		test organism <i>Staphylococcus aureus</i> SG511		
entry	compound	MHII + $Fe^{c}$	MHII – Fe <sup>d</sup>	
1	1b	16	16	
2	2b	128	64	
3	3b	128	64	
4	1c	64	32	
5	2c	64	64	
6	3c	4	1	
8	Lorabid	1	1	
9	ciprofloxacin	0.5	0.5	

<sup>*a*</sup>MIC values were determined using the broth microdilution method using visual end point analysis according to the CLSI guidelines.<sup>40</sup> <sup>*b*</sup>Each compound was tested in triplicate. <sup>*c*</sup>MHII + Fe: Mueller-Hinton Broth No. 2 + 100  $\mu$ M FeCl<sub>3</sub>. <sup>*d*</sup>MHII – Fe: Mueller-Hinton Broth No. 2 + 100  $\mu$ M 2,2'-bipyridine.

known to be antagonized in the presence of siderophores competing for uptake via the same membrane transport systems.<sup>44</sup> To screen for this type of antagonism, MIC values against S. aureus were determined under Fe(III)-restricted conditions using 1:1 molar mixtures of the hydroxamateantibiotic conjugates (1b-3b and 1c-3c) with the corresponding parent hydroxamate compounds (1a-3a and 3a-Fe) and the commercially available trihydroxamate siderophore desferrioxamine B (DFO-B) or its Fe(III) complex ferrioxamine B (FO-B), as shown in Table 3. The siderophore DFO-B and siderophore-Fe(III) complex FO-B were chosen, since they are known to be utilized by S. aureus for siderophore-mediated Fe(III)-acquisition and are structurally similar to danoxamine, the siderophore portion of the salmycins.<sup>45</sup> The antibacterial activity of the  $\beta$ -lactam-monohydroxamate conjugate 1b was not antagonized by the monohydroxamate 1a (Table 3, Entry 1) or **DFO-B** (Table 3, Entry 2). The activity of the  $\beta$ -lactam– bishydroxamate conjugate 2b was not antagonized, but was enhanced by the presence of bishydroxamate 2a (Table 3, Entry 3) or DFO-B (Table 3, Entry 4). Similarly, the activity of the  $\beta$ -lactam-trihydroxamate conjugate 3b was not antagonized by the presence of trihydroxamate-Fe(III) complex 3a-Fe (Table 3, Entry 6) or FO-B (Table 3, Entry 8) and was slightly

# **Bioconjugate Chemistry**

Table 3. MIC Values ( $\mu$ M) of Compounds 1b-3b and 1c-3c against *S. aureus* SG511 in the Presence of Competing Siderophores<sup>*a*</sup>

		test organism Staphylococcus aureus SG511
entry	$compound^b$	MHII – Fe <sup>c</sup>
1	1b + 1a	16
2	1b + DFO-B	16
3	2b + 2a	32
4	2b + DFO-B	16
5	3b + 3a	32
6	3b + 3a-Fe	64
7	3b + DFO-B	32
8	3b + FO-B	64
9	1c + 1a	32
10	1c + DFO-B	16
11	2c + 2a	64
12	2c + DFO-B	64
13	3c + 3a	1
14	3c + 3a-Fe	1
15	3c + DFO-B	128
16	3c + FO-B	>128

"MIC values were determined using the broth microdilution method using visual end point analysis according to the CLSI guidelines.<sup>40</sup> Each compound was tested in triplicate. Data presented here should be compared to MIC values in Table 1 for compounds **1b**–**3b** and **1c**– **3c**. <sup>6</sup>Compounds were tested as 1:1 molar mixtures. <sup>c</sup>MHII – Fe: Mueller-Hinton Broth No. 2 + 100  $\mu$ M 2,2'-bipyridine.

enhanced by the trihydroxamate **3a** (Table 3, Entry 5) and **DFO-B** (Table 3, Entry 7).

Neither the mono- or bishydroxamate-fluoroquinolone conjugates 1c and 2c, respectively, were antagonized by the presence of the parent hydroxamate compounds (1a and 2a, respectively) or DFO-B (Table 3, Entries 9–12). The fluoroquinolone-trihydroxamate conjugate 1c was not antagonized by the parent trihydroxamate siderophore 3a (Table 3, Entry 13) or the Fe(III)-complex 3a-Fe (Table 3, Entry 14). However, conjugate 3c was strongly antagonized by the trihydroxamate siderophore DFO-B (Table 3, Entry 15) and the Fe(III)-complex FO-B up to >128-fold. Antagonism of the antibacterial activity of 3c by DFO-B and FO-B suggests that these molecules are competing for entrance into *S. aureus* cells via siderophore-uptake pathways.

The influence of competing siderophores 3a, 3a-Fe, DFO-B, and FO-B on the antibacterial activity of the  $\beta$ -lactamtrihydroxamate conjugate 3b (Figure 5a,b) and the fluoroquinolone-trihydroxamate conjugate 3c (Figure 5c,d) against S. aureus was also studied using a qualitative agar diffusion competition experiment (Figure 5). Sterile paper strips soaked in solutions of the test compounds were layed in a "criss-cross" pattern on Fe(III)-deficient MHII agar inoculated with S. aureus. After incubation and growth of the organism, analysis of the intersecting and nonintersecting points of the paper strips revealed the effects of the compound mixtures relative to the single compounds alone. Data from these studies were in agreement with the data from the MIC studies (Table 3). The antibacterial activity of the  $\beta$ -lactam-trihydroxamate conjugate 3b was not antagonized by the presence of 3a-Fe and FO-B and was slightly enhanced by the presence of 3a and DFO-B, while the activity of Lorabid was unaffected (Figure 5a,b). The antibacterial activity of the fluoroquinolone-trihydroxamate conjugate 3c was antagonized by all the siderophores (3a, 3aArticle



**Figure 5.** Agar diffusion competition assay for conjugates **3b** and **3c** and siderophores **3a**, **3a-Fe**, **DFO-B**, and **FO-B** tested against *S. aureus* SG511. Siderophores **3a**, **3a-Fe**, **DFO-B**, and **FO-B** clearly antagonize the anti-*S. aureus* activity of the fluoroquinolone conjugate **3c** (c,d), but not the  $\beta$ -lactam conjugate **3b** (a,b). Images are not to actual scale.

Fe, DFO-B, and FO-B), while ciprofloxacin was unaffected. In agreement with the MIC data (Table 3), DFO-B and FO-B antagonized the activity of 3c more than siderophores 3a and 3a-Fe (Figure 5c,d).

Further Sideromycin Antibacterial Susceptibility Testing in Agar Diffusion Assays. A more quantitative agar diffusion assay was used to further verify the observed affects of variable Fe(III) concentrations and competing siderophores on the anti-S. aureus activity of the hydroxamate-antibiotic conjugates (Figures 6 and 7). The quantitative agar diffusion assay used was a modified<sup>46</sup> Kirby-Bauer assay.<sup>47</sup> Compound solutions were added to 9 mm wells of an agar Petri dish inoculated with S. aureus SG511 according to the conditions indicated in the captions and color-coded legends of Figures 6 and 7. After incubation, the diameters of growth inhibition zones were measured and are reported on the vertical axes as the average of three independent trials. As shown in Figure 6, the antibiotic activities of the  $\beta$ -lactam-hydroxamate conjugates 1b-3b and Lorabid against S. aureus were not significantly affected by changes in the concentration of Fe(III) in the testing media or by the presence of competing siderophores. These observations are consistent with the MIC values discussed previously (Tables 2 and 3).

As shown in Figure 7, all three of the fluoroquinolonehydroxamate conjugates (1c-3c) showed decreased potency (smaller diameter zone of growth inhibition) against *S. aureus* in Fe(III)-supplemented media (MHII + Fe) relative to Fe(III)-restricted media (MHII - Fe). Furthermore, the potency of conjugates 1c-3c all decreased in the presence of an equimolar amount of the competing parent siderophores (1a-3a, respectively) or DFO-B. Again, in agreement with MIC data from Table 3, antagonism of the fluoroquinolonetrihydroxamate conjugate 3c by DFO-B and FO-B was more pronounced than by the parent siderophore 3a and its Fe(III)complex 3a-Fe. These data again strongly support that the trihydroxamate conjugate 3c is utilizing the same siderophoretransport pathway as DFO-B to deliver the fluoroquinolone antibiotic to its cytoplasmic target (DNA gyrase) in *S. aureus*.



Figure 6. Effect of varying Fe(III) concentrations and the presence of competing siderophores (1a, 2a, 3a, 3a-Fe, DFO-B) on the antibacterial activity of  $\beta$ -lactam sideromycins (1b, 2b, and 3b) against *S. aureus* SG511 in the agar diffusion assay. The vertical axis indicates the diameter of the growth inhibition zone in millimeters and the horizontal axis represents the test compound. The assay was performed in triplicate and standard deviations are indicated by error bars. MHII-Fe: Mueller-Hinton agar No. 2 + 100  $\mu$ M 2,2'-bipyridine. MHII+Fe: Mueller-Hinton agar No. 2 + 100  $\mu$ M FeCl<sub>3</sub>. Parent siderophore refers to the iron-free siderophore (1a, 2a, and 3a) with the same compound numeral as the corresponding sideromycin (1b, 2b, and 3b, respectively). Data for all mixtures corresponds to a 1:1 molar ratio of test compounds. 50  $\mu$ L of sample at the concentration indicated in parentheses under the compound numbers was added to 9 mm wells as described in the Experimental Section. The tabulated numerical data for this chart are provided in the Supporting Information (Tables S2 and S3).



Figure 7. Effect of varying Fe(III) concentrations and presence of competing siderophores (1a, 2a, 3a, 3a-Fe, DFO-B, and FO-B) on the antibacterial activity of fluoroquinolone sideromycins (1c, 2c, and 3c) against *S. aureus* SG511 in the agar diffusion assay. The vertical axis indicates the diameter of the growth inhibition zone in millimeters and the horizontal axis indicates the test compound. The assay was performed in triplicate. Ciprofloxacin was used as a control. MHII-Fe: Mueller-Hinton agar No. 2 + 100  $\mu$ M 2,2'-bipyridine. MHII+Fe: Mueller-Hinton agar No. 2 + 100  $\mu$ M FeCl<sub>3</sub>. Parent siderophore refers to the iron-free siderophore (1a, 2a, and 3a) with the same compound numeral as the corresponding sideromycin (1c, 2c, and 3c, respectively). Data for all mixtures correspond to a 1:1 molar ratio of test compounds. S0  $\mu$ L of sample at the concentration indicated in parentheses under the compound numbers was added to 9 mm wells as described in the Experimental Section. The tabulated numerical data for this chart are provided in the Supporting Information (Tables S2 and S3).

# FURTHER DISCUSSION

The hydroxamate  $-\beta$ -lactam (1b-3b) and hydroxamate fluoroquinolone conjugates (1c-3c) from this study all showed a narrowed spectrum of activity compared to the parent antibiotics. The hydroxamate-antibiotic conjugates showed high selectivity toward the Gram-positive pathogen Staphylococcus aureus SG511. The trihydroxamate-fluoroquinolone conjugate 3c showed the most selective activity and retained equivalent potency as the parent antibiotic, ciprofloxacin, against S. aureus (MIC = 1  $\mu$ M). Sideromycin 3c represents one of the few synthetic siderophore-antibiotic conjugates using an antibiotic with an intracellular target that maintains the potency of the parent drug.<sup>27,35</sup> The trihydroxamate siderophore, desferridanoxamine (3a), clearly was responsible for the narrowed spectrum of activity of the trihydroxamateciprofloxacin conjugate (3c). Analogously, desferridanoxamine is also the likely source of the exquisite Gram-positive selectivity observed for the salmycins which contain a normally

broad-spectrum aminoglycoside antibiotic. The precise reason for the selective antibacterial activity against Gram-positive bacteria is still up for debate, but one possibility is that the more promiscuous membrane-anchored hydroxamate siderophore binding proteins in *S. aureus*, known as FhuD1 and FhuD2, are capable of binding to the salmycins and the trihydroxamate—ciprofloxacin conjugate with high affinity, while the more discriminative hydroxamate siderophore outer-membrane proteins found in Gram-negative bacteria have reduced affinity for the derivatized siderophores.<sup>45,48,49</sup>

Article

Antibacterial susceptibility testing in both the broth microdilution and agar diffusion assays revealed that the antibacterial activity of the hydroxamate- $\beta$ -lactam conjugates (1b-3b) against *S. aureus* showed no dependence on Fe(III) concentration in the media and was unaffected by the presence of competing siderophores. These observations are consistent with the fact that the  $\beta$ -lactam biological targets, the penicillin binding proteins (PBPs), are in the periplasmic peptidoglycan which is outside of the cell membrane for S. aureus. Therefore, active transport into the S. aureus cells actually denied the  $\beta$ lactam sideromycins access to S. aureus PBPs. The hydroxamate-fluoroquinolone conjugates required active transport across the cell membrane to reach their biological target (DNA gyrase) in the cytoplasm. This clearly demonstrated that siderophore-antibiotic conjugates targeting Gram-positive bacteria should only use antibiotics with cytoplasmic targets. Furthermore, antibacterial activity of the trihydroxamateciprofloxacin conjugate (3c) showed a significant dependence on Fe(III) concentration in the testing media and was antagonized by competing siderophores, while the antibacterial activities of the mono- and bishydroxamate-ciprofloxacin conjugates (1c and 2c, respectively) were less influenced by Fe(III) and competing siderophores in the testing media. This suggested that the entire trihydroxamate backbone of the siderophore is necessary to facilitate efficient active transport of the sideromycin by S. aureus. The likely reason for needing the full trihydroxamate siderophore is for competitive chelation of environmental Fe(III) and recognition of the siderophore-Fe(III) complex by the siderophore-binding receptor proteins, which preferentially bind to octahedral, trihydroxamate-Fe(III) complexes.14

The ability of the ferrioxamine siderophore DFO-B and its Fe(III) complex FO-B to antagonize the anti-S. aureus activity of sideromycin 3c to a much greater extent than the structurally similar trihydroxamate parent siderophore desferridanoxamine, 3a, and its Fe(III)-complex danoxamine, 3a-Fe, is attributed to the drastic difference in Fe(III)-affinity observed for these molecules. Desferrioxamine B has an Fe(III)-affinity ( $\sim 10^{30}$ ) which is  $\sim 7$  orders of magnitude greater than the Fe(III)affinity of trihydroxamate siderophore 3a (~10<sup>23</sup>) (unpublished data from our laboratory). In work unpublished so far, we have shown that a competing siderophore must have an Fe(III)affinity equal to or greater than that of the sideromycin in order to antagonize the antibacterial activity of the sideromycin. This observation is attributed to the fact that the sideromycin can only be transported into bacterial cells, as the Fe(III)-complex and the molecule with the higher Fe(III)-affinity will "win" the competition for entry into the bacterial cell by "winning" the competition for binding the scarce Fe(III) nutrient.

Since a stable amide bond linkage was formed between the antibiotic ciprofloxacin and the siderophore desferridanoxamine, it was suspected that the entire desferridanoxamineciprofloxacin was the active antibacterial agent inhibiting the action of DNA gyrase. This hypothesis is supported in the literature by examples of siderophore-ciprofloxacin conjugates linked through an amide bond on the distal nitrogen of the piperazine unit that still show high in vitro binding to DNA gyrase.<sup>35</sup> The question of whether or not the aminoglycoside antibiotic of the salmycins is released inside bacterial cells is still up for debate, but it is proposed that the ideal synthetic sideromycin should feature a microbe-triggered linker that releases the antibiotic inside the cell to give optimal affinity for the biological target.<sup>27</sup> In fact, the development of new linker systems that make such intracellular release possible have been the focus of much work in our group.<sup>50</sup>

The trihydroxamate—ciprofloxacin sideromycin 3c developed in this work has many potential applications. The exquisite selectivity against *S. aureus* makes sideromycin 3c an ideal candidate for a targeted microbe-selective antibacterial chemotherapy that could limit the spread of fluoroquinolone multidrug resistance by limiting exposure of nontargeted bacteria to the active ingredient. Sideromycin 3c has the potential for decreasing mammalian toxicity of fluoroquinolones since it selectively enters S. aureus cells via membrane transport systems that are absent in eukaryotic cells.<sup>51</sup> Trihydroxamate ferrioxamine siderophores, including FDAapproved DFO-B, have been proven safe for human consumption at extremely large doses (4-12 g/day) when given to patients for chelation treatment of iron-overload diseases.<sup>52</sup> It has been documented that the use of DFO-B to treat iron-overload during the course of an S. aureus infection can worsen the infection since S. aureus can use the Fe(III)complex FO-B for iron assimilation.<sup>45</sup> Using sideromycin 3c to treat iron-overload diseases might be an effective method for simultaneously treating or preventing S. aureus infections. In fact, the salmycins have been patented for such an intended use.<sup>53,54</sup> Additionally, ferrioxamine siderophores are not bound by the mammalian siderophore-binding protein siderocalin, so the bioavailability is predicted to be high and consequently the required dosing to treat bacterial infections might be low.<sup>5</sup>

One foreseeable limitation of sideromycin antibiotics, and most pathogen-targeted antibiotic agents, is that they might not be effective at treating intracellular infections caused by pathogenic microbes that survive inside mammalian cells since the delivery agent is not engineered to penetrate higher eukaryotic cells.<sup>56</sup> Fortunately, alternative methods for treating these types of infections with engineered delivery systems are in development.<sup>57</sup> Another potential limitation to sideromycin antibiotic chemotherapy born from their exquisite microbe selectivity is the need for sensitive pathogen detection and rapid diagnostics. Unlike broad-spectrum antibiotics, microbe-selective antibiotics such as sideromycins will require early pathogen identification to direct accurate antibacterial drug prescription.<sup>2</sup> Fortunately, major advancements in bacterial pathogen detection using siderophore-based sensors are in development by several groups and will certainly play an important role in the future for sideromycin antibiotics in the clinic.<sup>58,59</sup>

# SUMMARY AND CONCLUSIONS

Similar to the salmycins, the synthetic linear hydroxamate sideromycins studied here showed selective antibacterial activity against Gram-positive bacteria, especially Staphylococcus aureus SG511 (MIC = 1  $\mu$ M for the trihydroxamate-fluoroquinolone sideromycin 3c). Siderophore-sideromycin competition assays and bacterial growth inhibition assays using variable Fe(III) concentrations showed that only the fluoroquinolone sideromycins (1c-3c) required membrane passage to reach their cytoplasmic biological target and that the full trihydroxamate siderophore delivery vector was required for siderophoreassociated protein-mediated active transport into S. aureus cells. This work supports the hypothesis that linear trihydroxamate siderophores are selective antibiotic delivery vectors for Grampositive pathogens, as observed for the naturally occurring salmycins and the synthetic desferridanoxamine-ciprofloxacin sideromycin (3c), and teaches how to engineer more effective hydroxamate-based sideromycins.

# EXPERIMENTAL SECTION

**Materials and Instrumentation.** All reactions were performed under a dry argon atmosphere, unless otherwise stated. All solvents and reagents were obtained from commercial sources and used without further purification unless otherwise stated. Dichloromethane  $(CH_2Cl_2)$  was

distilled from calcium hydride. Dimethylformamide (DMF) and diisopropylethylamine ( $iPr_2EtN$ ) were used from Acros Seal anhydrous bottles. *N*-Boc-*O*-PNB-Lorabid (7) was a gift from Eli Lilly and Company. Ciprofloxacin (9) was purchased from Sigma-Aldrich (St. Louis, MO). Benzyl-protected hydroxamate siderophores **4**–**6** were prepared according to a previously described method.<sup>29</sup> Silica gel column chromatography was performed using Sorbent Technologies silica gel 60 (32–63  $\mu$ m).

All microbiological media and liquids were sterilized by autoclaving (121 °C, 15 min) before use unless otherwise stated. All aqueous solutions and media were prepared using distilled, deionized, and filtered water (Millipore Milli-Q Advantage A10 Water Purification System). Luria broth (LB) was purchased from VWR. Mueller-Hinton No. 2 broth (MHII broth; cation adjusted) was purchased from Sigma-Aldrich (St. Louis, MO). Iron-deficient MHII broth (MHII - Fe) was prepared by adding 0.8 mL of a 1 mg/mL filter-sterilized aqueous solution of 2,2'-bipyridine to 49.2 mL of sterile MHII broth. Iron-supplemented MHII broth (MHII + Fe) was prepared by adding 0.8 mL of a freshly prepared 1 mg/mL filter-sterilized aqueous solution of FeCl<sub>3</sub> to 49.2 mL of sterile MHII broth. Mueller-Hinton No. 2 agar (MHII agar; HiMedia Laboratories) was purchased from VWR. Iron-deficient MHII agar (MHII – Fe) was prepared by adding 0.5 mL of a 1 mg/ mL filter-sterilized aqueous solution of 2,2'-biypridine to 34 mL of melted MHII agar with gentle mixing. Iron-supplemented MHII agar (MHII + Fe) was prepared by adding 0.5 mL of a freshly prepared 1 mg/mL filter-sterilized aqueous solution of FeCl<sub>3</sub> to 34 mL of melted MHII agar with gentle mixing. McFarland BaSO<sub>4</sub> turbidity standards were purchased from bioMérieux, Inc. Sterile plastic Petri dishes (145 mm × 20 mm; Greiner Bio-One) and sterile polystyrene 96-well plates (BD Falcon) used for antibacterial susceptibility testing were purchased from VWR. Ferric chloride (anhydrous, reagent grade) and 2,2'-bipyridine (reagent plus grade) were purchased from Sigma-Aldrich. All test organisms used in this work were from sources given in the Supporting Information (Table S1).

<sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were obtained on a 300, 500, or 600 MHz Varian DirectDrive spectrometer and FIDs were processed using ACD/ChemSketch version 10.04. Chemical shifts ( $\delta$ ) are given in parts per million (ppm) and are referenced to residual solvent. Coupling constants (J) are reported in hertz (Hz). High-resolution, accurate mass measurements were obtained with a Bruker micrOTOF II electrospray ionization time-of-flight mass spectrometer in positive ion mode. Sample was introduced via flow injection at a rate of 4  $\mu$ L/min, and mass spectra were accumulated from 50 to 3000 m/z for 2 min. Preparative HPLC purifications were performed on a Waters preparative binary pump system at a flow rate of 15 mL/min with UV detection at 254 nm using a YMC-Pack Pro C18 column (150  $\times$  20 mm; 5  $\mu$ m particle size) fit with a guard column. Thin layer chromatography (TLC) was performed with Al-backed Merck 60-F<sub>254</sub> or Al-backed Merck RP-C18 F<sub>256</sub> silica gel plates using a 254 nm lamp and aqueous FeCl<sub>3</sub> for visualization. Melting points were determined in capillary tubes using a Thomas-Hoover melting point apparatus and are uncorrected.

**Representative Synthetic Procedures for the Synthesis of Siderophores and Sideromycins.** *Desferridanoxamine (3a).* Prepared as described previously.<sup>29,30</sup> Briefly, tetra-*O*-benzyldanoxamine 6 (340.0 mg, 0.35 mmol) was dissolved in 35 mL MeOH in an HCl-washed round-bottom flask sealed under argon. The flask was charged with 10% Pd-C (50.0 mg) and exposed to a balloon of hydrogen gas ( $\sim 1$  atm). Reaction progress was monitored by RP-C18 TLC (1.5:1 CH<sub>3</sub>CN:H<sub>2</sub>O; FeCl<sub>3</sub> stain), and after 11 h, there was no remaining starting material (6). The flask was flushed with argon, and the mixture was diluted with warm MeOH, vacuum filtered through a pad of Celite using a fine glass frit, and concentrated under reduced pressure. The resulting solid was dissolved in a minimal amount of MeOH and precipitated by addition of cold Et<sub>2</sub>O. After trituration with  $Et_2O$ , the siderophore desferridanoxamine (3a) was obtained in 97% yield as a white solid (208.0 mg, 0.34 mmol). Mp 134–140 °C; <sup>1</sup>H NMR (600 MHz, DMSO- $d_6$ )  $\delta$ 11.87 (br s, COOH, 1 H), 9.62 (br s, N-OH, 3 H), 7.78 (t, J = 5.3 Hz, 2 H), 4.35 (br s, 1 H), 3.47-3.42 (m, 6 H), 3.39-3.36 (m, 2 H), 3.02–2.97 (m, 4 H), 2.57 (t, J = 6.6 Hz, 6 H), 2.39 (t, *J* = 6.6 Hz, 2 H), 2.26 (t, *J* = 6.9 Hz, 4 H), 1.49 (dt, *J* = 14.2, 7.1 Hz, 6 H), 1.43–1.34 (m, 6 H), 1.27–1.17 (m, 6 H); <sup>13</sup>C NMR  $(150 \text{ MHz}, \text{DMSO-}d_6) \delta$  174.1, 172.0, 171.9, 171.6, 171.3, 60.6, 47.2, 47.1, 38.4, 32.2, 29.9, 28.8, 28.7, 27.6, 27.5, 27.1, 26.2, 26.0, 23.5, 23.4, 22.7; HRMS-ESI (m/z): [M+H]<sup>+</sup> calcd. for C<sub>27</sub>H<sub>50</sub>N<sub>5</sub>O<sub>11</sub>: 620.3501, found 620.3506.

Danoxamine (**3a-Fe**). Prepared as described previously.<sup>29,30</sup> Briefly, siderophore **3a** (154 mg, 0.25 mmol) was dissolved in 10 mL of MeOH at 40 °C (oil bath temperature). Fe(acac)<sub>3</sub> (97.0 mg, 0.28 mmol) was added, and the clear, orange solution was stirred for 2 h. The solution was cooled to rt and the MeOH was evaporated under reduced pressure. The resulting orange residue was dissolved in a minimal amount of MeOH and precipitated by addition of cold Et<sub>2</sub>O. After trituration with Et<sub>2</sub>O, the Fe(III)-complexed siderophore danoxamine (**3a-Fe**) was isolated in 99% yield as an orange powder (165.2 mg, 0.25 mmol). Mp 151–154 °C (dec.); HRMS-ESI (*m*/*z*): [M+H]<sup>+</sup> calcd. for C<sub>27</sub>H<sub>47</sub>FeN<sub>5</sub>O<sub>11</sub>: 673.2616, found 673.2647.

O-PNB-Lorabid TFA Salt (8). N-Boc-O-PNB-Lorabid (7; 208.0 mg, 0.36 mmol) was dissolved in 7 mL of anhydrous TFA:CH<sub>2</sub>Cl<sub>2</sub> (1:1). After 30 min, TLC (2:1 hexanes:EtOAc) showed complete consumption of the starting material. The TFA/CH<sub>2</sub>Cl<sub>2</sub> were removed by evaporation under reduced pressure giving TFA salt 8 as a light yellow viscous oil (208.0 mg, 0.36 mmol) that was used immediately in the next reaction without purification or characterization.

O-Benzyl-ciprofloxacin HCl Salt (10). Prepared by a modified literature procedure.<sup>31</sup> Ciprofloxacin (9; 1.01 g, 3.05 mmol), Boc anhydride (820.0 mg, 3.76 mmol), and NaHCO<sub>3</sub> (1.28 g, 15.24 mmol) were suspended in 25 mL of anhydrous DMF and stirred at rt for 2 h. Benzyl bromide (1.62 mL, 13.64 mL) was added and the mixture was heated to 90 °C (oil bath temperature). After 1 h of heating, the reaction mixture became homogeneous. After 25.5 h of heating, the mixture was cooled to rt and the DMF was removed using high-vacuum rotary evaporation ( $\sim$ 1 mmHg). The resulting solid was dissolved in 75 mL of CHCl<sub>3</sub> and 50 mL of H<sub>2</sub>O. The layers were separated and the CHCl<sub>3</sub> was washed with  $H_2O$  (25 mL) and brine (25 mL), dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated to give an orange solid. The crude solid was suspended in Et<sub>2</sub>O, filtered, and washed several times with Et<sub>2</sub>O. This provided N-Boc-O-benzyl-ciprofloxacin in 98% yield as a pale yellow solid (1.56 g, 2.99 mmol). Mp 191-193 °C; <sup>1</sup>H NMR (600 MHz,  $CDCl_3$ )  $\delta$  8.52 (s, 1 H), 8.05 (d, J = 13.5 Hz, 1 H), 7.52 (d, J = 7.0 Hz, 2 H), 7.38 (t, J = 7.5 Hz, 2 H), 7.33–7.29 (m, 1 H), 7.26 (d, J = 7.0 Hz, 1 H), 5.39 (s, 2 H), 3.69–3.62 (m, 4 H), 3.44-3.37 (m, 1 H), 3.24-3.19 (m, 4 H), 1.50 (s, 9 H), 1.32-1.27 (m, 2 H), 1.15-1.09 (m, 2 H); <sup>13</sup>C NMR (150 MHz,

CDCl<sub>3</sub>)  $\delta$  173.0, 165.5, 154.6, 153.3 (d,  $J_{C-F}$  = 248.5 Hz), 148.3, 144.4 (d,  $J_{C-F}$  = 10.1 Hz), 137.9, 136.4, 133.3, 129.3, 128.5, 127.9, 127.8, 123.3 (d,  $J_{C-F} = 6.2$  Hz), 113.4 (d,  $J_{C-F} =$ 24.1 Hz), 110.2, 105.0, 80.2, 66.3, 49.9, 34.5, 28.4, 8.1; HRMS-ESI (m/z):  $[M+H]^+$  calcd. for C<sub>29</sub>H<sub>33</sub>FN<sub>3</sub>O<sub>5</sub>: 522.2399, found 522.2423. The solid (1.50 g, 2.88 mmol) was dissolved in 25 mL of  $CH_2Cl_2$  and concentrated HCl (5 mL) was added slowly giving a dark orange solution. After 40 min, the CH<sub>2</sub>Cl<sub>2</sub>/HCl were evaporated giving a yellow solid. The solid was dissolved in hot MeOH (25 mL), activated charcoal was added, and the mixture was vacuum filtered through Celite. The MeOH was evaporated and the solid was recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/Et<sub>2</sub>O at -20 °C, which gave the desired O-benzyl-ciprofloxacin monohydrochloride salt (10) in 92% as a yellow solid (1.20 g, 2.62 mmol). Mp 250-255 °C (dec.); <sup>1</sup>H NMR (600 MHz, DMSO- $d_6$ )  $\delta$  9.71 (br s, 1 H), 9.62 (br s, 1 H), 8.48 (d, J = 3.5Hz, 1 H), 7.84-7.77 (m, 1 H), 7.51-7.44 (m, 3 H), 7.42-7.36 (m, 2 H), 7.32 (t, J = 7.0 Hz, 1 H), 5.27 (s, 2 H), 3.65–3.53 (m, 1 H), 3.53-3.46 (m, 4 H), 3.34-3.24 (m, 4 H), 1.28-1.23 (m, 2 H), 1.12–1.07 (m, 2 H); <sup>13</sup>C NMR (150 MHz, DMSO-d<sub>6</sub>)  $\delta$  171.6, 164.5, 152.4 (d,  $J_{C-F}$  = 246.8 Hz), 148.6, 142.8 (d,  $J_{C-F} = 10.7 \text{ Hz}$ ), 138.0, 136.6, 128.4, 127.7, 127.6, 122.5, 111.8 (d,  $J_{C-F} = 24.1$  Hz), 108.9, 106.7, 65.2, 46.4, 42.5, 34.9, 7.6; HRMS-ESI (m/z):  $[M+H]^+$  calcd. for C<sub>24</sub>H<sub>25</sub>FN<sub>3</sub>O<sub>3</sub>: 422.1874, found 422.1871.

tetra-O-Benzyldanoxamine-O-PNB-Lorabid Conjugate (15). O-PNB-Lorabid TFA salt (8; 85.7 mg, 0.15 mmol) was dissolved in 10 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> and iPr<sub>2</sub>EtN (0.13 mL, 0.75 mmol) was added slowly until the solution was basic (pH paper). tetra-O-Benzyl-trihydroxamate 6 (155.5 mg, 0.16 mmol), DMAP (8.0 mg, 0.06 mmol), and EDC-HCl (95.0 mg, 0.50 mmol) were then added, respectively. After 20 h at rt, TLC (3% MeOH in EtOAc; FeCl<sub>3</sub> stain) showed no remaining starting material (8). The CH<sub>2</sub>Cl<sub>2</sub> was removed under reduced pressure and the resulting oil was dissolved in EtOAc (20 mL), washed with 1 N HCl<sub>ag</sub> (15 mL), H<sub>2</sub>O (10 mL), 10% aqueous NaHCO<sub>3</sub> (10 mL), and brine (10 mL), dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated. The crude product was purified by silica gel column chromatography  $(1 \times 6$  in silica gel; 3%-5% MeOH in EtOAc) to give the desired product (15) in 87% yield as a clear, colorless oil (183.0 mg, 0.100.13) mmol). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.61 (d, J = 7.4 Hz, 1 H), 8.21 (d, J = 8.6 Hz, 2 H), 7.60 (d, J = 8.8 Hz, 2 H), 7.45-7.24 (m, 25 H), 6.75 (br s, 1 H), 6.45 (br s, 1 H), 5.65 (d, J = 7.0 Hz, 1 H), 5.46-5.42 (m, 1 H), 5.41 (d, I = 14.4 Hz, 1 H), 5.33–5.28 (m, 1 H), 4.86–4.77 (m, 5 H), 4.75 (d, J = 10.2 Hz, 1 H), 4.47 (s, 2 H), 4.03-3.93 (m, 1 H), 3.91-3.85 (m, 1 H), 3.66-3.56 (m, 2 H), 3.52-3.46 (m, 1 H), 3.46-3.37 (m, 3 H), 3.34-3.10 (m, 4 H), 3.00-2.92 (m, 1 H), 2.87-2.71 (m, 6 H), 2.65-2.50 (m, 5 H), 2.51-2.41 (m, 4 H), 1.68-1.55 (m, 6 H), 1.55–1.46 (m, 2 H), 1.46–1.39 (m, 3 H), 1.39–1.30 (m, 4 H), 1.24-1.16 (m, 4 H), 1.15-1.06 (m, 1 H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 174.3, 174.2, 173.8, 172.4, 172.2, 171.9, 171.4, 165.2, 160.0, 147.8, 142.1, 138.5, 136.9, 134.4, 134.2, 134.1, 130.8, 129.2, 129.1, 129.1, 129.1, 129.1, 129.0, 128.9, 128.8, 128.8, 128.7, 128.7, 128.7, 128.6, 128.6, 128.5, 128.3, 128.2, 127.7, 127.5, 127.4, 123.9, 123.7, 123.6, 123.1, 77.2, 76.3, 76.1, 72.8, 70.1, 66.2, 58.8, 57.6, 52.8, 45.5, 44.8, 43.7, 39.3, 39.2, 31.9, 30.6, 30.3, 30.1, 29.6, 29.3, 28.8, 28.1, 27.8, 27.7, 27.5, 26.7, 26.2, 25.7, 23.7, 23.4, 23.3, 21.3; HRMS-ESI (m/z): [M  $+H]^+$  calcd. for C<sub>78</sub>H<sub>93</sub>ClN<sub>9</sub>O<sub>16</sub>: 1446.6423, found 1446.6416. Desferridanoxamine-Lorabid Conjugate (3b). tetra-O-

mmol) and concentrated HCl (33.0  $\mu$ L, 24.0 mmol) were dissolved in 1.1 mL of DMF:H<sub>2</sub>O (95:5) in an HCl-washed, 10 mL round-bottom flask sealed under argon. The flask was charged with 10% Pd–C (23.0 mg) and exposed to a balloon of hydrogen gas ( $\sim 1$  atm). Reaction progress was monitored by RP-C18 TLC (2.5:1 H<sub>2</sub>O:CH<sub>3</sub>CN; FeCl<sub>3</sub> stain) and after 24 h there was no remaining starting material (15). The flask was flushed with argon and the mixture was diluted with MeOH, vacuum filtered through Celite, and concentrated using highvacuum rotary evaporation ( $\sim$ 1 mmHg). The crude product was sequentially purified by size exclusion chromatography (0.5 × 6 in Sephadex G50; 1:1 MeOH:EtOAc eluent), preparative HPLC using a  $150 \times 20$  mm YMC-Pack Proc C18 column fit with a guard column, 0.1% TFA in  $\mathrm{H_{2}O}$  (A) and 0.1% TFA in CH<sub>3</sub>CN (B) as mobile phases, and a flow rate of 15 mL/min. A gradient was formed from 20-50% of B over 4 min where the desired compound (3b) eluted at 4.1 min. Pure fractions were lyophilized and the obtained solid was recrystallized from MeOH/Et<sub>2</sub>O to give the desired product (3b) in 42% yield as an off-white solid (30.0 mg, 0.03 mmol). Mp 103–105  $^\circ \mathrm{C}$ (color change), 170-175 °C (dec.); <sup>1</sup>H NMR (600 MHz, CD<sub>3</sub>OD) δ 7.45–7.32 (m, 5 H), 5.43–5.42 (m, 1 H), 5.37 (d, J = 5.0 Hz, 1 H), 3.88 (td, J = 7.9, 3.8 Hz, 1 H), 3.83 (ddd, J = 14.0, 7.3, 7.1 Hz, 1 H), 3.62–3.53 (m, 4 H), 3.30–3.26 (m, 1 H), 3.18-3.13 (m, 4 H), 2.84 (t, J = 6.3 Hz, 2 H), 2.76 (td, J =7.1, 3.1 Hz, 3 H), 2.70–2.54 (m, 5 H), 2.53 (d, J = 5.6 Hz, 1 H), 2.50–2.43 (m, 4 H), 2.40 (dt, J = 15.2, 5.5 Hz, 1 H), 1.78– 1.72 (m, 2 H), 1.68-1.57 (m, 6 H), 1.57-1.54 (m, 2 H), 1.54-1.47 (m, 4 H), 1.39–1.35 (m, 2 H), 1.34–1.27 (m, 4 H); <sup>13</sup>C NMR (150 MHz, CD<sub>3</sub>OD) δ 175.6, 175.1, 174.6, 174.6, 174.6, 174.5, 173.9, 166.3, 163.9, 138.3, 130.0, 129.7, 129.4, 129.1, 126.0, 62.9, 59.9, 59.6, 54.1, 40.5, 40.4, 33.4, 32.5, 31.7, 31.6, 31.3, 30.5, 30.2, 30.1, 30.0, 29.9, 29.1, 29.0, 28.9, 27.6, 27.4, 27.2, 25.0, 24.8, 24.1, 23.1; HRMS-ESI (*m*/*z*): [M+H]<sup>+</sup> calcd. for C43H64ClN8O14: 951.4225, found 951.4234.

tetra-O-Benzyldanoxamine-O-benzylciprofloxacin Conjugate (16). O-Benzyl-ciprofloxacin hydrochloride salt (10) was free-based using Amberlite IR400(OH<sup>-</sup>) resin in CHCl<sub>3</sub> for 4 h. The resulting O-benzyl-ciprofloxacin amine (46.0 mg, 0.11 mmol), tetra-O-Benzyl-trihydroxamate 6 (103.0 mg, 0.10 mmol), *i*Pr<sub>2</sub>EtN (0.04 mL, 0.23 mmol), DMAP (3.2 mg, 0.03 mmol), and EDC-HCl (30.2 mg, 0.16 mmol) were dissolved in 10 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub>, respectively. After 4.5 h at rt, TLC (9% MeOH in CH<sub>2</sub>Cl<sub>2</sub>; FeCl<sub>3</sub> stain) showed no remaining starting material 6. The mixture was diluted with  $CH_2Cl_2$  (20 mL), washed with 10% aqueous citric acid (15 mL), H<sub>2</sub>O (15 mL), 10% aqueous NaHCO<sub>3</sub> (15 mL), and brine (15 mL), dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated. The crude product was purified by silica gel column chromatography  $(1 \times 4 \text{ in silica gel; } 3\% \text{ MeOH in})$  $CH_2Cl_2$ ) to give the desired product (16) in 84% yield as a clear, yellow oil (121.5 mg, 0.09 mmol). <sup>1</sup>H NMR (600 MHz,  $CDCl_3$ )  $\delta$  8.53 (s, 1 H), 8.07 (d, J = 12.9 Hz, 1 H), 7.52 (d, J = 7.3 Hz, 2 H), 7.43-7.24 (m, 24 H), 6.46 (br s, 1 H), 6.33 (br s, 1 H), 5.39 (s, 2 H), 4.91 (s, 2 H), 4.84 (s, 4 H), 4.47 (s, 2 H), 3.86-3.82 (m, 2 H), 3.76-3.73 (m, 2 H), 3.69-3.58 (m, 6 H), 3.44 (t, J = 6.6 Hz, 2 H), 3.42–3.37 (m, 1 H), 3.31–3.27 (m, 2 H), 3.24–3.16 (m, 6 H), 2.86–2.76 (m, 6 H), 2.70 (t, J = 6.0 Hz, 2 H), 2.47 (t, J = 6.2 Hz, 4 H), 1.68–1.58 (m, 8 H), 1.52– 1.45 (m, 4 H), 1.42–1.24 (m, 8 H), 1.12–1.08 (m, 2 H); <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>) δ 174.0, 173.8, 173.7, 173.0, 172.9, 172.1, 172.0, 170.4, 165.5, 153.3 (d,  $J_{C-F}$  = 249.1 Hz), 148.3, 144.1 (d,  $J_{C-F} = 10.7$  Hz), 138.5, 137.9, 136.4, 134.5, 134.3,

129.1, 129.0, 128.9, 128.8, 128.8, 128.6, 128.5, 128.3, 127.9, 127.8, 127.5, 127.4, 123.5 (d,  $J_{C-F} = 6.7$  Hz), 113.5 (d,  $J_{C-F} = 23.6$  Hz), 110.2, 105.1 (d,  $J_{C-F} = 2.24$  Hz), 76.4, 76.3, 76.2, 72.8, 70.1, 66.3, 50.3, 50.2, 49.6, 49.5, 45.7, 45.5, 45.2, 45.1, 44.8, 41.5, 39.3, 34.5, 30.7, 30.5, 29.3, 29.0, 28.5, 28.1, 27.9, 27.3, 27.2, 26.6, 26.5, 26.4, 23.9, 23.6, 23.4, 8.1; HRMS-ESI (m/z):  $[M+H]^+$  calcd. for  $C_{79}H_{96}FN_8O_{13}$ : 1383.7075, found 1383.7064.

Desferridanoxamine-Ciprofloxacin Conjugate (3c). tetra-O-Benzyl-trihydroxamate-O-benzylciprofloxacin conjugate (16; 55.0 mg, 0.04 mmol) was dissolved in 6 mL of MeOH in an HCl-washed, 10 mL rb flask sealed under argon. The flask was charged with 10% Pd-C (15.0 mg) and exposed to a balloon of hydrogen gas (~1 atm). Reaction progress was monitored by RP-C18 TLC (1.5:1 CH<sub>3</sub>CN:H<sub>2</sub>O; FeCl<sub>3</sub> stain) and after 16 h there was no remaining starting material (16). The flask was flushed with argon and the mixture was diluted with MeOH, vacuum filtered through Celite, and concentrated under reduced pressure. The resulting solid was dissolved in a minimal amount of MeOH and precipitated by addition of cold  $Et_2O$ . After trituration with  $Et_2O$ , the desired product (3c) was obtained in 97% yield as a light yellow solid (36.1 mg, 0.039 mmol). Mp 94–96 °C (color change), 153–160 °C (dec.); <sup>1</sup>H NMR (600 MHz, DMSO- $d_6$ )  $\delta$  9.66 (br s, N–OH, 1 H), 9.62 (br s, N–OH, 2 H), 8.67 (s, 1 H), 7.93 (d, I = 13.2 Hz, 1 H), 7.78 (t, J = 4.4 Hz, 1 H), 7.58 (d, J = 7.3 Hz, 1 H), 4.35 (t, J =5.0 Hz, 1 H), 4.04 (s, 1 H), 3.86–3.79 (m, 1 H), 3.74–3.66 (m, 4 H), 3.49-3.41 (m, 6 H), 3.39-3.34 (m, 4 H), 3.31-3.27 (m, 2 H), 3.03-2.95 (m, 4 H), 2.65 (t, J = 6.5 Hz, 2 H), 2.61-2.53(m, 6 H), 2.30–2.21 (m, 4 H), 1.54–1.45 (m, 6 H), 1.43–1.34 (m, 6 H), 1.34–1.30 (m, 2 H), 1.27–1.16 (m, 8 H); <sup>13</sup>C NMR (150 MHz, DMSO-d<sub>6</sub>) δ 176.3, 172.0, 171.9, 171.9, 171.3, 171.2, 170.2, 165.9, 152.9 (d,  $J_{C-F}$  = 249.1 Hz), 148.1, 144.9 (d,  $J_{C-F} = 10.7$  Hz), 139.1, 111.0 (d,  $J_{C-F} = 23.0$  Hz), 106.6, 60.6, 60.5, 49.6, 49.6, 49.2, 47.2, 47.1, 47.1, 44.4, 40.8, 38.4, 38.4, 38.3, 38.3, 35.9, 32.2, 32.1, 29.9, 29.8, 28.8, 27.6, 27.2, 26.9, 26.2, 26.1, 26.0, 23.5, 22.7, 7.6; HRMS-ESI (m/z): [M+H]<sup>+</sup> calcd. for C44H66N8O14: 933.4728, found 933.4744.

Determination of MIC Values by the Broth Microdilution Method. Antibacterial activity of the compounds was determined by measuring their minimum inhibitory concentrations (MICs) using the broth microdilution method according to the Clinical and Laboratory Standards Institute (CLSI, formerly the NCCLS) guidelines.<sup>40</sup> Each well of a 96well microtiter plate was filled with 50  $\mu$ L of sterile broth media (MHII, MHII – Fe, or MHII + Fe). Each test compound was dissolved in DMSO or H<sub>2</sub>O making a 20 mM solution, then diluted with sterile MHII broth to 512  $\mu$ M. Exactly 50  $\mu$ L of the compound solution was added to the first well of the microtiter plate and 2-fold serial dilutions were made down each row of the plate. Exactly 50  $\mu$ L of bacterial inoculum (5 × 10<sup>5</sup> CFU/ mL) was then added to each well giving a total volume of 100  $\mu$ L/well and a final compound concentration gradient of 128  $\mu$ M-0.0625  $\mu$ M. The plate was incubated at 37 °C for 18 h and then each well was examined for bacterial growth. The MIC was recorded as the lowest compound concentration  $(\mu M)$ inhibiting visible bacterial growth as judged by turbidity of the culture media relative to a row of wells filled with a DMSO standard. Ciprofloxacin was included in a control row at a concentration gradient of 32–0.0156  $\mu$ M as well as a solvent control row.

For MIC studies using mixtures of compounds, both test compounds were diluted to 20 mM in DMSO as one

homogeneous solution. The DMSO solution was then diluted with sterile broth media to give a solution at 512  $\mu$ M for both compounds. Finally, 50  $\mu$ L of the mixed compound broth solution was used in the assay exactly as described previously for single compound experiments.

General Procedure for the Agar Diffusion Paper Strip Siderophore–Sideromycin Competition Assay. A culture of S. aureus SG511 was grown in LB broth for 18-24 h and a standard cell suspension of  $1.5 \times 10^6$  CFU/mL was prepared in saline solution (0.9% NaCl) according to a 0.5 BaSO<sub>4</sub> McFarland Standard.<sup>60</sup> The standardized cell suspension (0.1 mL) was added to 34 mL of sterile, melted MHII - Fe irondeficient agar tempered to 47-50 °C. After gentle mixing, the inoculated agar media was poured into a sterile plastic Petri dish (145 mm  $\times$  20 mm) and allowed to solidify near a flame with the lid cracked for  $\sim 30$  min. Solutions of the test compounds dissolved in a 1:10 mixture of DMSO:MeOH (3a, 3b, and 3c) or sterile, distilled, and deionized  $H_2O$  (3a-Fe, DFO-B, FO-B, Lorabid, and ciprofloxacin) were prepared at the desired concentration (2.0 mM for 3a, 3a-Fe, 3b, DFO-B, and FO-B; 0.5 mM for 3c; 0.1 mM for Lorabid; 5  $\mu$ g/mL for ciprofloxacin). Sterile filter paper strips (Whatman No. 1 Standard grade; cut to  $\sim 1 \times 8$  cm) were soaked in the test compound and laid on the surface of the inoculated agar media as shown in Figure 5. The Petri dishes were incubated at 37 °C for 18-24 h and photographed after 48 h.

General Procedure for the Quantitative Agar Diffusion Antibiotic Susceptibility Assay. Antibacterial activity of the compounds was determined by a modified Kirby-Bauer agar diffusion assay.46,47 Overnight cultures of test organisms were grown in LB broth for 18-24 h and standard cell suspensions of  $1.5 \times 10^6$  CFU/mL were prepared in saline solution (0.9% NaCl) according to a 0.5 BaSO<sub>4</sub> McFarland Standard.<sup>60</sup> Each standardized cell suspension (0.1 mL) was added to 34 mL of sterile, melted agar (MHII, MHII - Fe, or MHII + Fe) tempered to 47-50 °C. After gentle mixing, the inoculated agar media was poured into a sterile plastic Petri dish (145 mm  $\times$  20 mm) and allowed to solidify near a flame with the lid cracked for  $\sim$ 30 min. Wells of 9.0 mm diameter were cut from the Petri dish agar and filled with exactly 50  $\mu$ L of the test sample solution. For studies using mixtures of compounds, both test compounds were diluted to equal concentrations as one homogeneous solution and 50  $\mu$ L of the mixed compound solution was added to the sample well. The Petri dish was incubated at 37 °C for 18-24 h and the inhibition zone diameters were measured (mm) with an electronic caliper after 24-48 h.

# ASSOCIATED CONTENT

#### Supporting Information

List of strains, markers, and origins for microorganisms used in this work (Table S1). Tabulated biological data collected during this study (Tables S2–S3). Experimental procedures for the syntheses of compounds 1a–c, 2a–c, and 11–14. Copies of <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra for compounds 1a-c, 2a-c, 3a-c, and 11–16. This material is available free of charge via the Internet at http://pubs.acs.org.

# AUTHOR INFORMATION

#### **Corresponding Author**

\*Phone (574) 631-7571; fax (574) 631-6652; e-mail mmiller1@nd.edu.

# **Bioconjugate Chemistry**

#### Present Address

<sup>#</sup>Department of Pharmaceutical and Biomedical Sciences, University of Georgia, Athens, GA 30602, United States.

#### Notes

The authors declare no competing financial interest.

# ACKNOWLEDGMENTS

We gratefully acknowledge the use of NMR facilities provided by the Lizzadro Magnetic Resonance Research Center at The University of Notre Dame (UND) and the mass spectrometry services provided by The UND Mass Spectrometry & Proteomics Facility (Mrs. N. Sevova, Dr. W. Boggess, and Dr. M. V. Joyce; supported by the National Science Foundation under CHE-0741793). We thank Dr. Viktor Krchňák (UND) for assistance with HPLC. We are grateful to Prof. Shahriar Mobashery (UND) and Dr. Sergei Vakulenko (UND) for providing the ESKAPEE panel of bacteria and Dr. Nuno Tiago GiaoAntunes (UND) for determination of some preliminary MIC values against the panel. TAW gratefully acknowledges The UND Chemistry-Biochemistry-Biology (CBBI) Interface Program funded by the NIH (T32GM075762) for three years of fellowship support and funding an internship at the Hans Knöll Institute (HKI) in Jena, Germany to learn microbiological assays. T.A.W. sincerely thanks the HKI for hosting such an internship. T.A.W. also thanks the UND Department of Chemistry and Biochemistry (Grace Fellowship), the UND Center for Environmental Science and Technology (CEST), and Bayer for additional fellowship support. We acknowledge The University of Notre Dame and the NIH for financial support of this work.

# ABBREVIATIONS

ABC, ATP-binding cassette transporter; acac, acetylacetonate; ATCC, American Type Culture Collection; Bn, benzyl; Boc, tert-butoxycarbonyl; CFU, colony forming unit; DFO-B, desferrioxamine B; DMAP, 4-dimethylaminopyridine; DMF, dimethylformamide; DMSO, dimethylsulfoxide; ESI, electrospray ionization; ExbB/D, inner membrane bound protein of TonB complex; FO-B, ferrioxamine B; Fur, ferric uptake regulator protein; HPLC, high-performance liquid chromatography; HRMS, high-resolution mass spectrometry; MHII, Mueller-Hinton media no. II; MIC, minimum inhibitory concentration; NMR, nuclear magnetic resonance; OMR, outer membrane receptor protein; OMT, TonB-dependent outer membrane siderophore transport protein; PNB, para-nitrobenzyl; Red, iron reductase enzyme; SPBP, siderophore periplasmic binding protein; TFA, trifluoroacetic acid; TLC, thin layer chromatography; TonB, membrane-spanning protein complex involved in siderophore transport

## REFERENCES

(1) Boucher, H. W., Talbot, G. H., Bradley, J. S., Edwards, J. E., Jr., Gilbert, D., Rice, L. B., Scheld, M., Spellberg, B., and Bartlett, J. (2009) Bad bugs, no drugs: No ESKAPE! An update from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 48, 1–12.

(2) Fischback, M. A., and Walsh, C. T. (2009) Antibiotics for emerging pathogens. *Science* 325, 1089–1093.

(3) Payne, D. J., Gwynn, M. N., Holmes, D. J., and Pompliano, D. L. (2007) Drugs for bad bugs: Confronting the challenges of antibacterial discovery. *Nat. Rev. Drug Discovery* 6, 29–40.

(4) Pereira, M. P., and Kelley, S. O. (2011) Maximizing the therapeutic window of an antimicrobial drug by imparting

mitochondrial sequestration in human cells. J. Am. Chem. Soc. 133, 3260–3263.

(5) Silver, L. L. (2008) Are natural products still the best source for antibacterial discovery? The bacterial entry factor. *Expert Opin. Drug Discovery* 3, 487–500.

(6) O'Shea, R., and Moser, H. E. (2008) Physiochemical properties of antibacterial compounds: Implications in drug discovery. *J. Med. Chem.* 51, 2871–2878.

(7) Ranny, D. F. (2000) Biomimetic transport and rational drug delivery. *Biochem. Pharmacol.* 59, 105–114.

(8) Christofferson, R. E. (2006) Antibiotics: An investment worth making? Nat. Biotechnol. 24, 1512–1514.

(9) Braun, V., and Endri $\beta$ , F. (2007) Energy-coupled outer membrane transport proteins and regulatory proteins. *Biometals 20*, 219–231.

(10) Ji, C., Juárez-Hernández, R. E., and Miller, M. J. (2012) Exploiting bacterial iron acquisition: siderophore conjugates. *Future Med. Chem.* 4, 297–313.

(11) Roosenberg, J. M., Lin, Y.-M., Lu, Y., and Miller, M. J. (2000) Studies and syntheses of siderophores, microbial iron chelators, and analogs as potential drug delivery agents. *Curr. Med. Chem.* 7, 159–197.

(12) Frederick, R. E., Mayfield, J. A., and DuBois, J. L. (2009) Iron trafficking as an antimicrobial target. *Biometals* 22, 583–593.

(13) Miethke, M., and Marahiel, M. A. (2007) Siderophore-based iron acquisition and pathogen control. *Microbiol. Mol. Biol. Rev.* 71, 413–451.

(14) Hider, R. C., and Kong, X. (2010) Chemistry and biology of siderophores. *Nat. Prod. Rep.* 27, 637–657.

(15) Wandersman, C., and Delepelaire, P. (2004) Bacterial iron sources: From siderophores to hemophores. *Annu. Rev. Microbiol.* 58, 611–647.

(16) Challis, G. L., and Hopwood, D. A. (2003) Synergy and contingency as driving forces for the evolution of multiple secondary metabolite production by *Streptomyces* species. *Proc. Natl. Acad. Sci.* U.S.A. 100, 14555–14561.

(17) Braun, V., Pramanik, A., Gwinner, T., Köberle, M., and Bohn, E. (2009) Sideromycins: Tools and antibiotics. *Biometals* 22, 3–13.

(18) Reynolds, D. M., Schatz, A., and Waksman, S. A. (1947) Grisein, a new antibiotic produced by a strain of *Streptomyces griseus*. *Proc. Soc. Exp. Biol. Med.* 64, 50–54.

(19) Benz, G., Schroder, T., Kurz, J., Wünsche, C., Karl, W., Steffens, G., Pfitzner, J., and Schmidt, D. (1982) Konstitution der deferriform der albomycine  $\delta_1$ ,  $\delta_2$ , und  $\varepsilon$ . Angew. Chem., Int. Ed. Engl. 21, 527–528. (20) Ferguson, A. D., Braun, V., Fiedler, H. P., Coulton, J. W., Diedrichs, K., and Welte, W. (2000) Crystal structure of the antibiotic albomycin in complex with the outer membrane transporter FhuA. *Protein Sci. 9*, 956–963.

(21) Clarke, T. E., Braun, V., Winkelmann, G., Tari, L. W., and Vogel, H. J. (2002) X-ray crystallographic structures of the *Escherichia coli* periplasmic protein FhuD bound to hydroxamate-type siderophores and the antibiotic albomycin. *J. Biol. Chem.* 277, 13966–13972.

(22) Braun, V., Günthner, K., Hantke, K., and Zimmermann, L. (1983) Intracellular activation of albomycin in *Escherichia coli* and *Salmonella typhimurium. J. Bacteriol.* 156, 308–315.

(23) Zeng, Y., Kulkarni, A., Yang, Z., Patil, P. B., Zhou, W., Chi, X., Van Lanen, S., and Chen, S. (2012) Biosynthesis of albomycin  $\delta_2$  provides a template for assembling siderophore and aminoacyl-tRNA synthetase inhibitor conjugates. *ACS Chem. Biol.* 7, 1565–1575.

(24) Vértesy, L., Aretz, W., Fehlhaber, H.-W., and Koger, H. (1995) Salmycin A-D, antibiotika aus *Streptomyces violaceus*, DSM 8286, mit siderophore-aminoglycosid-struktur. *Helv. Chim. Acta* 78, 46–60.

(25) Dong, L., Roosenberg, J. M., and Miller, M. J. (2002) Total synthesis of desferrisalmycin B. J. Am. Chem. Soc. 124, 15001–15005. (26) Bunet, R., Brock, A., Rexer, H.-U., and Takano, E. (2006) Identification of genes involved in siderophore transport in Streptomyces coelicolor A3(2). FEMS Microbiol. Lett. 262, 57–64.

(27) Wencewicz, T. A., Möllmann, U., Long, T. E., and Miller, M. J. (2009) Is drug release necessary for antimicrobial activity of siderophore-drug conjugates? Syntheses and biological studies of the naturally occurring salmycin "Trojan Horse" antibiotics and synthetic desferridanoxamine-antibiotic conjugates. *Biometals*, 22.

(28) Walsh, C. (2003) Antibiotics: Actions, origins, and resistance, ASM Press, Washington, D.C.

(29) Roosenberg, J. M., Jr., and Miller, M. J. (2000) Total synthesis of the siderophore danoxamine. J. Org. Chem. 65, 4833-4838.

(30) Wencewicz, T. A., Oliver, A. G., and Miller, M. J. (2012) Iron(III)-templated macrolactonization of trihydroxamate sidero-phores. *Org. Lett.* 14, 4390–4393.

(31) Jung, M. E., Yang, E. C., Vu, B. T., Kiankarimi, M., Spyrou, E., and Kaunitz, J. (1999) Glycosylation of fluoroquinolones through direct oxygenated polymethylene linkages as a sugar-mediated active transport system for antimicrobials. *J. Med. Chem.* 42, 3899–3909.

(32) Testero, S. A., Fisher, J. F., and Mobashery, S. (2010)  $\beta$ -Lactam antibiotics, In Burger's Medicinal Chemistry, Drug Discovery and Development (Abraham, D. J., and Rotella, D. P., Eds.) 7th ed., pp 259–404, Wiley, Hoboken.

(33) Mitscher, L. A. (2005) Bacterial topoisomerase inhibitors: Quinolone and pyridone antibacterial agents. *Chem. Rev.* 105, 559–592.

(34) Brochu, A., Brochu, N., Nicas, T. I., Parr, T. R., Jr., Minnick, A. A., Jr., Dolence, E. K., McKee, J. A., Miller, M. J., Lavoie, M. C., and Malouin, F. (1992) Modes of action and inhibitory activities of new siderophore- $\beta$ -lactam conjugates that use specific iron uptake pathways for entry into bacteria. *Antimcrob. Agents Chemother.* 36, 2166–2175.

(35) Hennard, C., Truong, Q. C., Desnottes, J.-F., Paris, J.-M., Moreau, N. J., and Abdallah, M. A. (2001) Synthesis and activities of pyoverdin-quinolone adducts: A prospective approach to a specific therapy against *Pseudomonas aeruginosa*. J. Med. Chem. 44, 2139–2151.

(36) Ghosh, A., Ghosh, M., Niu, C., Malouin, F., Moellmann, U., and Miller, M. J. (1996) Iron transport-mediated drug delivery using mixed-ligand siderophore- $\beta$ -lactam conjugates. *Chem. Biol.* 3, 1011–1019.

(37) McKee, J. A., Sharma, S. K., and Miller, M. J. (1991) Iron transport mediated drug delivery systems: Synthesis and antibacterial activity of spermidine- and lysine-based siderophore- $\beta$ -lactam conjugates. *Bioconjugate Chem. 2*, 281–291.

(38) Ghosh, A., and Miller, M. J. (1993) Synthesis of novel citratebased siderophores and siderophore- $\beta$ -lactam conjugates. J. Org. Chem. 58, 7652–7659.

(39) Ghosh, M., and Miller, M. J. (1994) Iron transport-mediated drug delivery: Synthesis and biological evaluation of cyanuric acid-based siderophore analogs and  $\beta$ -lactam conjugates. J. Org. Chem. 59, 1020–1026.

(40) CLSI (2009) Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 8th ed, Villanova, PA.

(41) de Lorenzo, V., Perez-Martín, J., Escolar, L., Pesole, G., and Bertoni, G. (2004) Mode of binding of the Fur protein to target DNA: Negative regulation of iron-controlled gene expression, In *Iron Transport in Bacteria* (Crosa, J. H., Mey, A. R., and Payne, S. M., Eds.) pp 185–196, ASM Press, Washington, DC.

(42) Möllmann, U., Heinisch, L., Bauernfeind, A., Köhler, T., and Ankel-Fuchs, D. (2009) Siderophores as drug delivery agents: Application of the "Trojan Horse" strategy. *Biometals* 22, 615–624.

(43) Heinisch, L., Wittmann, S., Stoiber, T., Berg, A., Ankel-Fuchs, D., and Möllmann, U. (2002) Highly antibacterial active aminoacyl penicillin conjugates with bis-catecholate siderophores based on secondary diamino acids and related compounds. *J. Med. Chem.* 45, 3032–3040.

(44) Pramanik, A., and Braun, V. (2006) Albomycin uptake via a ferric hydroxamate transport system of *Streptococcus pneumoniae* R6. *J. Bacteriol.* 188, 3878–3886.

(45) Reniere, M. L., Pishchany, G., and Skaar, E. P. (2010) Iron uptake in *Staphylococci*, In *Iron uptake and homeostasis in microorganisms* (Cornelis, P., and Andrews, S. C., Eds.) pp 247–264, Caister Academic Press, Norfolk.

(46) Wencewicz, T. A., Yang, B., Rudloff, J. R., Oliver, A. G., and Miller, M. J. (2011) N-O Chemistry for antibiotics: Discovery of N- alkyl-*N*-(pyridin-2-yl)hydroxylamine scaffolds as selective antibacterial agents using nitroso Diels-Alder and ene chemistry. *J. Med. Chem. 54*, 6843–6858.

(47) Bauer, A. W., Kirby, W. M. M., Sherris, J. C., and Turck, M. (1966) Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* 45, 493–496.

(48) Krewulak, K. D., and Vogel, H. J. (2008) Structural biology of bacterial iron uptake. *Biochim. Biophys. Acta* 1778, 1781–1804.

(49) Krewulak, K. D., Peacock, R. S., and Vogel, H. J. (2010) Periplasmic binding proteins involved in bacterial iron uptake, In *Iron uptake and homeostasis in microorganisms* (Cornelis, P., and Andrews, S. C., Eds.) pp 113–129, Caister Academic Press, Norfolk.

(50) Ji, C., and Miller, M. J. (2012) Chemical syntheses and in vitro antibacterial activity of two desferrioxamine B-ciprofloxacin conjugates with potential esterase and phosphatase triggered drug release linkers. *Bioorg. Med. Chem.* 20, 3828–3836.

(51) De Sarro, A., and De Sarro, G. (2001) Adverse reactions to fluoroquinolones: An overview on mechanistic aspects. *Curr. Med. Chem.* 8, 371–384.

(52) Kontoghiorghes, G. J., Eracleous, E., Economides, C., and Kolnagou, A. (2005) Advances in iron overload therapies. Prospects for effective use of deferipron (L1), deferoxamine, the new experimental chelators ICL670, GT56–252, L1NAII and their combinations. *Curr. Med. Chem.* 12, 2663–2681.

(53) Vertesy, L., Aretz, W., Fehlhaber, H.-W., and Ganguli, B. N. Salmycins, a process for their preparation and their use as a pharmaceutical, U.S. Patent 5475094, December 12, 1995.

(54) Vertesy, L., Werner, A., and Fehlhaber, H.-W. Chelating agents, their preparation from the antibiotics salmycin A, B, C, or D, and their use, U.S. Patent 5519123, May 21, 1996.

(55) Clifton, M. C., Corrent, C., and Strong, R. K. (2009) Siderocalins: Siderophore-binding proteins of the innate immune system. *Biometals* 22, 557–564.

(56) Barza, M. (1994) Challenges to antibiotic activity in tissues. Clin. Infect. Dis. 19, 910-915.

(57) Imbuluzqueta, E., Gamazo, C., Ariza, J., and Blanco-Prieto, M. J. (2010) Drug delivery systems for potential treatment of intracellular bacterial infections. *Front. Biosci.* 15, 397–417.

(58) Heo, J., and Hua, S. Z. (2009) An overview of recent strategies in pathogen sensing. *Sensors 9*, 4483–4502.

(59) Zheng, T., and Nolan, E. M. (2012) Siderophore-based detection of Fe(III) and microbial pathogens. *Metallomics* 4, 866–880.

(60) Murray, P. R., Baron, E. J., Pfaller, M. A., Tenover, F. C., and Yolken, R. H. (1999) *Manual of clinical microbiology*, 7th ed., American Society for Microbiology, Washington, DC.