


RSC Advances



This article can be cited before page numbers have been issued, to do this please use: L. Josa-Culleré, Y. A. Wainman, K. Brindle and F. J. Leeper, *RSC Adv.*, 2014, DOI: 10.1039/C4RA08861A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.


Diazo group as a new chemical reporter for bioorthogonal labelling of biomolecules

Laia Josa-Culleré, Yelena A. Wainman, Kevin M. Brindle and Finian J. Leeper

RSC Adv., 2014, **4**,

DOI: 10.1039/

Diazoacetyl groups undergo spontaneous cycloaddition with strained alkenes and alkynes and can be bioorthogonal reporter groups labelling proteins and glycans.



ARTICLE

Diazo group as a new chemical reporter for bioorthogonal labelling of biomolecules

Cite this: DOI: 10.1039/C4RA08861A

Laia Josa-Culleré,^a Yelena A. Wainman,^a Kevin M. Brindle^b and Finian J. Leeper^aReceived 10th January 2014,
Accepted 10th January 2014

DOI: 10.1039/C4RA08861A

www.rsc.org/

Diazoacetyl groups have been shown to undergo spontaneous cycloaddition reactions with strained alkenes and alkynes. The rates of reaction are similar to the equivalent reactions of azide groups. This allows diazo groups to be used as bioorthogonal reporter groups. This is demonstrated by fluorescent labelling of a diazoacetylated protein and of cell-surface glycans *via* metabolic incorporation of a diazoacetylated sugar.

Introduction

For *in vivo* molecular imaging, small molecules have better access to extravascular compartments and show better clearance of unbound material than macromolecules such as antibodies and lectins. Their use as imaging agents requires them to target selectively the desired biomolecule. One way to achieve this is to use the bio-orthogonal chemical reporter strategy, in which the target of interest is labelled in a two-step process. Firstly, a bio-orthogonal chemical reporter is incorporated into the target of interest using the cell's own biosynthetic machinery. This can be achieved by attaching a suitable reporter to substrates that can be metabolised by the cell. For example, some unnatural amino acids can be incorporated into proteins or monosaccharides bearing bioorthogonal functional groups can be introduced into cell-surface glycans by enzymes that tolerate the unnatural motif. Secondly, the chemical reporter is then reacted with a small molecule probe that has a complementary bioorthogonal reactive moiety. The bioorthogonal reaction partners should react rapidly under physiological conditions (37 °C, pH 6–8) and remain inert to other biological functional groups, they should form a stable and non-toxic product and have no or inert by-products. The bioorthogonal reporter strategy has proved to be a powerful tool to image glycans,^{1–4} proteins^{5–7} and lipids⁸.

The number of functional groups that have been shown to possess all the requisites to be useful chemical reporters is limited. This group comprises peptide motifs that can be tagged with small-molecule imaging probes,^{5,9} ketones and aldehydes that can be ligated with hydrazine and hydroxylamine derivatives,^{1,6,10} azides that can be modified with phosphines or reactive alkynes,^{7,8,11–13} terminal alkynes that can be tagged with azides,¹⁴ and strained alkenes^{15–20} and isonitriles²¹ that can be ligated with tetrazines. Amongst these, azides have been the

most widely used because of their chemical stability and small size.

The reaction of diazo compounds with alkenes and alkynes has been known for many years²² yet little attention has been paid to its potential as a new chemical reporter group. In this paper, we present a proof-of-principle study where we first evaluated the rate of cycloaddition reactions of a diazo compound with a strained alkene and an alkyne. We also demonstrated the bioorthogonality of the diazo group by labelling a protein and tested the incorporation of diazo-bearing sugars into cell-surface glycans.

Results and discussion

Kinetic studies with the diazo group

The highly strained (*E,E*)-1,5-cyclooctadiene (COD) **2** has been reported to perform very efficient [3+2] cycloadditions with 1,3-dipoles and inverse-electron-demand Diels Alder reactions with tetrazines, making it suitable for applications in chemical ligation.²³ In order to test whether the diazo group could be used as a chemical reporter in a two-step labelling process, its reaction with (*E,E*)-COD followed by reaction of the adduct formed with tetrazine was explored. (*E,E*)-COD was prepared from (*Z,Z*)-COD *via* a three-step phosphine oxide-mediated olefin inversion.²⁴ It was found to react cleanly with ethyl diazoacetate **1** by a [3+2] cycloaddition to give pyrazoline **3** after tautomerisation (Fig. 1). The second-order rate constant of this reaction was measured by UV spectroscopy (Fig. S3a[†]) in pentane at room temperature to be $0.045 \pm 0.002 \text{ M}^{-1} \text{ s}^{-1}$. The rate constant of the second [3+2] cycloaddition, giving **4a** and **4b**, was measured by ¹H NMR spectroscopy in CD₃OD at room temperature and is two orders of magnitude lower, which can be explained by the relief of ring strain once the first *trans* double bond of **2** has reacted. Thus pyrazoline **3** can be

ARTICLE




Fig. 1 Cycloaddition of ethyl diazoacetate with *(E,E)*-COD **2** and TMDIBO **7** and inverse-electron-demand Diels Alder with tetrazine **5**, with measured rate constants.

obtained in almost quantitative yield by reaction of equimolar amounts of **1** and **2**. The inverse-electron-demand Diels Alder reaction of the remaining *trans* double bond of **3** with tetrazine **5** to give the dihydropyridazine adduct **6** was then measured by UV spectroscopy in MeOH at room temperature and proceeded, as expected, at a much faster rate, $206 \pm 6 \text{ M}^{-1} \text{ s}^{-1}$.

Therefore, using **2** as an intermediary, two consecutive click reactions can ligate a diazo ester to a tetrazine, which could potentially bear an imaging agent. The rates of these two click reactions are of the same order of magnitude as for an azide group reacting with first **2** and then **5**, the first one being 2-fold faster and the second one 2-fold slower. However, because of the lack of solubility in water and volatility of **2**, it cannot be used as a reagent with a diazo chemical reporter in biological systems. Therefore, we used instead an alternative [3+2] cycloaddition with TMDIBO **7**, a stable dibenzocyclo-octyne developed previously by our group and used for imaging of cell-surface azido sugars.²⁵ Its reaction with ethyl diazoacetate **1** to give **8a** and **b** was followed by ^1H NMR spectroscopy in CDCl_3 at room temperature and proved to have almost exactly the same rate as for the cycloaddition of **1** with *(E,E)*-COD **2**, with a second-order rate constant of $0.043 \pm 0.004 \text{ M}^{-1} \text{ s}^{-1}$. This cycloaddition of diazo compounds with strained alkynes has only recently been reported by McGrath and Raines.²⁶

Bioorthogonal labelling of diazo-modified sugars and proteins

The bioorthogonality of the diazo group was tested by incubating ethyl diazoacetate with 10 mM glutathione at pH 7.4

and 37°C . ^1H NMR spectroscopy showed no significant reaction after 24 h (see ESI, Fig. S5[†]), indicating that diazoesters do not react with biologically relevant functional groups (thiols, carboxylic acids and amines).

In order to demonstrate the use of the diazo group as a bioorthogonal labelling agent, its ability to label proteins was tested. A solution of lysozyme was incubated with the diazoacetate NHS ester **9**²⁷ for 3 h to prepare the diazo-modified protein (Fig. 2). The protein was then purified by gel filtration, labelled with different concentrations of a TMDIBO-fluorescein complex **10** and then run on a SDS-polyacrylamide gel. Fluorescence imaging of the gel showed concentration-dependent labelling of the protein, co-localised with the protein bands as shown by Coomassie Blue staining. (see Fig. S7[†]).

We next studied the application of the diazo group for the metabolic labelling of cell-surface glycans. Tetra-*O*-acetyl-galactosamine hydrochloride ($\text{Ac}_4\text{GalNH}_3\text{Cl}$ **12**) was prepared in anomerically pure form from galactosamine **11** by the three-step process of Faroux-Corlay *et al.*,²⁸ involving protection of the amine as its imine with anisaldehyde. The diazoacetyl derivative ($\text{Ac}_4\text{GalNDiaz}$, **13**) was then synthesised using NHS ester **9** (Fig. 3A). In metabolic labelling of glycans, the sugar precursor is normally administered in *O*-peracetylated form as this increases cellular uptake; the sugars are deacetylated inside the cell by endogenous esterases.²⁹




Fig. 2. Fluorescence-labelling of diazo-modified lysozyme. The star represents the fluorophore (fluorescein)

Lewis lung carcinoma (LL2) cells were incubated with a medium containing 0 or 200 μM of the diazo sugar for 24 h. To detect diazo groups incorporated into the cell-surface glycans, the cells were washed, treated with TMDIBO-PEG-biotin (Fig. 3B) for 30 min, washed again, treated with Neutravidin-Dylight680 for 15 min, washed again, and analysed by flow cytometry. This two-step labelling procedure, labelling first with biotin and then visualising this with fluorescently labelled neutravidin gives an improved signal-to-background ratio compared to a one-step procedure using, for example, a TMDIBO-fluorophore conjugate such as **10**.³⁰ Furthermore, Neutravidin, being a large protein, is completely impermeable to the cells. Therefore any increase in fluorescence must be due to diazo groups on the cell-surface and not intracellular diazo groups.

The flow cytometry results showed a clear increase in fluorescence in cells incubated with the diazosugar compared to untreated cells. As shown in Fig. 3C, the mean fluorescence almost doubled (see ESI† for a representative histogram). This shows that some diazo groups have survived and been incorporated into cell-surface glycans. However the level of labelling is much lower than one would typically expect with the equivalent azido sugar.²⁵ The reason(s) for this are not known at present. *N*-diazoacetylgalactosamine may be a poor substrate for one or more enzymes of the galactosamine salvage

pathway or of glycan biosynthesis, even though the diazoacetyl group is smaller than the azidoacetyl group. However it is a different shape, preferring a planar conformation, and this may be detrimental to binding to some enzymes. Alternatively it may be that the *N*-diazoacetylgalactosamine is diverted into a different pathway, involving either enzymic degradation or incorporation into some intracellular entity. It may well be that other diazo-labelled sugars would be better incorporated - the enzymes required for incorporation of *N*-acetylmannosamine




Fig. 3. (A) Synthesis of tetra-*O*-acetyl-*N*-diazoacetyl-galactosamine **13**; (B) Detection of diazo-modified glycans on LL2 cancer cells surface (C) Flow cytometry results. M.F.I. = mean fluorescence intensity in the far-red channel. Error bars represent standard deviation for three repeats.

ARTICLE

into sialic acid residues of cell-surface glycans are known to be particularly tolerant of additional bioorthogonal reporter groups.³¹

Conclusions

We have been able to show that the diazo group can be used as a new chemical reporter for bioorthogonal imaging. The reaction rates with the strained *trans*-cyclooctene **2** and dibenzo-cyclooctyne **7** are of the same order of magnitude as for azides, which is a widely used chemical reporter. The labelling of lysozyme shows that diazoacetyl groups are compatible with all the functional groups present in proteins, while the labelling of cell-surface glycans shows that metabolic incorporation of diazo groups is also a possibility. The small size of the diazo group (smaller even than azide) may well make it the chemical reporter group of choice in certain instances.

Acknowledgements

We thank Will Crone and Terence Kwan for technical assistance and Henning Stöckmann for the TMDIBO-PEG-biotin. L.J.C. was in receipt of the scholarship La Caixa and Y.A.W. was recipient of Cancer Research UK funding.

Notes and references

^a University of Cambridge, Department of Chemistry, Lensfield Road, Cambridge, CB2 1EW, UK.

^b Cancer Research UK, Cambridge Institute, Li Ka Shing Centre, Robinson Way, Cambridge CB2 0RE, UK.

† Electronic Supplementary Information (ESI) available: Experimental data and NMR spectra. See DOI: 10.1039/b000000x/

- L. K. Mahal, K. J. Yarema and C. R. Bertozzi, *Science*, 1997, **276**, 1125-1128.
- E. Saxon, S. J. Luchansky, H. C. Hang, C. Yu, S. C. Lee and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2002, **124**, 14893-14902.
- S. J. Luchansky, H. C. Hang, E. Saxon, J. R. Grunwell, C. Y. Danielle, D. H. Dube and C. R. Bertozzi, *Methods Enzymol.*, 2003, **362**, 249-272.
- S. J. Luchansky, S. Goon and C. R. Bertozzi, *ChemBioChem*, 2004, **5**, 371-374.
- B. A. Griffin, S. R. Adams and R. Y. Tsien, *Science*, 1998, **281**, 269-272.
- Z. W. Zhang, B. A. C. Smith, L. Wang, A. Brock, C. Cho and P. G. Schultz, *Biochemistry*, 2003, **42**, 6735-6746.
- A. J. Link, M. K. S. Vink and D. A. Tirrell, *J. Am. Chem. Soc.*, 2004, **126**, 10598-10602.
- Y. Kho, S. C. Kim, C. Jiang, D. Barma, S. W. Kwon, J. K. Cheng, J. Jaunbergs, C. Weinbaum, F. Tamanoi, J. Falck and Y. M. Zhao, *Proc. Natl. Acad. Sci. USA*, 2004, **101**, 12479-12484.
- S. R. Adams, R. E. Campbell, L. A. Gross, B. R. Martin, G. K. Walkup, Y. Yao, J. Llopis and R. Y. Tsien, *J. Am. Chem. Soc.*, 2002, **124**, 6063-6076.
- I. Chen, M. Howarth, W. Y. Lin and A. Y. Ting, *Nat. Methods*, 2005, **2**, 99-104.
- A. E. Speers, G. C. Adam and B. F. Cravatt, *J. Am. Chem. Soc.*, 2003, **125**, 4686-4687.
- J. A. Prescher, D. H. Dube and C. R. Bertozzi, *Nature*, 2004, **430**, 873-877.
- N. J. Agard, J. A. Prescher and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2005, **127**, 11196-11196.
- A. E. Speers and B. F. Cravatt, *Chem. Biol.*, 2004, **11**, 535-546.
- R. Rossin, P. R. Verkerk, S. M. van den Bosch, R. C. M. Vulders, I. Verel, J. Lub and M. S. Robillard, *Angew. Chem. Int. Ed.*, 2010, **49**, 3375-3378.
- N. K. Devaraj, R. Weissleder and S. A. Hilderbrand, *Bioconjugate Chem.*, 2008, **19**, 2297-2299.
- D. M. Patterson, L. A. Nazarova, B. Xie, D. N. Kamber and J. A. Prescher, *J. Am. Chem. Soc.*, 2012, **134**, 18638-18643.
- C. M. Cole, J. Yang, J. Seckute and N. K. Devaraj, *ChemBioChem*, 2013, **14**, 205-208.
- J. Yang, J. Seckute, C. M. Cole and N. K. Devaraj, *Angew. Chem. Int. Ed.*, 2012, **51**, 7476-7479.
- J. Schoch, M. Wiessler and A. Jaschke, *J. Am. Chem. Soc.*, 2010, **132**, 8846-+.
- S. Stairs, A. A. Neves, H. Stockmann, Y. A. Wainman, H. Ireland-Zecchini, K. M. Brindle and F. J. Leeper, *ChemBioChem*, 2013, **14**, 1063-1067.
- R. Huisgen, in *1,3-Dipolar Cycloaddition Chemistry*, ed. A. Padwa, Wiley, New York, 1984.
- H. Stockmann, A. A. Neves, H. A. Day, S. Stairs, K. M. Brindle and F. J. Leeper, *Chem. Commun.*, 2011, **47**, 7203-7205.
- D. Boeckh, R. Huisgen and H. Noth, *J. Am. Chem. Soc.*, 1987, **109**, 1248-1249.
- H. Stockmann, A. A. Neves, S. Stairs, H. Ireland-Zecchini, K. M. Brindle and F. J. Leeper, *Chem. Sci.*, 2011, **2**, 932-936.
- N. A. McGrath and R. T. Raines, *Chem. Sci.*, 2012, **3**, 3237-3240.
- A. Ouhia, L. Rene, J. Guilhem, C. Pascard and B. Badet, *J. Org. Chem.*, 1993, **58**, 1641-1642.
- B. Faroux-Corlay, L. Clary, C. Gadras, D. Hammache, J. Greiner, C. Santaella, A. M. Aubertin, P. Vierling and J. Fantini, *Carbohydr. Res.*, 2000, **327**, 223-260.
- C. L. Jacobs, K. J. Yarema, L. K. Mahal, D. A. Nauman, N. W. Charters and C. R. Bertozzi, *Methods Enzymol.*, 2000, **327**, 260-275.
- A. A. Neves, H. Stockmann, Y. A. Wainman, J. C. H. Kuo, S. Fawcett, F. J. Leeper and K. M. Brindle, *Bioconjugate Chem.*, 2013, **24**, 934-941.
- J. Du, M. A. Meledeo, Z. Wang, H. S. Khanna, V. D. P. Paruchuri, and K. J. Yarema, *Glycobiology*, 2009, **19**, 1382-1401.