



Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

Spiroadamantyl 1,2,4-trioxolane, 1,2,4-trioxane, and 1,2,4-trioxepane pairs: Relationship between peroxide bond iron(II) reactivity, heme alkylation efficiency, and antimalarial activity

Xiaofang Wang^a, Darren J. Creek^b, Charles E. Schiaffo^c, Yuxiang Dong^a, Jacques Chollet^d, Christian Scheurer^d, Sergio Wittlin^d, Susan A. Charman^b, Patrick H. Dussault^c, James K. Wood^e, Jonathan L. Vennerstrom^{a,*}

^a College of Pharmacy, University of Nebraska Medical Center, 986025 Nebraska Medical Center, Omaha, NE, USA

^b Centre for Drug Candidate Optimisation, Monash Institute of Pharmaceutical Sciences, Monash University (Parkville Campus), 381 Royal Parade, Parkville, Victoria 3052, Australia

^c Department of Chemistry, University of Nebraska-Lincoln, Lincoln, NE, USA

^d Swiss Tropical Institute, Socinstrasse 57, CH-4002 Basel, Switzerland

^e Department of Chemistry, University of Nebraska at Omaha, Omaha, NE, USA

ARTICLE INFO

Article history:

Received 29 May 2009

Revised 20 June 2009

Accepted 2 July 2009

Available online 9 July 2009

Keywords:

1,2,4-Trioxolane

1,2,4-Trioxane

1,2,4-Trioxepane

Peroxide

Antimalarial

Artemisinin

ABSTRACT

These data suggest that iron(II) reactivity for a set of homologous spiroadamantyl 1,2,4-trioxolane, 1,2,4-trioxane, and 1,2,4-trioxepane peroxide heterocycles is a necessary, but insufficient, property of antimalarial peroxides. Heme alkylation efficiency appears to give a more accurate prediction of antimalarial activity than FeSO₄-mediated reaction rates, suggesting that antimalarial activity is not merely dependent on peroxide bond cleavage, but also on the ability of reactive intermediates to alkylate heme or other proximal targets.

© 2009 Elsevier Ltd. All rights reserved.

The semisynthetic artemisinins rapidly reduce parasite burden and are particularly effective when used in a 3-day artemisinin combination treatment (ACT) regimen.¹ One hypothesis^{2–4} that accounts for the antimalarial specificity of artemisinin⁵ is that the peroxide bond undergoes reductive activation by heme released by parasite hemoglobin digestion to produce carbon-centered free radicals or carbocations that alkylate heme⁶ or parasite proteins. Although the 1,2,4-trioxane heterocycle in artemisinin (Fig. 1) is the critical pharmacophore, its presence alone⁷ is insufficient for high antimalarial activity as most synthetic 1,2,4-trioxanes are less active than artemisinin.⁸ The discovery⁹ of ozonide (1,2,4-trioxolane) drug development candidate OZ277 (arterolane) (Fig. 1), now entering Phase III clinical trials in the form of an arterolane maleate + piperazine phosphate combination, prompted us to investigate a set of structurally related 1,2,4-trioxolanes **1**, 1,2,4-trioxanes **2**, and 1,2,4-trioxepanes **3**¹⁰ (Fig. 2) to better understand the relationship between peroxide bond iron(II) reactivity and antimalarial activity for these homologous peroxide heterocycles.

Trioxolanes **1b**¹¹ and **1c** were obtained by Griesbaum cozonolysis¹² of the corresponding oxime ether/ketone^{13,14} pairs **4/5** and **6/7** in low yields (Scheme 1). Trioxolane **1a** was similarly obtained as previously described.¹⁵

Trioxanes **2b** and **2c** were prepared as shown in Scheme 2. In the presence of catalytic *p*-toluenesulfonic acid (PTSA), **2b** was readily formed by condensation of β -hydroperoxy alcohol **8**¹⁶ with **5**. Attempts to perform the corresponding reaction of β -hydroperoxy alcohol **9**¹⁶ with **5** provided only small quantities of trioxane **2c** accompanied by a complex mixture of decomposition products.






Figure 1. Artemisinin and OZ277.

* Corresponding author. Tel.: +1 402 559 5362; fax: +1 402 559 9543.

E-mail address: jvenners@unmc.edu (J.L. Vennerstrom).

Figure 2. Trioxolanes **1**, trioxanes **2**, and trioxepanes **3**.


Scheme 1.

Characterization of the reaction products indicated that β -hydroperoxy alcohol **9** underwent acid-catalyzed O–O bond (pathway I) and C–O bond (pathway II) heterolysis followed by rearrangement. Formation of ketone **7** (pathway I) can be rationalized by a Grob-type fragmentation or by migration of the hydroxymethyl group.¹⁷ This contrasts with the phenyl migration seen in the acid-catalyzed decomposition of cumene hydroperoxide to form phenol and acetone.¹⁸ Aldehyde **11** may arise from the formation and rearrangement of a relatively stable benzylic tertiary carbocation in the transition state for fragmentation, with the co-generated hydrogen peroxide producing lactone **12** via an acid-catalyzed Baeyer–Villiger oxidation of adamantanone. Thus, acid-catalyzed heterolytic peroxide bond fragmentation (pathway II) was the dominant reaction course for **9**. Therefore, we tried protecting the hydroperoxide functional group of **9** to suppress the fragmentation.¹⁹ Treatment of **9** with *N,O*-bis-(trimethylsilyl)acetamide (BSA) in CH_2Cl_2 gave the corresponding bis(trimethylsilylated) product **10** in high yield. Subsequent condensation of **10** with **5**


in the presence of 10–20% CSA in CH_2Cl_2 afforded the desired **2c** (Scheme 2). Trioxane **2a** was prepared as previously described.²⁰

Trioxepanes **3a** and **3b** were prepared via parallel routes (Scheme 3) beginning with alcohol **13**²¹ and cinnamyl alcohol (**14**). The corresponding acetates (**15**, **16**) underwent successive Co-mediated dioxygenation²² and chemoselective reduction to furnish triethylsilyl peroxyalcohols **17** and **18**.²³ The peroxyalcohols underwent HF-mediated condensation with ketone **5** to produce **3a** and **3b**.

In vitro and in vivo antimalarial activities⁹ (Table 1) were measured using the chloroquine-resistant K1 and chloroquine-sensitive NF54 strains of *Plasmodium falciparum* and *Plasmodium berghei*-infected mice. Several observations arise from the antimalarial data. First, trioxolane **1b** was more than two orders of magnitude less potent than trioxolanes **1a** and **1c**. Because it has a α -H atom, **1b** may lack the necessary chemical stability²⁴ required for antimalarial activity. This contrasts with the good antimalarial



Scheme 3.



Scheme 2.

Table 1Activity of **1–3** against *P. falciparum* in vitro and *P. berghei* in vivo

Compd	IC ₅₀ (ng/ml) ^a		Activity ^b (%) po
	K1	NF54	
NONE	—	—	0
1a ^c	0.97	1.4	>99.99
1b	400	>1000	25
1c	2.9	3.4	99.0
2a ^d	54	44	0
2b	9.1	19	96
2c	19	22	0
3a	ND ^e	>1000	0
3b	ND ^e	>1000	ND ^e
ART ^{c,f}	1.6	2.8	98

^a Mean from $n = 2–3$ against chloroquine-resistant (K1) and chloroquine-sensitive (NF54) strains of *P. falciparum*.

^b Groups of three *P. berghei*-infected NMRI mice were treated orally one day post-infection with trioxolanes (100 mg/kg) dissolved or suspended in 3% ethanol and 7% Tween 80. Antimalarial activity was measured by percent reduction in parasitemia on day 3 post-infection. Individual measurements generally differed by less than 10%.

^c Data from Dong et al.¹⁵

^d Data from Tang et al.²⁰


^e Not determined.

^f ART = artemisinin.

activity of the corresponding and presumably more chemically stable trioxane **2b**. Trioxanes **2a** and **2c** were less potent than the corresponding trioxolanes **1a** and **1c**; neither **2a** nor **2c** had any activity in vivo. The less accessible peroxide bond in the trioxane vs trioxolane heterocycle²⁰ may explain why trioxanes **2a** and **2c** are less active than their trioxolane chemical cousins **1a** and **1c**. Similarly, the complete lack of activity of trioxepanes **3a** and **3b** may arise from peroxide bonds that are too sterically hindered (vide infra). These data suggest that antimalarial activity for these spiroadamantyl trioxolanes, trioxanes, and trioxepanes depends on a finely tuned balance between peroxide bond shielding and accessibility. To provide mechanistic insight into this hypothesis, we examined the reactivity of several of these compounds with iron(II) in the form of FeSO₄ and heme.

The reactivity of **1–3** with inorganic ferrous iron was determined using previously established standardized conditions.²⁵ In these experiments, pseudo-first order reaction rate constants for **1b**, **2a**, **2b**, **3a**, and **3b** (0.03 mM) with FeSO₄ (3 mM) in 50% acetonitrile/water at 37 °C under an argon atmosphere were determined. These data ($n = 3$) were corrected for non-specific degradation in iron-free controls. Under these experimental conditions, the previously determined pseudo-first order rate constants (k) for **1a** and artemisinin were $0.41 \pm 0.02 \text{ h}^{-126}$ and $0.054 \pm 0.006 \text{ h}^{-1}$.²⁵ The weakly active **1b** reacted considerably more rapidly ($k = 1.77 \pm 0.06 \text{ h}^{-1}$) with iron(II) than did **1a** confirming our previous observations^{26,27} that chemically reactive peroxides do not necessarily possess high antimalarial activities. Trioxane **2b** reacted with iron(II) at a rate ($k = 0.12 \pm 0.03 \text{ h}^{-1}$) between that of **1a** and artemisinin, whereas trioxepane **3b** reacted much more slowly ($k = 0.021 \pm 0.006 \text{ h}^{-1}$), and neither **2a** nor **3a** underwent statistically significant degradation over the 24 h time course studied ($k = 0.011 \pm 0.013 \text{ h}^{-1}$ and $0.017 \pm 0.012 \text{ h}^{-1}$, respectively).

As has been previously demonstrated for **1a** versus **2a**,²⁰ the difference in iron-mediated reactivity between **1a** and **3a** can be explained by considering peroxide bond accessibility of the more reactive equatorial peroxide bond conformers²⁸ as determined by molecular modeling (Spartan'08, Wavefunction, Inc.). As shown in Figure 3, the peroxide bond LUMO in **1a** is easily accessible from the cyclohexane side, whereas the peroxide bond LUMO in **3a** is

**Figure 3.** LUMO (blue) mapped on isodensity surfaces of **1a** and **3a**.

not accessible from either direction, consistent with the low reactivity of **3a** with iron(II). The greater steric crowding in **3a** is due to the methylene groups of the trioxepane ring decreasing the angle between the adamantane and cyclohexane rings. This lack of iron-mediated reactivity for trioxepane **3a** is consistent with its lack of antimalarial activity; similar arguments have been put forth to explain the low antimalarial activity of other sterically congested 1,2,4-trioxepanes.²⁹

We have previously shown³⁰ a good correlation between in vitro antimalarial activity and heme alkylation efficiency for a series of 22 trioxolanes. We measured the efficiency of heme alkylation for these trioxolanes, trioxanes and trioxepanes using these same experimental conditions.³⁰ Significant heme alkylation was observed within 30 s for the active **1c** ($67 \pm 2\%$) and moderately active **2b** ($55 \pm 4\%$), and to a slightly lesser extent for the less active **1b** ($50 \pm 1\%$) and **2a** ($45 \pm 1\%$); however the extent of heme alkylation by these compounds was far lower than previously reported for the highly active trioxolanes **1a** ($86 \pm 0.2\%$) and OZ277 ($83 \pm 2\%$).³⁰ Trioxepanes **3a** and **3b** mediated very little heme alkylation ($25 \pm 1\%$ and $1 \pm 0.4\%$), in agreement with their lack of antimalarial activity. The investigation of heme alkylation appears to give a more accurate prediction of antimalarial activity than investigation of iron-mediated reaction rates, suggesting that antimalarial activity is not merely dependant on peroxide bond cleavage, but also on the ability of reactive intermediates to alkylate heme or other proximal targets. This further supports a likely role for heme alkylation in the antimalarial mechanism of action of peroxide antimalarials.⁶

Acknowledgments

We thank Josefina Santo Tomas and Christopher Snyder for assistance in performing the in vivo antimalarial assays. This investigation received financial support from Medicines for Malaria Venture (MMV).

References and notes

- White, N. J. *Science* **2008**, 320, 330.
- Jefford, C. W. *Curr. Med. Chem.* **2001**, 8, 1803.
- Meshnick, S. R. *Int. J. Parasitol.* **2002**, 32, 1655.
- O'Neill, P. M.; Posner, G. H. *J. Med. Chem.* **2004**, 47, 2945.
- Kaiser, M.; Wittlin, S.; Nehrbass-Stuedli, A.; Dong, Y.; Wang, X.; Hemphill, A.; Matile, H.; Brun, R.; Vennerstrom, J. L. *Antimicrob. Agents Chemother.* **2007**, 51, 2991.
- Robert, A.; Benoit-Vical, F.; Claparols, C.; Meunier, B. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, 102, 13676.
- Casteel, D. A.; Peri, S. P.; Gerena, L. *Bioorg. Med. Chem. Lett.* **1993**, 3, 1707.
- Tang, Y.; Dong, Y.; Vennerstrom, J. L. *Med. Res. Rev.* **2004**, 24, 425.
- Vennerstrom, J. L.; Arbe-Barnes, S.; Brun, R.; Charman, S. A.; Chiu, F. C. K.; Chollet, J.; Dong, Y.; Dorn, A.; Hunziker, D.; Matile, H.; McIntosh, K.; Padmanilayam, M.; Santo Tomas, J.; Scheurer, C.; Scorneaux, B.; Tang, Y.; Urwyler, H.; Wittlin, S.; Charman, W. N. *Nature* **2004**, 430, 900.
- Adamantane-2-spiro-3'-5'-phenyl-1',2',4'-trioxolane (**1b**): ¹H NMR (500 MHz, CDCl₃) δ 1.70–1.93 (m, 9H), 2.03–2.16 (m, 4H), 2.22–2.26 (m, 1H), 6.09 (s, 1H), 7.36–7.46 (m, 3H), 7.50–7.56 (m, 2H). ¹³C NMR (125.7 MHz, CDCl₃) δ 26.5, 26.8, 34.2, 34.6, 34.9, 35.7, 35.8, 36.78, 36.8, 103.7, 113.0, 127.9, 128.5, 130.3, 132.8. Anal. Calcd for C₁₇H₂₀O₃: C, 74.97; H, 7.40. Found: C, 74.94; H, 7.19. Adamantane-2-spiro-3'-5'-methyl-5'-phenyl-1',2',4'-trioxolane (**1c**): mp 70–

- 72 °C. ^1H NMR (500 MHz, CDCl_3) δ 1.72 (s, 3H), 1.56–2.20 (m, 14H), 7.28–7.38 (m, 3H), 7.50–7.56 (m, 2H). ^{13}C NMR (125.7 MHz, CDCl_3) δ 25.9, 26.5, 26.9, 34.5, 34.75, 34.78, 35.3, 35.4, 36.6, 36.8, 108.7, 112.9, 125.2, 128.06, 128.11, 142.6. Anal. Calcd for $\text{C}_{18}\text{H}_{22}\text{O}_3$: C, 75.50; H, 7.74. Found: C, 75.58; H, 7.63. Adamantane-2-spiro-3'-6'-phenyl-1',2',4'-trioxane (**2b**): mp 110–111 °C. ^1H NMR (500 MHz, CDCl_3) δ 1.60–2.16 (m, 13H), 3.07 (br s, 1H), 3.83 (dd, $J_1 = 12$ Hz, $J_2 = 3.5$ Hz, 1H), 4.12 (dd, $J_1 = 12$ Hz, $J_2 = 10.5$ Hz, 1H), 5.30 (dd, $J_1 = 10.5$ Hz, $J_2 = 3.5$ Hz, 1H), 7.30–7.40 (m, 5H). ^{13}C NMR (125.7 MHz, CDCl_3) δ 27.16, 27.17, 29.4, 33.0, 33.3, 33.5, 33.7, 36.3, 37.2, 62.7, 81.3, 104.5, 127.3, 128.7, 129.1, 135.3. Anal. Calcd for $\text{C}_{18}\text{H}_{22}\text{O}_3$: C, 75.50; H, 7.74. Found: C, 75.76; H, 7.60; Adamantane-2-spiro-3'-6'-methyl-6'-phenyl-1',2',4'-trioxane (**2c**): mp 61–62 °C. ^1H NMR (500 MHz, CDCl_3) δ 1.44–2.04 (br m, 16H), 3.02 (br s 1H), 3.75 (br s 1H), 4.16 (br s 1H), 7.25–7.60 (m, 5H). ^{13}C NMR (125.7 MHz, CDCl_3) δ 21.9 (br), 25.4 (br), 27.14, 27.17 (br), 29.7 (br), 33.2 (br), 33.3, 33.5, 36.2 (br), 37.2, 65.7 (br), 79.8 (br), 104.4, 124.7 (br), 127.6 (br), 128.3, 141.6 (br). Anal. Calcd for $\text{C}_{19}\text{H}_{24}\text{O}_3$: C, 75.97; H, 8.05. Found: C, 76.16; H, 8.24; Adamantane-2-spiro-3'-7'-cyclohexyl-1',2',4'-trioxepane (**3a**): ^1H NMR (400 MHz, CDCl_3) δ 1.30–1.86 (20H), 1.89–2.02 (m, 3H), 2.02–2.15 (m, 2H), 2.40 (br s, 1H), 3.69 (dt, $J_{\text{Ha}} = 3.7$, $J_{\text{Hb}} = 3.7$, 1H), 3.69 (dt, $J = 12.4$, 3.7, 1H), 3.90 (apparent pentet, $J = 6.2$, probably dt, $J = 12.2$, 6.2, 1H). ^{13}C NMR (100.6 MHz, CDCl_3) δ 22.0, 22.4, 26.0, 27.3, 32.9, 33.6, 33.7, 33.8, 33.9, 34.2, 34.6, 35.7, 37.5, 42.3, 58.1, 81.7, 107.5. HRMS-FAB calcd for $\text{C}_{18}\text{H}_{26}\text{O}_3$ (MH^+): 293.2117; found 293.2103. Adamantane-2-spiro-3'-7'-phenyl-1',2',4'-trioxepane (**3b**): ^1H NMR (400 MHz, CDCl_3) δ 1.59–1.75 (m, 6H), 1.79–1.86 (m, 2H), 1.89–2.14 (m, 6H), 2.22–2.42 (m, 2H), 3.85 (dt, $J = 12.2$, 3.4, 1H), 4.13 (app t, $J = 11.3$, 1H), 5.14 (dd, $J = 11.5$, 3.5, 1H), 7.29–7.40 (m, 5H). ^{13}C NMR (75.5 MHz, CDCl_3) δ 27.1, 27.2, 33.1, 33.6, 33.9, 34.1, 34.7, 35.0, 37.5, 39.4, 60.0, 76.6, 86.1, 108.6, 127.3, 128.4, 128.5, 138.5. HRMS-FAB calcd for $\text{C}_{19}\text{H}_{26}\text{O}_3$ (MH^+): 301.1804; found 301.1809.
11. Keul, H. *Chem. Ber.* **1975**, *108*, 1207.
 12. Griesbaum, K.; Liu, X.; Kassiaris, A.; Scherer, M. *Liebigs Ann. Recl.* **1997**, 1381.
 13. Konakahara, T.; Matsuki, M.; Sugimoto, S.; Sato, K. *J. Chem. Soc., Perkin Trans. 1* **1987**, 1489.
 14. Dong, Y.; Vennerstrom, J. L. *J. Org. Chem.* **1998**, *63*, 8582.
 15. Dong, Y.; Chollet, J.; Matile, H.; Charman, S. A.; Chiu, F. C. K.; Charman, W. N.; Scorneaux, B.; Urwyler, H.; Santo Tomas, J.; Scheurer, C.; Snyder, C.; Dorn, A.; Wang, X.; Karle, J. M.; Tang, Y.; Wittlin, S.; Brun, R.; Vennerstrom, J. L. *J. Med. Chem.* **2005**, *48*, 4953.
 16. Ogata, Y.; Sawaki, Y.; Shimizu, H. *J. Org. Chem.* **1978**, *43*, 1760.
 17. Kagan, J.; Przybytek, J. T. *Tetrahedron* **1973**, *29*, 1163.
 18. Kharasch, M. S.; Fono, A.; Nudenberg, W. *J. Org. Chem.* **1950**, *15*, 748.
 19. Kim, H.-S.; Tsuchiya, K.; Shibata, Y.; Wataya, Y.; Ushigoe, Y.; Araki, M.; Nojima, M.; McCullough, K. J. *J. Chem. Soc., Perkin Trans. 1* **1999**, 1867.
 20. Tang, Y.; Dong, Y.; Wang, X.; Sriraghavan, K.; Wood, J. K.; Vennerstrom, J. L. *J. Org. Chem.* **2005**, *70*, 5103.
 21. Bourhani, Z.; Malkov, A. *Chem. Commun.* **2005**, 4592.
 22. Tokuyasu, T.; Kunikawa, S.; Masuyama, A.; Nojima, M. *Org. Lett.* **2002**, *4*, 3595.
 23. Jin, H.; Liu, H.; Zhang, Q.; Wu, Y. *J. Org. Chem.* **2005**, *70*, 4240.
 24. Kornblum, N.; DeLaMare, H. E. *J. Am. Chem. Soc.* **1951**, *73*, 880.
 25. Creek, D. J.; Chiu, F. C. K.; Prankerd, R. J.; Charman, W. N.; Charman, S. A. *J. Pharm. Sci.* **2005**, *94*, 1820.
 26. Creek, D. J.; Charman, W. N.; Chiu, F. C. K.; Prankerd, R. J.; McCullough, K. J.; Dong, Y.; Vennerstrom, J. L.; Charman, S. A. *J. Pharm. Sci.* **2007**, *96*, 2945.
 27. Wang, X.; Dong, Y.; Wittlin, S.; Creek, D.; Chollet, J.; Charman, S. A.; Santo Tomas, J.; Scheurer, C.; Snyder, C.; Vennerstrom, J. L. *J. Med. Chem.* **2007**, *50*, 5840.
 28. The equatorial peroxide conformer provides greater access to the peroxide bond from the cyclohexane side. For **1a** and **3a**, the energy difference between the two chair conformers ($\Delta E = E_{\text{eq}} - E_{\text{ax}}$) is 0.280 and 1.71 kcal/mol, respectively.
 29. Amewu, R.; Stachulski, A. V.; Berry, N. G.; Ward, S. A.; Davies, J.; Labat, G.; Rossignol, J.-F.; O'Neill, P. M. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 6124.
 30. Creek, D. J.; Charman, W. N.; Chiu, F. C. K.; Prankerd, R. J.; Dong, Y.; Vennerstrom, J. L.; Charman, S. A. *Antimicrob. Agents Chemother.* **2008**, *52*, 1291.