# Practical Application of Oxidation Using a Novel Na<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> System under Neutral Conditions for Scale-Up Manufacturing of 12 $\alpha$ -Hydroxy-3-oxooleanano-28,13-lactone: Key Intermediate of Endothelin A Receptor Antagonist S-0139

Takemasa Hida,<sup>†,</sup>\* Yuuki Fukui,<sup>‡</sup> Kyozo Kawata,<sup>‡</sup> Mikio Kabaki,<sup>‡</sup> Toshiaki Masui,<sup>§</sup> Masataka Fumoto,<sup>§,⊥</sup> and Hideo Nogusa<sup>‡</sup>

CMC Research Laboratories, Shanghai Office, Shionogi & Co., Ltd., 306A, 3F, Building A, Far East International Plaza, No. 319, Xianxia Road, Shanghai 200051, China, CMC Research Laboratories, Shionogi & Co., Ltd., 1-3, Kuise Terajima 2-chome, Amagasaki, Hyogo 660-0813, Japan, Nichia Pharmaceutical Industry, 224-20, Kawauchi Cho Hiraishi Ebisuno, Tokushima 771-0132, Japan, and Shionogi Discovery Research Laboratories, Shionogi & Co., Ltd., 12-4, Sagisu 5-chome, Fukushima-ku, Osaka 553-0002, Japan

### **Abstract:**

Novel alcohol oxidation using a  $Na_2WO_4$ -H<sub>2</sub>O<sub>2</sub> system was applied to manufacture 12 $\alpha$ -hydroxy-3-oxooleanano-28,13lactone which is a key intermediate of S-0139. Oxidation of the hydroxyl group of oleanolic acid was optimized based on the design of experiment (DoE) approach. Statistical analysis was used to maximize the yield of the corresponding ketone and minimize the generation of byproducts. The safety evaluation of this oxidation was also conducted in detail, and pilot manufacturing was achieved.

# Introduction

The oxidation of alcohol to ketone is one of the most important reactions in organic chemistry. Various oxidation methods have been reported, including Jones oxidation,<sup>1</sup> Swern oxidation,<sup>2</sup> Dess–Martin oxidation,<sup>3</sup> Oppeneure oxidation,<sup>4</sup> TEMPO oxidation,<sup>5,6</sup> Ca(OCl)<sub>2</sub> with acetic acid,<sup>7</sup> and TPAP oxidation.<sup>8</sup> However, oxidation is not often used in large-scale manufacturing because of inherent safety concerns and waste of byproducts.<sup>9</sup> In manufacturing a new type of endothelin A receptor antagonist S-0139, **4**, discovered at Shionogi Research Laboratories,<sup>10–12</sup> oxidation of the hydroxyl group posed a

- Bowden, K.; Heilbron, I. M.; Jones, E. R. H.; Weedon, B. C. L. J. Chem. Soc. 1946, 39.
- (2) Omura, K.; Sharma, A. K.; Swern, D. J. Org. Chem. 1976, 41, 957.
- (3) Dess, D. B.; Martin, J. C. J. Org. Chem. 1983, 48, 4155.
- (4) Djerassi, C. Org. React. 1944, 6, 207.
- (5) (a) Anelli, P. L.; Biffi, C.; Montanari, F.; Quici, S. J. Org. Chem.
- **1987**, *52*, 2562. (b) Zanka, A. *Chem. Pharm. Bull.* **2003**, *51*, 888. (6) Kartrin, K.; Brünjes, M.; Kunst, E.; Jöge, T.; Gallier, F.; Adibekian,
- A.; Kirschning, A. Adv. Synth. Catal. 2005, 347, 1423.
  (7) Ohta, T.; Zhang, H.; Torihara, Y.; Furukawa, I. Org. Process, Res. Dev. 1997, 1, 420.
- (8) Griffith, W. P.; Ley, S. V.; Whitcombe, G. P.; White, A. D. J. Chem. Soc., Chem. Commun. 1987, 1625.
- (9) Dugger, R. W.; Ragan, J. A.; Ripin, D. H. B. Org. Process Res. Dev. 2005, 9, 253.
- (10) Mihara, S.; Nakajima, S.; Matsumura, S.; Konoike, T.; Fujimoto, M. J. Pharmacol. Exp. Ther. 1994, 268, 1122.
- (11) Konoike, T.; Takahashi, K.; Araki, Y.; Horibe, I. J. Org. Chem. 1997, 62, 960.

Scheme 1. Synthetic route to S-0139 4 from oleanolic acid 1



serious issue. The synthetic route to 4 from oleanolic acid 1, of which the key step is the Barton reaction, was developed by Konoike et al.<sup>11,12</sup> The hydroxyl group at the 3-C position of **1** was oxidized by Jones oxidation, and then a stream of ozonized oxygen was introduced to obtain 12a-hydroxy-3-oxooleanano-28,13-lactone, **3**, which is a key intermediate for the Barton reaction (Scheme 1). Jones oxidation, however, is not suitable for bulk chemical development because of the toxicity of chromium(VI) and a large quantity of metal waste. Although some oxidations, for example TEMPO oxidation with NaOCl, oxidation using Ca(OCl)<sub>2</sub> and acetic acid, Swern oxidation, and other methods were screened to avoid using chromium(VI), the corresponding ketone could not be obtained. Oxidations using hypochlorite such as NaOCl and Ca(OCl)<sub>2</sub> gave the chlorinated compound at 12-en, while Swern oxidation gave methylthioether at the 3-C position as a byproduct. During the screening process, oxidation using tungstic acid (H<sub>2</sub>WO<sub>4</sub>) and hydrogen peroxide  $(H_2O_2)$  was also investigated. The corresponding ketone 2 was obtained, but many kinds of byproduct were also generated.

We previously reported a novel alcohol oxidation using a  $Na_2WO_4-H_2O_2$  system.<sup>13</sup> We applied it to the oxidation of **1** and achieved scale-up manufacturing. The reaction conditions were optimized by a statistical method (design of experiment, DoE), and a detailed safety evaluation study was conducted.

(13) Hida, T.; Nogusa, H. Tetrahedron 2009, 65, 270.

<sup>\*</sup> To whom correspondence should be addressed. Fax: 81-6-6401-1371. Telephone: 81-6-6401-8198. E-mail: takemasa.hida@shionogi.co.jp.

<sup>&</sup>lt;sup>†</sup>CMC Research Laboratories, Shanghai Office.

<sup>\*</sup> CMC Research Laboratories

<sup>§</sup> Nichia Pharmaceutical Industry.

<sup>&</sup>lt;sup>⊥</sup> Shionogi Discovery Research Laboratories.

<sup>(12)</sup> Konoike, T.; Oda, K.; Uenaka, M.; Takahashi, K. Org. Process Res. Dev. **1999**, *3*, 347.



*Figure 1.* Byproducts on oxidation of 1 using a catalytic amount of  $H_2WO_4$  or  $Na_2WO_4$  and  $H_2O_2$ .

### Scheme 2. Over-oxidation of 1



The pilot manufacturing was successful. Here we report the details of the practical oxidation for scale-up manufacturing.

## **Results and Discussion**

The reaction solvent selected was N,N-dimethylacetamide (DMA), which is the most suitable for the oxidation as described in a previous report.<sup>13</sup> Oxidation of the hydroxyl group of **1** and the formation of hydroxyl lactone **3** by ozonized oxygen gas were carried out by a telescoping process.

Preliminary Experimental Study of Oxidation. First, oxidation of 1 using H<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> was carried out in DMA at 90 °C. However, more than 2.0 equiv of H<sub>2</sub>O<sub>2</sub> was required to complete the reaction, many side reactions occurred, and the yield of 2 was only 81.1%. The main byproduct was allylic hydroperoxide 5, at a yield of 12.9%. Although some oxidation at the allylic position was reported,<sup>14</sup> allylic hydroperoxidation like this is a new type of reaction (Scheme 2). Decomposition of 5 gave other byproduct 6-8 (see Figure 1). Next, we investigated Na<sub>2</sub>WO<sub>4</sub> instead of H<sub>2</sub>WO<sub>4</sub>. Although the yield of 2 increased to 91.4% and the yield of byproduct 5 decreased to 6.9%, more than 2.0 equiv of H<sub>2</sub>O<sub>2</sub> was still required. These byproducts contaminated the isolated crystals 3; decontamination of 6 was impossible. Thus, control of 5 is the most important point for controlling the quality of 3 as well as improving the yield of **2**.

**Optimization of Oxidation Conditions (DoE).** In our previous study, we found that addition of phosphate salt accelerated the oxidation.<sup>13</sup> Without phosphate salt, excess  $H_2O_2$  is required to complete the reaction. Therefore, we sought to optimize the reaction conditions by DoE. Factors to consider included reaction temperature, concentration, the amount of  $H_2O_2$  and  $Na_2WO_4 \cdot 2H_2O$ , and the pH of the phosphate buffer. The reaction temperature was fixed at 90 °C because the reaction was slow at a lower temperature. The amount of solvent DMA was fixed at 5.4 mL vs 1.0 g of **1**, which is the minimum volume needed to dissolve **1** at 90 °C. We thus focused on the

### Table 1. Values for the response surface study (CCD)

factor	unit	values chosen
H <sub>2</sub> O <sub>2</sub> <sup><i>a</i></sup>	equiv	1.0 to 1.4
Na <sub>2</sub> WO <sub>4</sub> •2H <sub>2</sub> O	mol %	3 to 7
phosphate buffer <sup><i>b</i></sup>	pH	1.5 to 9.1

 $^a$  Using 35% aqueous solution.  $^b$  Preparation of 0.1 mol/L aqueous solution with using H\_3PO\_4, NaH\_2PO\_4  $\cdot$  2H\_2O and Na\_2HPO\_4  $\cdot$  12H\_2O.

following three factors: the amounts of  $H_2O_2$  and  $Na_2WO_4$  ·  $2H_2O$  and the pH value of the phosphate buffer. The ranges of these factors are shown in Table 1. The points for the response surface study model were chosen according to a central composite design (CCD), which comprises a factorial design with additional axial points and two center points, leading to a set of 16 experiments. The reaction time was fixed at 5 h.

Viewing the linear model for the yield of **5** as a threedimensional (3D) surface graph (Figure 2a) shows that the trend of the generation of **5** can be estimated as follows: using more  $H_2O_2$  and a phosphate buffer of lower pH increases the yield of **5**. On the other hand, viewing the quadratic model for the yield of **2** as 3D surface graphs (Figure 3), approximately 1.15-1.25 equiv of  $H_2O_2$  gives the maximum yield of **2** at any pH value of the phosphate buffer. If 1.2 equiv of  $H_2O_2$  is fixed, the 3D surface graph of **5** (Figure 2b) shows that a neutral pH value of phosphate buffer is better, and Na<sub>2</sub>WO<sub>4</sub>·2H<sub>2</sub>O at 5 mol % gives the best result.

The pH value of the Na<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> mixture is dependent on the ratio of the quantity of Na<sub>2</sub>WO<sub>4</sub>·2H<sub>2</sub>O and the pH value of the phosphate buffer. Finally, we optimized the pH value of Na<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> (Table 2). Using the Na<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> mixture at a higher pH led to a decrease in the amount of **5** and an increase in the yield of **2**. However, the Na<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> mixture under basic conditions is unstable even at 10 °C and thus is not suitable for semibatch-style and scale-up manufacturing. Therefore, the conditions of entry 4 were chosen for an actual manufacturing method. The phosphate buffer of entry 4 is used at pH 6.8.

**Safety Evaluation Study of the Oxidation.** In general, oxidation is undesirable for scale-up manufacturing because of inherent safety concerns, such as a large heat of reaction and the potential risk of explosion of the reagent.<sup>9</sup> A mixture of Na<sub>2</sub>WO<sub>4</sub>•2H<sub>2</sub>O, phosphate buffer of pH 6.8, and aqueous H<sub>2</sub>O<sub>2</sub> is also unstable to heat. However, as the reaction temperature is 90 °C, careful safety evaluation is necessary.

First, the thermal stability of the reaction mass was investigated by DSC (Table 3). The final reaction mass did not show any exothermic peak until 300 °C. The reaction mixture, which was stopped at the point of 42% conversion of the oxidation, showed an exothermic peak from 90 to 210 °C, and no other exothermic peak was detected. Thus, this peak was confirmed to be that of the normal heat of reaction. From these results, as long as the normal reaction continues, there should be no large risk. However, safety precautions should take into consideration unexpected accidents. The risk of decomposition of the mixture

<sup>(14)</sup> Konoike, T.; Araki, Y.; Kanda, Y. Tetrahedron Lett. 1999, 40, 6971.



(a)  $Na_2WO_4 \cdot 2H_2O 5 \text{ mol}\%$ Figure 2. Simulated 3D surface graphs for yield of 5.





Figure 3. Simulated 3D surface graphs for yields of 2 using phosphate buffer of different pH's.

Table 2.	Optimization	of pH	value o	of Na <sub>2</sub> W	$O_4 - H_2O$	$\mathbf{D}_{2}$
----------	--------------	-------	---------	----------------------	--------------	------------------

	nH of		yield % (HPLC peak area)			
entry	oxidant <sup>a</sup>	5	6	7	8	2
1	1.9	1.90	N.D	0.12	0.14	96.36
2	2.7	1.65	0.02	0.06	0.14	96.61
3	5.4	1.56	0.02	0.05	0.15	96.84
4	7.1	0.73	N.D	0.04	0.05	97.86
5	8.1	0.63	N.D	0.04	0.05	98.14

 $^{\it a}$  Prepared by combination of 5 mol % of  $H_2WO_4$  or  $Na_2WO_4{\cdot}2H_2O$  and 0.1 mol/L of phosphate buffer adjusted as follows: pH 1.5, 6.8, and 9.1.

of  $Na_2WO_4$ -aq  $H_2O_2$ , in the case where oxidation of 1 does not proceed for unforeseen reasons,<sup>15</sup> was investigated.

The exothermic heat of a mixture of  $Na_2WO_4$ -aq  $H_2O_2$  and DMA was measured by DSC. This mixture shows two exothermic peaks (Table 3). The first exothermic peak started from about 100 °C, which seemed to be the decomposition of  $H_2O_2$  by the tungsten catalyst because its energy value was similar to that of a mixture of  $Na_2WO_4$ -aq  $H_2O_2$  without DMA.

<sup>(15)</sup> Errors that can be anticipated include the following: stopping of agitation due to electrical power outage, error in temperature control due to problems with the thermo sensor, human error of dropwise addition speed and jacket control, stopping of the reaction or undesirable reaction due to contamination by a foreign object.

Table 3. Values of thermal screening by DSC<sup>a</sup>

			heat	energy
	onset <sup>c</sup> (°C)	peak top (°C)	J/g	kJ/mol
reaction mixture at 42% conversion <sup>b</sup>	111.41	150.76	34.97	205.7 <sup>c</sup>
final reaction mass <sup>b</sup>	N.D	N.D	N.D	N.D
DMA/water solution of	103.76	121.39	22.93	$54.6^{e}$
$Na_2WO_4$ -aq $H_2O_2^d$	264.95	275.98	61.41	$146.2^{e}$
aqueous solution of Na <sub>2</sub> WO <sub>4</sub> -H <sub>2</sub> O <sub>2</sub> <sup>d</sup>	77.27	80.81	23.53	56.0 <sup>e</sup>

<sup>*a*</sup> Heating rate; 10 °C/min. <sup>*b*</sup> Concentration of charged 1: 0.17 mmol/g. <sup>*c*</sup> Based on charged 1. From this value and conversion of this sample, total heat of reaction energy is calculated to be 354.7 kJ/mol. <sup>*d*</sup> Na<sub>2</sub>WO<sub>4</sub>•2H<sub>2</sub>O: 1.74 × 10<sup>2</sup> mmol/g, H<sub>2</sub>O<sub>2</sub>: 0.42 mmol/g. <sup>*e*</sup> Based on charged H<sub>2</sub>O<sub>2</sub>.



Figure 4. RC1e heat flow of oxidation of 1.

### Table 4. Result of RC1e experiment

operation	$\Delta H_{\rm react}$ (kJ/mol) <sup>b</sup>	specific heat of reaction mass (J/(g•K))	$\Delta T_{ad}$ (K)
dosing <sup>a</sup>	315.1	2.19	53.7
stirring	35.0		6.0

<sup>*a*</sup> Dropwise addition of the mixture of  $Na_2WO_4 \cdot 2H_2O$  (5 mol %), 0.05 mol/L of aq  $Na_2HPO_4 \cdot 12H_2O$  (2 mol %), 0.05 mol/L of aq  $NaH_2PO_4 \cdot 2H_2O$  (2 mol %) and 35% aq  $H_2O_2$  (1.2 equiv) for 1 h. <sup>*b*</sup> Based on charged **1**.

Another peak of unknown exothermic energy was observed near 200 °C. It shows that inadequate reaction control due to operation error or decomposition of residual oxidant may cause a runaway reaction. We next investigated whether the adiabatic temperature rise by the primary exothermic decomposition or heat of reaction can reach the secondary decomposition temperature.

The reaction calories were measured by RC1e as shown in Figure 4 and Table 4. The mixture of Na<sub>2</sub>WO<sub>4</sub>-aq H<sub>2</sub>O<sub>2</sub> containing 0.1 mol/L of pH 6.8 phosphate buffer, was added dropwise over 1 h. The total reaction calorie level was 350.1 kJ/mol, and the adiabatic temperature rise ( $\Delta T_{ad1}$ ) was 59.7 K. The process temperature is 90 °C, and the maximum temperature of the synthetic reaction (MTSR) by this  $\Delta T_{ad1}$  is 149.7 °C. The MTSR is close to the starting point of the secondary exothermic peak of the mixture of Na<sub>2</sub>WO<sub>4</sub>-aq H<sub>2</sub>O<sub>2</sub> and DMA. If the adiabatic temperature rise, due to this secondary exothermic decomposition ( $\Delta T_{ad2} = 28.0$  K), is added to the MTSR, the inside temperature rises to 177.7 °C, which is higher than the boiling temperature of DMA (Figure 5).

Adiabatic temperature rise by the primary exothermic decomposition of the mixture of  $Na_2WO_4$ -aq  $H_2O_2$  and DMA is only 10.5 °C, and the risk of a runaway reaction is low.



*Figure 5.* Simulation of an unexpected scenario of oxidation of 1.

**Table 5.** MTSR at several  $T_{\rm P}$ 's and dropwise addition times

$T_{\rm P}$ (°C)	dropwise addition time (min)	X <sub>ST</sub> (%)	MTSR (°C) <sup>a</sup>
50	10	2	108.0
90	10	25	134.4
90	60	79	102.4

<sup>*a*</sup> MTSR =  $T_p - (X_{ST})/100$  ×  $\Delta T_{ad}$  ×  $(M_F)/(M_{ST})$ .  $T_P$  = process temperature,  $X_{ST}$  = conversion at dropwise addition of stoichiometric reagent,  $M_{ST}$  = mass of reaction mixture at dropwise addition of stoichiometric reagent,  $M_F$  = mass of reaction mixture at finish of dropwise addition,  $\Delta T_{ad}$  = adiabatic temperature rise of reaction heat.



*Figure 6.* Comparison between simulation and actual pilotplant manufacturing.

Therefore, control of the reaction heat is the key point for ensuring safety. We calculated the MTSR at several process temperatures and from the dropwise addition time of a mixture of Na<sub>2</sub>WO<sub>4</sub>-aq H<sub>2</sub>O<sub>2</sub> including pH 6.8 of phosphate buffer (Table 5). These findings indicated that safety could be ensured if the mixture of Na<sub>2</sub>WO<sub>4</sub>-aq H<sub>2</sub>O<sub>2</sub> is added dropwise over 1 h at 90 °C. At a reaction temperature of 50 °C, reaction for more than 24 h is necessary to complete the oxidation.

Simulation and Actual Pilot-Plant Manufacturing for Oxidation of 1. Pilot-plant manufacturing of the oxidation of 1 was carried out using a 300 L glass-lined reactor. The heat-transfer jacket system contains ethylene glycol—water. The reactor was charged with 22 kg of 1 and 107 kg of DMA and heated to 90 °C, followed by stirring. A mixture of Na<sub>2</sub>WO<sub>4</sub>–aq H<sub>2</sub>O<sub>2</sub> including phosphate buffer of pH 6.8 was added dropwise for about 1 h at the same temperature. The process was simulated and compared with the experimental result (Figure 6). It shows that the profile predicted for the reaction temperature is similar to the profile of the actual temperature, leading us to conclude that this oxidation can be safely scaled up.



Figure 7. 12-Oxolactone 10.

Scheme 3. Synthetic process of 3 from 2



Process of Manufacturing 3. The compound isolated in this step was not 2 but hydroxy lactone 3. After the oxidation of 1, toluene was used to extract 2, followed by its condensation, and then DMA and methanol were added as reaction solvents. The mixture was ozonized to obtain 3. However, epoxide compound 9 was obtained under these conditions (Scheme 3). According to a previous report using dichloromethane and methanol as a reaction solvent,<sup>11</sup> **3** was directly obtained with a small amount (6-7%) of 12-oxolactone 10 (Figure 7) under a stream of ozonized oxygen gas. DMA seems to include a small amount of amine, and the reaction mixture shows weakly basic conditions. Therefore, the carboxylic acid at the 17-C position in 2 is masked by amine, and the opening of epoxide seems to be difficult to do. Although 3 was oxidized to 10 in a stream of ozonized oxygen gas,<sup>16</sup> 9 remained in the course of the reaction and did not give 10 under our conditions. The quality of 3 was improved. Epoxide 9 dissolves in DMA/ methanol cosolvent at room temperature, while hydroxy lactone 3 was difficult to dissolve. Epoxide 9 was opened by addition of sulfuric acid, and crystals of 3 were obtained by this ringopening reaction.

Steric Conformation of Epoxide of 9 and Reaction Mechanism. Coupling between 11-H and 12-H showed that a conformation of epoxide of 9 is  $\alpha$ -face. The  $\beta$ -Face of 12-en in 2 seems to be sterically hindered by carboxylic acid at the 17-C position, and ozone cannot attack from the  $\beta$ -face (Figure 8). In general, ozonolysis causes a cleavage reaction of olefins via primary oxonide 11;<sup>17</sup> however, epoxidation proceeded in this case. We think that the E-ring of 2 arches out into an  $\alpha$ -face,<sup>18</sup> and primary ozonide 11<sup>17</sup> cannot be formed by this steric hindrance. Therefore, epoxide 9 was produced in this case (Figure 8).



Figure 8. Mechanism for  $\alpha$ -epoxidation of 2 by ozone.

# Conclusion

A novel alcohol oxidation system using  $Na_2WO_4-H_2O_2$ under neutral conditions was applied to pilot manufacturing of  $12\alpha$ -hydroxy-3-oxooleanano-28,13-lactone, which is a key intermediate of S-0139. The reaction conditions were optimized by DoE, the safety evaluation was conducted in detail, and then scale-up manufacturing was achieved on a scale of 22 kg.

### **Experimental Section**

NMR spectra were measured on a Varian<sup>Unity</sup> Inova-500 or Inova-600. High-resolution mass spectra were recorded on JEOL JMS-SX/SX102A. High performance liquid chromatographic (HPLC) analysis was carried out using a Shimadzu LC-2010HT. DoE was designed and analyzed by Design Expert ver. 7. DSC were measured on METTLER TOLEDO DSC822e.

**Procedure for Manufacturing 3 in a Pilot Plant.** Na<sub>2</sub>HPO<sub>4</sub>•12H<sub>2</sub>O (0.34 kg, 0.95 mol) and NaH<sub>2</sub>PO<sub>4</sub>•2H<sub>2</sub>O (0.15 kg, 0.95 mol) were dissolved in water (18.9 kg). Na<sub>2</sub>WO<sub>4</sub>·2H<sub>2</sub>O (0.79 kg, 2.40 mol) was dissolved in cooled 35% aq H<sub>2</sub>O<sub>2</sub> (5.6 kg, 57.65 mol) (Caution! Exothermic dissolution. Keep the temperature below 20  $^{\circ}C!$ ) and mixed with a previously prepared phosphate buffer. Oleanolic acid 1 (22.0 kg, 48.17 mol) and DMA (107 kg, 114 L) were added to a glass-lined reactor (300 L) with heating to 90 °C. The cooled mixture of Na<sub>2</sub>WO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> with phosphete buffer was added dropwise to the DMA solution of **1** for over 1 h followed by stirring for 1 h under the same reaction conditions. After termination of the reaction was confirmed by HPLC, the reaction mixture was transferred to a 500 L reactor, and the reaction temperature was cooled to 45 °C. Toluene (96 kg, 110 L) and 110 kg of 5% aq NaCl including L-ascorbic acid (1.69 kg) were added to deactivate excess H<sub>2</sub>O<sub>2</sub> at 45 °C followed by stirring for 5 min. After checking for disappearance of peroxide using KI starch paper, the organic layer was separated at the same temperature. The organic layer was condensed to ca. 31 kg under reduced pressure. DMA (157 kg, 167 L) and methanol (35 kg, 44 L) were added to the residue, and then the mixture was cooled to -40 °C and stirred under a stream of ozonized oxygen gas over 2.5 h. After termination of the reaction had been confirmed by HPLC, excess ozone was removed by bubbling with a N2 stream, and a mixture of L-ascorbic acid (4.40 kg) and DMA (21 kg, 22 L) was added to deactivate peroxide. After checking for disappearance of peroxide using KI starch paper, the reaction mixture was heated to 30 °C. Next, 62% sulfuric acid (3.80 kg, 24.09 mol) was added dropwise over 5 min to obtain crystals of 3. White crystals of **3** were separated with a centrifuge and washed with a mixture

<sup>(16)</sup> Waters, W. L.; Rollin, A. J.; Bardwell, C. M.; Schneider, J. A.; Aanerud, T. W. J. Org. Chem. 1976, 41, 889.

<sup>(17)</sup> Criegee, R.; Schröder, G. Chem. Ber. 1960, 93, 689.

<sup>(18) (</sup>a) Barton, D. H. R.; Holness, N. J. J. Chem. Soc. 1952, 78. (b) Abel el, A. M.; Carlisle, C. H. Chem. Ind. (London) 1954, 279.

of water and methanol (79 kg, water 44 kg, and methanol 35 kg). The collected crystals were dried by double-corn dryer under reduced pressure at 45 °C. By this procedure, 17.4 kg of **3** was obtained (isolated yield: 76.8%).<sup>11</sup>

Preparation of 11α-Hydroperoxy-3-keto-oleanolic Acid 5.  $H_2WO_4$  (33.3 mg, 0.14 mmol) was dissolved in 35% aq  $H_2O_2$ (7.86 g, 80.10 mmol) by stirring for 40 min at room temperature. Oleanolic acid 1 (6.10 g, 13.35 mmol) and DMA (37 mL) were added to a flask (100 mL) which was heated to 90 °C. A mixture of H<sub>2</sub>WO<sub>4</sub> and H<sub>2</sub>O<sub>2</sub> was added dropwise over 10 min at 90 °C followed by stirring for 2 h. Generation of 5 was confirmed by HPLC (yield; 17.2 peak area %). The reaction mixture was cooled to 60 °C and toluene (50 mL) was added. Excess H<sub>2</sub>O<sub>2</sub> was deactivated by 40% aq glyoxylic acid (13.5 g, 66.75 mmol). The organic layer was separated and washed with 5% aq NaCl (25 g). The organic layer was evaporated to  $\sim 10$  g, and methanol (18 mL) was added to the residue. A small amount of seed crystals of 3-oxoolean-12-en-28-oic acid 2 was added to crystallize 3. After filtering 3, the mother liquid was collected and concentrated. This residue was purified by silica gel chromatography (eluent: dichloromethane/ethyl acetate = 7:2), and 38 mg of 5 was obtained. Compound 5: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.85 (s, 3H), 0.93 (s, 3H), 0.96 (s, 3H), 1.05 (s, 3H), 1.10 (s, 3H), 1.15 (s, 3H), 1.17 (m, 1H), 1.25 (s, 3H), 1.25 (m, 1H), 1.26 (m, 1H), 1.31 (m, 1H), 1.39 (m, 2H), 1.49 (m, 1H), 1.54 (m, 1H), 1.58 (m, 1H), 1.62 (m, 1H), 1.66 (m, 1H), 1.67 (m, 2H), 1.68 (m, 1H), 1.79 (td, J = 13.8, 4.3 Hz, 1H), 1.94 (d, J = 8.9 Hz, 1H), 2.02 (td, J = 13.1, 3.7 Hz, 1H), 2.29 (td, J = 14.0, 7.1, 3.6 Hz, 1H), 2.40 (ddd, J = 16.0, 6.7, 3.6 Hz, 1H), 2.57 (ddd, J = 16.0, 11.2, 7.4 Hz, 1H), 2.93 (dd, J = 14.1, 4.2 Hz, 1H), 4.57 (dd, J = 8.9, 3.7 Hz, 1H), 5.56 (d, J = 3.7 Hz, 1H). <sup>13</sup>C NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  16.3, 18.8, 19.8, 21.7, 22.9, 23.7, 24.9, 26.6, 28.0, 30.9, 32.3, 32.8, 33.2, 33.9, 34.4, 37.8, 40.0, 40.9, 42.2, 43.0, 45.8, 46.4, 47.9, 49.2, 55.5, 81.6, 121.9, 151.4, 183.3, 217.6. HRMS (FAB<sup>+</sup>) Calcd for  $C_{30}H_{47}O_5$ : ([M + H]<sup>+</sup>) 487.3418. Found: m/z 487.3064.

**Compounds 6–8.** Oxidation of **1** was carried out as follows based on the section, Preparation of  $11\alpha$ -hydroperoxy-3-ketooleanolic acid **5**. After the oxidation, toluene was added, and 10% aq Na<sub>2</sub>SO<sub>3</sub> was added dropwise to reduce the excess H<sub>2</sub>O<sub>2</sub>. By this reduction, **5** was decomposed, and byproducts **6–8** were generated. The organic layer was separated, washed with 5% aq NaCl, and evaporated. The structures of of **6–8** were identified by LC–MS and LC–NMR.

Compound 6: <sup>1</sup>H NMR (600 MHz, D<sub>2</sub>O/CD<sub>3</sub>CN = 25:75, 25 °C)  $\delta$  0.88 (s, 3H), 0.96 (s, 3H), 1.01 (s, 3H), 1.02 (s, 3H), 1.04 (s, 3H), 1.05 (s, 3H), 1.08 (s, 3H), 1.33 (br d,  $J = \sim$ 13 Hz), 1.45 (m, 1H), 1.86 (t, J = 13.6 Hz, 1H), 2.09 (m, 1H), 2.14 (br d,  $J = \sim$ 14 Hz), 2.39 (ddd, J = 15.9, 7.1, 3.4 Hz, 1H), 2.645 (ddd, J = 15.9, 7.1, 3.4 Hz, 1H), 5.49 (dd, J = 10.2, 2.9 Hz, 1H), 6.13 (d, J = 10.2 Hz). <sup>13</sup>C NMR (600 MHz, D<sub>2</sub>O/CD<sub>3</sub>CN = 25:75, 25 °C)  $\delta$  23.0, 25.2, 30.3, 31.3, 32.5, 41.4, 36.3, 36.9, 38.5, 52.3, 54.0, 90.8, 219.4. LC-MS m/z = 453.4 [M + H]<sup>+</sup>. Compound 7: <sup>1</sup>H NMR (600 MHz, D<sub>2</sub>O: CD<sub>3</sub>CN = 25:75, 25 °C) 0.92 (s, 3H), 0.93 (s, 3H), 0.95 (s, 3H), 0.95 (s).

3H), 1.02 (s, 3H), 1.05 (s, 3H), 1.17 (s, 3H), 1.38 (s, 3H), 1.52 (m, 1H), 2.37 (ddd, J = 15.7, 6.9, 4.7 Hz, 1H), 2.55 (ddd, J =15.7, 10.2, 7.2 Hz, 1H), 2.76 (ddd, J = 13.9, 7.2, 4.7 Hz, 1H), 2.96 (dd, J = 13.3, 4.0 Hz, 1H), 5.58 (s, 1H, O=C-CH=C). <sup>13</sup>C NMR (600 MHz,  $D_2O/CD_3CN = 25.75$ , 25 °C)  $\delta$  15.5, 18.6, 20.6, 22.8, 26.0, 27.6, 30.3, 31.9, 32.0, 33.3, 36.7, 39.3, 42.0, 44.0, 44.4, 47.6, 54.4, 171.9, 201.4, 219.9. LC-MS m/z  $= 469.4 ([M + H]^{+})$ . Compound 8: <sup>1</sup>H NMR (600 MHz, D<sub>2</sub>O/  $CD_3CN = 25.75, 25 \ ^{\circ}C) \delta \ 0.92 \ (s, 3H), 0.93 \ (s, 3H), 0.95 \ (s, 3$ 3H), 1.02 (s, 3H), 1.05 (s, 3H), 1.17 (s, 3H), 1.38 (s, 3H), 1.52 (m, 1H), 2.37 (ddd, J = 15.7, 6.9, 4.7 Hz, 1H), 2.55 (ddd, J =15.7, 10.2, 7.2 Hz, 1H), 2.76 (ddd, *J* = 13.9, 7.2, 4.7 Hz, 1H), 2.96 (dd, J = 13.3, 4.0 Hz, 1H), 5.58 (s, 1H). <sup>13</sup>C NMR (600 MHz,  $D_2O/CD_3CN = 25:75, 25 \ ^{\circ}C) \delta 15.5, 18.6, 20.6, 22.8,$ 26.0, 27.6, 30.3, 31.9, 32.0, 33.3, 36.7, 39.3, 42.0, 44.0, 44.4, 47.6, 54.4, 171.9, 201.4, 219.9. LC-MS m/z = 469.6 ([M +  $H^{+}).$ 

Preparation of 12α-Epoxy-3-keto-oleanolic Acid 9. Compound 9 was crystallized from the ozonized reaction mixture of 2 at -40 °C, as described in Procedure for Manufacturing 3 in a Pilot Plant. The crystals of 9 were separated by filtration at low temperature. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.82 (s, 3H), 0.88 (s, 3H), 0.95 (s, 3H), 0.98, (s, 3H), 0.99 (s, 3H), 1.07 (s, 3H), 1.14 (s, 3H), 1.22 (m, 1H), 1.25 (m, 1H), 1.33 (m, 1H), 1.36 (m, 1H), 1.39 (m, 1H), 1.43 (m, 1H), 1.40 (m, 1H), 1.45 (m, 1H), 1.47 (m, 1H), 1.49 (m, 1H), 1.65 (m, 1H), 1.67 (m, 1H), 1.69 (m, 2H), 1.78 (m, 2H), 1.91 (ddd, *J* = 13.3, 7.2, 5.1 Hz), 1.98 (m, 1H), 2.00 (m, 1H), 2.01 (m, 1H), 2.44 (ddd, J = 16.2, 8.7, 7.2 Hz, 1H), 2.47 (ddd, J = 16.2, 8.3, 5.1 Hz, 1H), 3.20 (t, J = 1.8 Hz, 1H). <sup>13</sup>C NMR (200 MHz, CDCl<sub>3</sub>, 5 °C) δ 16.8, 19.3, 20.0, 21.3, 22.4, 22.6, 23.3, 24.2, 27.3, 29.2, 30.4, 32.6, 33.1, 33.2, 33.9, 34.0, 36.2, 37.9, 39.0, 39.2, 40.3, 40.5, 42.7, 47.0, 47.6, 54.4, 63.6, 67.0, 183.1, 218.3. HRMS  $(FAB^+)$  Calcd for C<sub>30</sub>H<sub>47</sub>O<sub>4</sub>:  $([M + H]^+)$  471.3469. Found: m/z471.000.

### Acknowledgment

We thank Dr. Junko Kikuchi in Shionogi Discovery Research Laboratories for NMR analysis of compounds 5-9, Dr. Sohei Omura in Shionogi CMC Research Laboratories and Mr. Takahiro Hasegawa in Shionogi Biostatistics Department for helpful discussions on statistical science, and Dr. Yusuke Sato in Shionogi CMC Research Laboratories for helpful discussions on safety evaluation study.

# **Supporting Information Available**

Copies of <sup>1</sup>H NMR and C <sup>13</sup>NMR spectra of compound **5** and **9**; copies of LC–NMR of **6–8**; tables showing individual runs, details of statistical analysis (DoE); copies of DSC charts of a mixture of  $Na_2WO_4$ – $H_2O_2$  and reaction mass. This material is available free of charge via the Internet at http://pubs.acs.org.

Received for review October 16, 2009.

OP900265H