This article was downloaded by: [Pennsylvania State University] On: 06 June 2014, At: 07:00 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

New Approaches for the Synthesis of 1,3,4-Thiadiazole and 1,2,4-Triazole Derivatives with Antimicrobial Activity

Wagnat W. Wardakhan ^a & Nahed N. E. El-Sayed ^a ^a National Organization for Drug Control and Research (NODCAR), Cairo, Egypt Published online: 18 Feb 2009.

To cite this article: Wagnat W. Wardakhan & Nahed N. E. El-Sayed (2009) New Approaches for the Synthesis of 1,3,4-Thiadiazole and 1,2,4-Triazole Derivatives with Antimicrobial Activity, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:3, 790-804, DOI: <u>10.1080/10426500802274534</u>

To link to this article: http://dx.doi.org/10.1080/10426500802274534

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



New Approaches for the Synthesis of 1,3,4-Thiadiazole and 1,2,4-Triazole Derivatives with Antimicrobial Activity

Wagnat W. Wardakhan and Nahed N. E. El-Sayed National Organization for Drug Control and Research (NODCAR), Cairo, Egypt

The thiosemicarbazide derivatives 3a and 3b were cyclized in the presence of concentrated sulfuric acid to give the 5-cyanomethyl-1,3,4-thiadiazole derivatives 4a and 4b, respectively. The latter products were used for many heterocyclic transformations to form coumarin, 1,3,4-thiadiazolo[4,5-a]pyridine, and 5-thiophenylthiophene. In addition, compound 3b underwent cyclization in NaOH (2 N) solution to give the 1,2,4-triazole derivative 16. The reactivity of the latter product towards some chemical reagents was studied. The antimicrobial activities of the newly synthesized products were measured and showed high activities.

Keywords Antimicrobial; coumarin; thiadiazole; triazole

INTRODUCTION

The cyclization of suitable linear compounds is one of the most common and popular methods for preparing heterocyclic compounds. Asymmetrical ureas have been cyclized to produce several heterocycles such as 1,3,4-thiadiazoles, 1,2,4-triazoles, and 1,3,5triazines. 2,4-Disubstituted semicarbazones have been proposed as dipeptide isoesters² and could be a new class of urea peptide mimetics. The possible biological properties of semi- and thiosemicarbazone derivatives make it attractive to study the chemical reactivity of these compounds. For instance, Kabashima et al. demonstrated that the reaction of α -halocarbonyl compounds with alkyland arylidenephenylthiosemicarbazones gave 1,2,4-triazoline and

Received 13 May 2008; accepted 16 June 2008.

Address correspondence to Wagnat W. Wardakhan, National Organization for Drug Control and Research (NODCAR) P.O. Box 29, Cairo, A. R. Egypt. E-mail: wagnatward@ hotmail.com

1,2,4-dithiazolidines.³ 2,4-Disubstituted thiosemicarbazides were cyclized to 1,2,4-triazoline-3-thiones and 1,3,4-thiadiazolines when treated with acyl isothiocyanates.⁴ Oxidative cyclization of substituted aldehyde thiosemicarbazones, induced by different metallic salts, led to 1,2,4-triazoline derivatives.⁵⁻⁹ Alternatively, the interaction of thiosemicarbazide and dithiocarbazate derivatives with some π -acceptors such as propanedinitrile and benzoquinone, as well as naphthoquinone, afforded thiazines, thiadiazines, thiadiazoles, indazoles, pyridazines, oxathiadiazoles, and various fused heterocyclic compounds possible via a single electron transfer before the ring closure step.¹⁰⁻¹⁴

RESULTS AND DISCUSION

In this work, we use the thiosemicarbazide derivatives **3a** and **3b**, which were synthesized according to the literature procedure^{15,16} in the synthesis of potentially biologically active 1,3,4-thiadiazole and 1,2,4-triazole derivatives. Therefore, compounds **3a** and **3b** underwent cyclization in the presence of concentrated sulfuric acid to give the 5-cyanomethyl-1,3,4-thiadiazole derivatives **4a** and **4b**, respectively. The assignment of the structures of compounds **4a,b** was based on analytical and spectroscopic data. Thus, the ¹H NMR spectrum of compound **4a** showed the presence of a singlet at δ 4.68 ppm for the CH₂ group, a multiplet at δ 7.26–7.33 ppm corresponding to the phenyl protons, and a singlet at δ 8.69 ppm for the NH group.

The reaction of compounds **4a** or **4b** with benzenediazonium chloride (**5**) in ethanol/sodium hydroxide solution at $0-5^{\circ}$ C gave the phenyl hydrazone derivatives **6a** and **6b**, respectively. The reaction of compounds **4a** and **4b** with benzaldehyde (**7**) gave the benzal derivatives **8a** and **8b**, respectively. In addition, the reaction of compounds **4a** and **4b** with salicylaldehyde (**9**) gave the coumarin derivatives **10a** and **10b**, respectively. Formation of coumarins from the reaction of cyanomethylene reagents with salicylaldehyde has been reported previously in the literature.^{17,18} The analytical and spectroscopic data of **8a,b** and **10a,b** were in agreement with the proposed structures. Thus the ¹H NMR spectrum **10a** a singlet at δ 6.69 ppm corresponding to coumarin H-4, a multiplet at δ 7.31–7.44 ppm for aromatic protons and a singlet at δ 8.35 ppm (D₂O exchangeable) for an NH group (Scheme 1).



SCHEME 1

The reaction of compounds **4a** and **4b** with either malononitrile (**11a**) or ethyl cyanoacetate (**11b**) afforded the 1,3,4-thiadiazolo[4,5-a]pyridine derivatives **12a-d**, respectively. The structures of the latter products were established based on analytical and spectroscopic data. On the other hand, the reaction of compound **4a** and **4b** with

the cinnamonitrile derivatives **13a** and **13b** resulted in the formation of the 7-phenyl-1,3,4-thiadiazolo[4,5-a]pyridine derivatives **14a–d**, respectively. The obtained analytical and spectral data are in agreement with the proposed structures. Thus, the ¹H NMR spectrum of **14a** showed the presence of a singlet at δ 4.68 ppm corresponding to NH₂ group, a singlet at δ 5.36 for pyridine H-4, a multiplet at δ 7.29–7.38 ppm for aromatic protons, and a singlet at δ 8.36 (D₂O exchangeable) corresponding to an NH group (Scheme 2).



SCHEME 2

The reaction of either compound 4a or 4b with any of the cyanomethyleno reagents 11a or 11b and elemental sulfur gave the 5-thiophenylthiophene derivatives 15a-d respectively. The reaction was produced in the same way as the reported in Gewald's thiophene synthesis.^{19,20}

Compound **3b** underwent ready cyclization in sodium hydroxide (2.0 N) solution to give the 1,2,4-triazole derivative **16**. Assignment of the structure of the latter product was based on analytical and spectroscopic data. The ¹H NMR spectrum of compound **16** showed the presence of a singlet at δ 4.82 corresponding to the CH₂ group, a broad singlet at δ 5.26 for the SH group, and a multiplet at δ 7.30–7.37 for the phenyl protons. Further confirmation for the structure of compound **16** was obtained through studying its reactivity with some chemical reagents. Thus when compound **16** was allowed to react with hydrazine hydrate (**17a**) and phenylhydrazine (**17b**), the hydrazone derivatives **18a** and **18b**, respectively, were formed. Moreover, the reaction of compound **16** with benzenediazonium chloride **5** gave the phenylhydrazone derivative **19**. The analytical and spectroscopic data of compounds **18a,b** and **19** were consistent with the proposed structures (see the Experimental section and Scheme 3).

The reaction of compound 16 with benzaldehyde gave the benzal derivative 20. Furthermore, the reaction of compound 16 with salicylaldehyde (9) gave the 3-(1,2,4-triazol-3-yl)-coumarin derivative 21 (Scheme 4).

ANTIMICROBIAL ACTIVITY

The in vitro antimicrobial activity of the structurally promising 24 heterocyclic derivatives were tested against two strains of Gram-positive bacteria (*Bacillus subtilis* CECT 498 and *Bacillus cereus* CECT 148), two strains of Gram-negative bacteria (*Escherichia coli* ECT 101 and *Pseudomonas aeruginosa*), and*Candida albicans* CECT 1394 as a representative species of fungi. The newly synthesized products were dissolved in aqueous ethanol to give a logarithmic series of concentrations from 2 to 256 mg/L upon tenfold dilution with the growth medium and spore suspension of the test fungi. The toxicity of compounds was determined via pipette additions into the wells of multi-well slides, followed by with 25 μ L of the culture medium. The inoculated slides were then incubated at 25 °C until short germ tubes appeared of approximately 50 μ m in length (at 0 h).

Five μ L volumes of the prepared compound test solutions were added to the inoculated wells, one control well on each slide being treated



SCHEME 3

with solvent only. The slides were then returned to the incubator until germ tubes 400 μ m (±50 μ m) long were visible in the control wells. Further growth was arrested by the addition of lactophenol aniline blue to each of the wells. The minimal inhibitory concentration [(MIC)



in μ g mL⁻¹] was determined using an adaptation of agar streak dilution method based on radial diffusion.^{21,22} Under the same conditions, ampicillin (antibacterial) and cycloheximide (antifungal) were used as standards. The MIC was considered to be the lowest concentration of the tested compound (in dimethylformamide) that inhibits growth of bacteria or fungi on the plate (Table I). The diameters of the inhibition zones corresponding to the MICs are also presented in Table I.

It is obvious from Table I that compounds **10b** (N(5-oxo-2Hchromen-3-yl)-1,3,4-thiadiazol-2-yl)benzamide), **12d** (N-(7-amino-8cyano-5-oxo-5H-[1,3,4-]thiadiazolo[3,2-a]pyridine-2-yl)benzamide), and **15c** (N-(5-(3,5-diamino-4-cyanothiophene-2-yl)-1,3,4-thiadiazol-2yl)benzamide) are the most active towards *E. coli*, and their activity is higher than that of ampicilin. Alternatively, compounds **4a**, **4b**, **8b**, **12b**, **12d**, and **16** are more active towards *B. Cereus* than ampicilin. Compounds **4a**, **8b**, **12b**, **12d**, and **15a** showed high activity against *B. Subtilis*. Moreover, compounds **6a**, **6b**, **8a**, **10a**, **10b**, **14b**, **15a**, and **15c** showed high activity towards *C. Albicans*, where compounds **10b** and **15b** (both of them with the 2-substituted thiadiazole derivatives) showed the maximum activities. It is clear that compound **8b** (N-(5-(1-cyano-2-phenylvinyl)-1,3,4-thiadiazol-2-yl)benzamide) showed the highest activity towards *E. Coli*, *B. Subtilis*, and *C. Albicans* (Tables II and III).

	$MIC~(\mu gmL^{-1})~(\text{zone of inhibition, }mm)$				
Compound No.	E. coli ECT	B. cereus CECT	B. subtilis CECT	C. albicans CECT	
4a	_	22.52 (8)	20.55 (4)	8.65 (4)	
4b	_	11.32(3)	21.01 (8)	18.73 (8)	
6a	_	6.05 (15)	16.88 (2)	22.89 (3)	
6b	_	12.30 (4)	12.40(4)	33.23 (6)	
8a	_	3.72(6)	18.22 (8)	16.58 (12)	
8b	_	22.38 (8)	24.18 (3)	22.83 (6)	
10a	_	20.46 (9)	19.15 (4)	23.64 (6)	
10b	12.50 (6)	18.12 (5)	16.45 (6)	100.00	
12a	_	0.40 (10)	18.23 (5)	16.42(2)	
12b	_	18.24(7)	22.01 (8)	4.13 (10)	
12c	_	10.05 (6)	0.61 (6)	30.23 (6)	
12d	11.34(4)	12.25(2)	22.45(8)	6.18 (4)	
14a	_	4.55(10)	6.25(15)	24.44(2)	
14b	_	16.22 (3)	18.32(8)	10.38 (4)	
14c	_	6.22(6)	14.40 (4)	0.03 (9)	
14d	_	0.40 (5)	12.55(12)	25.36 (8)	
15a	_	23.63 (6)	21.15(4)	50.17	
15b	_	26.12 (3)	8.22(2)	100.00 (5)	
15c	14.84	18.32(5)	10.33(5)	23.16 (9)	
15d	_	0.08(2)	8.36 (4)	12.42(2)	
16	_	22.33(5)	12.22 (8)	8.66 (6)	
18a	_	6.05 (6)	4.55 (10)	12.34(7)	
18b	_	3.13(10)	6.13 (4)	12.34(7)	
19	_	12.50 (10)	6.13(4)	12.50	
20	_	0.40 (5)	8.25 (6)	22.82 (8)	
21	_	0.05 (9)	14.36 (8)	20.11(4)	
Ampicillin	6.25	3.13	12.50 (10)	_	
Cycloheximide	—	—	—	12.50 (6)	

TABLE I Antimicrobial Activities of the Tested Compounds

EXPERIMENTAL

All melting points are uncorrected. IR spectra were recorded (KBr) on a Pye Unicam SP-1000 spectrophotometer. ¹H NMR spectra were obtained on a Varian EM-390 MHz spectrophotometer in DMSO- d_6 as the solvent and TMS as internal reference. Chemical shifts are expressed in δ or ppm. Antibacterial results were recorded by a research group at the Botany Department at Cairo University.

5-Cyanomethyl-2-phenylamino-1,3,4-thiadiazole (4a) and 2-Benzoylamino- 5-cyanomethyl-1,3,4-thiadiazole (4b)

A suspension of the corresponding thiosemicarbazide either 3a (2.34 g, 0.01 mol) or 3b (2.62 g, 0.01 mol) in cold concentrated sulfuric acid

Compd	Viold (%)	M n °C Cryst	Mol	Calcd/Found (%)			6)
No.	color	solvent	Formula (Mr)	С	Н	Ν	S
4a	70 Buff	173 - 176	$\mathrm{C_{10}H_8N_4S}$	55.54	3.73	25.91	14.83
		Ethanol	(216.26)	55.62	4.03	26.24	14.71
4b	74 Yellow	262 - 264	$C_{11}H_8N_4OS$	54.09	3.30	22.94	13.13
		Ethanol	(244.27)	53.96	3.53	23.31	13.04
6a	70 Reddish-	194–196	$C_{16}H_{12}N_6S$	59.98	3.78	26.23	10.01
	brown	Ethanol	(320.37)	60.33	4.01	26.52	9.83
6b	86 Red	252 - 255	$C_{17}H_{12}N_6OS$	58.61	3.47	24.12	9.20
		Ethanol	(348.38)	58.88	3.26	24.31	8.95
8a	62 Yellow	180-183	$\mathrm{C}_{17}\mathrm{H}_{12}\mathrm{N}_4\mathrm{S}$	67.08	3.97	18.41	10.53
		Ethanol	(304.37)	66.89	4.08	18.17	10.33
8b	50 Yellow	222-224	$C_{18}H_{12}N_4OS$	65.04	3.64	16.86	9.65
		Ethanol	(332.38)	64.82	3.91	17.04	9.36
10a	93 Yellow	>300 Ethanol	$C_{17}H_{11}N_3O_2S$	63.54	3.45	13.08	9.98.
			(321.35)	63.43	3.66	12.79	10.26
10b	69 Yellow	246-248	$C_{18}H_{11}N_3O_3S$	61.88	3.17	12.03	9.18
		Ethanol	(349.36)	62.16	3.43	11.89	9.04
12a	70 Orange	166-169	$C_{13}H_{10}N_6S$	55.30	3.57	29.77	11.36
	0	1.4–dioxan	(282.32)	55.09	3.69	29.53	11.08
12b	66 Yellow	132-135	C13HoN5OS	55.11	3.20	24.72	11.32
		1.4–Dioxan	(283.31)	54.89	2.93	24.48	11.17
12c	78 Orange	222–225 DMF	$C_{14}H_{10}N_6OS$	54.18	3.25	27.08	10.33
			(310.33)	53.87	3.38	26.85	10.61
12d	73 Pale	189–193	C14H9N5O2S	54.01	2.91	22.50	10.30
	vellow	1.4–Dioxan	(311.32)	53.85	3.27	22.83	10.17
14a	70 Pale	98–100 Ethanol	C20H14N6S	64.85	3.81	22.69	8.66
	brown		(370.43)	64.47	4.25	23.05	8.89
14b	54 Reddish	230-233	CooH10N=OoS	63.29	4.59	16.78	7.68
	brown	Ethanol	(417.48)	63.62	4.31	17.16	7.95
14c	60 Reddish	98–102 Ethanol	Co1H14NeOS	63.30	3.54	21.09	8.05
	brown		(398.45)	63 49	3 76	20.86	7.82
14d	78 Reddish	230-233	$C_{99}H_{10}N_{\pi}O_{9}S$	62.01	4.30	15.72	7.20
	brown	Ethanol	(445,50)	62.17	4 38	15 15	7 27
15a	73 Yellow	258-260	$C_{12}H_{10}N_eS_{2}$	49.66	3.21	26.73	20.40
100	10 10110	Ethanol	$(314\ 39)$	49 44	3 50	26 53	20.72
15h	75 Vellow	240-245	C1rH1rNrOoSo	49.84	4 18	19.39	17 74
100	10 10110.0	Ethanol	(361 44)	48.98	4 33	19 27	18.09
15c	64 Brown	>300 Ethanol	C14H10NcOS0	49 11	2.94	24 54	18 73
100	04 DIOWII		(342.41)	49.04	3 35	24.04	19.16
15d	50 Brown	>300 Ethanol	C16H1ENEO2S	49 34	3.88	17 98	16 47
104	50 DIOWII		(389.45)	49.66	3 72	17.81	16.97
16	57 White	>300	C ₁₁ H ₀ N ₂ OS	54 09	3 30	22.94	13 13
10	51 WIII0	1 4-Dioxan	$(244\ 27)$	53.85	3.67	23 37	12.84
18a	65 Yellow	282-284	$C_{11}H_{10}N_{2}O$	54 54	4 14	34 69	12.01
104	55 ICHOW	1 1_Diovon	(9/1 9/)	54 10	1 1 9	34 89	
		1,4-DI0Xall	(241.24)	04.10	4.40	04.00	

 TABLE II Physical and Analytical Data of the Newly Synthesized

 Products

(Continued on next page)

				Found/Calcd (%)			
Compd No.	Yield (%) color	M.p. °C Cryst. solvent	Mol. Formula (Mr)	С	Н	N	S
18b	50 Yellow	276–279	$\mathrm{C_{17}H_{14}N_{6}O}$	64.14	4.43	26.40	
		1,4–Dioxan	(318.33)	64.09	4.11	26.32	
19	75 Yellow	280 - 282	$C_{17}H_{12}N_6OS$	58.61	3.47	24.12	9.20
		Ethanol	(348.38)	58.89	4.11	24.32	9.33
20	50 Pale	226-229	$C_{18}H_{12}N_4OS$	65.04	3.64	16.86	9.65
	brown	Methanol	(332.38)	65.11	3.88	17.17	9.44
21	92 Orange	242-244	$C_{18}H_{11}N_3O_3S$	61.88	3.17	12.03	9.18
		Ethanol	(349.36)	62.26	3.45	12.48	8.76

 TABLE II Physical and Analytical Data of the Newly Synthesized

 Products (Continued)

(28 mL) was stirred for 10 min. Then the mixture, in each case, was allowed to warm to room temperature and stirred for an additional 30 min. The resulting mixture, in each case, was poured onto an ice/water mixture and made alkaline to pH 8 with aqueous ammonia. The solid product formed in each case was collected by filtration.

5-(α-Phenylhydrazocyanomethyl-2-phenylamino-1,3,4thiadiazole (6a), 2-Benzoylamino-5-(αphenylhydrazocyanomethyl-1,3,4-Thiadiazole (6b), and 4-Benzoyl-(a-phenylhydrazocyanomethyl)-3-thioxo-1,2,4-triazole (19)

To a cold solution $(0-5^{\circ}C)$ of aniline (1.02 g, 1.0 mL, 1.1 equiv.), conc. HCl (1.09 g, 0.92 mL, 3 equiv.) was added. The temperature was maintained at $0-5^{\circ}C$ and an aqueous solution of sodium nitrite (0.76 g, 1.1 equiv.) was added dropwise to the aniline solution. The reaction mixture was stirred for 15 min. The clear diazonium salt solution **5** was then added dropwise to a solution of either thiadiazole **4a** (2.16 g, 0.01 mol, 1 equiv.) in ethanol (10 mL), thiadiazole **4b** (2.44 g, 0.01 mol, 1 equiv.) in an ethanol/acetic acid mixture (10:2 mL), or triazole **16** (2.44 g, 0.01 mol) in ethanol/DMF (10:2 mL) containing sodium acetate (4.0 g) at $0-5^{\circ}C$. After the addition of the diazonium salt was complete, the reaction mixture was stirred at room temperature for 2 h and left in the refrigerator overnight. The solid product formed in each case was collected by filtration.

Compd No. IR $(\nu/cm-1)$ 1H NMR (δ ppm) (DMSO-d6) 3465-3341 (NH), 3068 (CH aromatic), 4.68 (s, 2H, CH₂), 7.26-7.33 (m, 4a 2989 (CH₂), 2227 (CN), 1620 (C=C), 5H, C₆H₅), 8.69 (s, 1H, NH) 1585 (C=N) **4b** 3455–3337 (NH), 3059 (CH aromatic), 4.70 (s, 2H, CH₂), 7.28–7.37 (m, 2981 (CH₂), 2224 (CN), 1687 (CO), 5H, C₆H₅), 8.85 (s, 1H, NH) 1623 (C=C), 1578 (C=N) 6a 3474-3328 (2 NH), 3055 (CH 7.31-7.45 (m, 10H, 2 C₆H₅), aromatic), 2224 (CN), 1629 (C=C), 8.64, 8.95 (2s, 2H, 2NH) 1576 (C=N) 6b 3465-3317 (2 NH), 3072 (CH 7.27-7.40 (m, 10H, 2 C₆H₅), 8.54, 9.66 (2s, 2H, 2NH) aromatic), 2220 (CN), 1685 (CO), 1636 (C=C), 1582 (C=N) 19 3544-3328 (NH), 3055 (CH aromatic), 5.66 (s, br, 1H, SH), 7.29-7.36 2222 (CN), 1687 (C=O), 1638 (C=C), (m, 10H, 2 C6H5), 8.61 (s, 1H, NH) 1577 (C=N) 6.91 (s, 1H, CH=C), 7.30-7.38 8a 3489-3331 (NH), 3058 (CH aromatic), 2225 (CN), 1643 (C=C), 1582 (C=N) (m, 10 H, 2 C₆H₅), 8.73 (s, 1H, NH). **8b** 3476-3340 (NH), 3055 (CH aromatic), 5.82 (s, 1H, CH=C), 6.87 (s, 1H, NH), 7.27-7.40 (m, 10H, 2 2223 (CN), 1685 (CO), 1639 (C=C), 1589 (C=N) C_6H_5 10a 3458-3339 (NH), 3054 (CH aromatic), 6.69 (s, 1H, coumarin H-4), 1687 (CO), 1648 (C=C), 1593 (C=N) 7.31–7.44 (m, 9H, C₆H₅, C₆H₄), 8.35 (s, 1H, NH) 10b 3473-3329 (NH), 3059 (CH aromatic), 6.54 (s, 1H, coumarin H-4), 1693, 1688 (2 CO), 1641 (C=C), 1585 7.28-7.42 (m, 9H, C₆H₅, C₆H₄), 8.32 (s, 1H, NH) (C=N)3450–3328 (NH₂, 2NH), 3046 (CH 12a 4.66 (s, 2H, NH₂), 7.03-7.36 (m, aromatic), 2220 (CN), 1641 (C=C), 6H, C₆H₅, pyridine H-3), 8.68, 1586 (C=N) 9.40 (2s, 2H, NH) 12b 3466-3332 (NH2, NH), 3048 (CH 4.68 (s, 2H, NH₂), 7.19-7.38 (m, aromatic), 2221 (CN), 1687 (CO), 6H, C₆H₅, pyridine H-3), 8.66 1640 (C=C), 1593 (C=N) (s, 1H, NH) 12c 3469-3330 (NH2, 2NH), 3055 (CH 4.71 (s, 2H, NH₂), 7.21-7.44 (m, aromatic), 2226 (CN), 1689 (CO), 6H, C₆H₅, pyridine H-3), 8.68, 1644(C=C), 1590 (C=N) 9.39 (2s, 2H, 2NH) 12d 3485-3341 (NH₂, NH), 3059 (CH 4.67 (s, 2H, NH₂), 7.23-7.46 (m, aromatic), 2224 (CN), 1692, 1688 6H, C₆H₅, pyridine H-3), 8.65 (2CO), 1640(C=C), 1588 (C=N) (s, 1H, NH) 14a 3477–3332 (NH₂, NH), 3054 (CH 4.68 (s, 2H, NH₂), 5.36 (s, 1H, aromatic), 2228, 2223 (2 CN), 1656 pyridine H-4), 7.29-7.38 (m, (C=C), 1588 (C=N) 10H, 2C₆H₅), 8.36 (s, 1H, NH) 3469-3343 (NH2, NH), 3053 (CH 14b $1.36 (t, 3H, J = 7.09 Hz, CH_3),$ aromatic), 2987, 2879 (CH₃, CH₂), $4.23 (q, 2H, J = 9.07 Hz, CH_2),$ 2220 (CN), 1689 (CO), 1654 (C=C), 4.67 (s, 2H, NH₂), 5.63 (s, 1H, pyridine H-4), 7.25-7.37 (m, 1587 (C=N) 10H, 2C₆H₅), 8.59 (s, 1H, NH)

TABLE III Spectral Data of the Newly Synthesized Products

Compd No.	IR (v/cm-1)	1H NMR (δ ppm) (DMSO-d6)
14c	3479–3338 (NH ₂ , NH), 3057 (CH aromatic), 2225, 2222 (2CN), 1705 (CO), 1659 (C=C), 1584 (C=N)	4.64 (s, 2H, NH ₂), 5.72 (s, 1H, pyridine H-4), 7.28–7.39 (m, 10H, 2C ₆ H ₅), 8.63 (s, 1H, NH)
14d	3459–3329 (NH ₂ , NH), 3059 (CH aromatic), 2229 (CN), 1689, 1684 (2 CO), 1648 (C=C), 1568 (C=N)	1.35 (t, 3H, J = 6.77, CH ₃), 4.26 (q, 2H, J = 6.77 Hz, CH ₂), 4.57 (s, 2H, NH ₂), 5.68 (s, 1H, pyridine H-4), 7.33–7.47 (m, 10H, $2C_6H_5$), 8.60 (s, 1H, NH)
15a	3475–3331 (2NH2, NH), 3053 (CH aromatic), 2219 (CN), 1645 (C=C), 1587 (C=N)	$\begin{array}{c} 4.47,5.60(2s,4H,2NH_2),7.277.39\\ (m,5H,C6H5),8.31(s,1H,NH) \end{array}$
15b	$\begin{array}{l} 3468-3319\ (2NH_2,\ NH),\ 3059\ (CH \\ aromatic),\ 2987,\ 2855\ (CH_3,\ CH_2), \\ 1680\ (CO),\ 1645\ (C=C),\ 1587\ (C=N) \end{array}$	$\begin{array}{l} \text{1.14 (t, 3H, J} = 7.66 \text{ Hz}, \text{CH}_3), 4.23 \\ \text{(q, 2H, J} = 7.66 \text{ Hz}, \text{CH}_2), 4.45, \\ \text{5.68 (2s, 4H, 2NH}_2), 7.287.37 \\ \text{(m, 5H, C}_6\text{H}_5), 8.44 (s, 1H, NH) \end{array}$
15c	3488–3328 (2NH ₂ , NH), 3066 (CH aromatic), 2221 (CN), 1689 (CO), 1645 (C=C), 1587 (C=N)	$\begin{array}{l} \text{4.67, 5.37 (2s, 4H, 2NH_2), 7.32-7.38} \\ \text{(m, 5H, C_6H_5), 8.48 (s, 1H, NH)} \end{array}$
15d	$\begin{array}{l} 3473-3321\ (2NH_2,\ NH),\ 3056\ (CH \\ aromatic),\ 2980\ (CH3),\ 2877\ (CH_2), \\ 1689,\ 1686\ (2\ CO),\ 1642\ (C=C),\ 1581 \\ (C=N) \end{array}$	$\begin{array}{l} 1.42 \ (t, 3H, J{=} 6.62 \ Hz, CH_3), 4.21 \\ (q, 2H, J{=} 6.62 \ Hz, CH_2), 4.60, \\ 5.41 \ (2s, 4H, 2NH_2), 7.34{-}7.48 \\ (m, 5H, C_6H_5), 8.52 \ (s, 1H, NH) \end{array}$
16	3310 (SH), 3058 (CH aromatic), 1715 (CO), 2897 (CH ₂), 1652 (C=C), 1580 (C=N)	$\begin{array}{l} \text{4.82 (s, 2H, CH_2), 5.26 (s, br, 1H, } \\ \text{SH). 7.30-7.37 (m, 5H, C_6H_5)} \end{array}$
18a	3483–3322 (NH ₂ , NH), 3058 (CH aromatic), 2227 (CN), 1710 (CO), 1651 (C=C), 1590 (C=N)	$\begin{array}{l} 4.73~(s,2H,NH_2),4.85~(s,2H,\\ CH_2),7.277.40~(m,5H,C_6H_5),\\ 8.42~(s,1H,NH) \end{array}$
18b	3483–3322 (2 NH), 3061 (CH aromatic), 2224 (CN), 1713 (CO), 1657 (C=C), 1588 (C=N)	$\begin{array}{l} 4.84\ (s,2H,CH_2),7.29{-}7.38\ (m,\\ 10H,2C_6H_5),8.39,8.68\ (2s,2H,\\ 2NH) \end{array}$
20	3055 (CH aromatic), 2223 (CN), 1688 (CO), 1639 (C=C), 1589 (C=N)	5.83 (s, br, 1H, SH), 6.79 (s, 1H, CH=C), 7.27–7.40 (m, 10H, 2 C_6H_5)
21	3058 (CH aromatic), 1693, 1682 (2 CO), 1652 (C=C), 1578 (C=N)	$\begin{array}{l} 6.34\ ({\rm s},\ 1{\rm H},\ coumarin\ {\rm H}\text{-}4),\ 6.61\ ({\rm s},\\ {\rm br},\ 1{\rm H},\ {\rm SH}),\ 7.357.41\ ({\rm m},\ 9{\rm H},\\ {\rm C}_{6}{\rm H}_{5},\ {\rm C}_{6}{\rm H}_{4}) \end{array}$

TABLE III Spectral Data of the Newly Synthesized Products

 (Continued)

3-Phenyl-2-(5-(phenylamino)-1,3,4-thiadiazol-2-yl)acrylonitrile (8a), N-(5-(1-Cyano-2-phenylvinyl)-1,3,4-thiadiazol-2-yl) benzamide (8b), and 2-(4,5-Dihydro-5-thioxo-3H-1,2,4triazol-3-yl-3-phenylacrylonitrile (20)

To a solution of either **4a** (1.08 g, 0.005 mol), **4b** (1.22 g, 0.005 mol), or **16** (1.22 g, 0.005 mol) in DMF (25 mL) containing piperidine (0.5 mL),

benzaldehyde (0.53 g, 0.51 mL, 0.005 mol) was added. The reaction mixture, in each case, was heated under refluxed for 4 h, and then cooled and poured onto an ice/water mixture. The pH was made acidic using hydrochloric acid. The solid product formed, in each case, was collected by filtration.

7-Amino-5-imino-2-(phenylamino)-*5H*-[1,3,4]thiadiazolo[3,2a]pyridine-8-carbonitrile (12a), 7-Amino-5-oxo-2-(phenylamino)-*5H*-[1,3,4]thiadiazolo[3,2-a]pyridine-8carbonitrile (12b), N-(7-Amino-8-cyano-5-imino-5H-[1,3,4-]thiadiazolo[3,2-a]pyridine-2-yl)benzamide (12c), and N-(7-Amino-8-cyano-5-oxo-5H-[1,3,4-]thiadiazolo [3,2-a]pyridine-2-yl)benzamide (12d)

To a solution of either thiadiazole 4a (2.16 g, 0.01 mol) or thiadiazole 4b (2.44 g, 0.01 mol) in 1,4-dioxane (50 mL) containing triethylamine (1.01 g, 1.4 mL, 1.0 equiv.), either malononitrile (0.66 g, 0.01 mol) or ethyl cyanoacetate (1.07 g, 0.01 mol) was added. The reaction mixture in each case was heated under reflux for 3 h then poured onto ice water containing a few drops of hydrochloric acid. The formed solid product, in each case was collected by filtration.

2,4-Diamino-5-(5-phenylamino)-1,3,4-thiadiazol-2yl)thiophene-3-carbonitrile (15a), Ethyl 2,4-diamino-5-(5-(phenylamino)-1,3,4-thiadiazol-2-yl)thiophene-3-carboxylate (15b), N-(5-(3,5-Diamino-4-cyanothiophene-2-yl)-1,3,4thiadiazol-2-yl)benzamide (15c), and Ethyl 2,4-Diamino-5-(5-benzamido)-1,3,4-thiazol-2-yl)thiophene-3-carboxylate (15d)

To a solution of either thiadiazole 4a (2.16 g, 0.01 mol) or thiadiazole 4b (2.44 g, 0.01 mol) in 1,4-dioxane (50 mL) containing triethylamine (1.01 g, 1.4 ml, 1.0 equiv.) and elemental sulfur (0.32 g, 0.01 mol), either malononitrile 11a (0.66 g, 0.01 mol) or ethyl cyanoacetate 11b (1.13 g, 1.1 ml, 0.01 mol) was added separately. The resulting reaction mixture in each case was heated under reflux for 4 h then poured onto an ice/water mixture, and the pH was made acidic using hydrochloric acid. The formed solid product, in each case, was collected by filtration.

3-(5-Phenylamino-1,3,4-thiadiazol-2-yl)-2H-chromen-2-one (10a), N(5-Oxo-2H-chromen-3-yl)-1,3,4-thiadiazol-2-yl) benzamide (10b), and 3-(4,5-Dihydro-5-thioxo-3H-1,2, 4-triazol-3-yl)-2H-chromen-2-one (21)

To a solution of either thiadiazole 4a (1.08 g, 0.005 mol), thiadiazole 4b, (1.22 g, 0.005 mol), or triazole 16 (1.22 g, 0.005 mol) in 1,4-dioxane (25 mL) containing a few drops of piperidine, salicylaldehyde (0.67 g, 0.58 mL, 0.005 mol, 1.1 equiv.) was added. The reaction mixture, in each case, was heated under reflux for 4 h, and then cooled and poured onto an ice/water mixture. Then the pH was made acidic using hydrochloric acid. The solid product formed, in each case, was collected by filtration.

2-(1-Benzoyl-4,5-dihydro-5-thioxo-3H-1,2, 4-triazol-3-yl)acetonitrile (16)

A solution of thiosemicarbazide **3b** (2.62 g, 0.01 mol) in sodium hydroxide (2 N, 28 mL) was refluxed for 3 h. The resulting solution was cooled to room temperature and acidified to pH 3–4 with 37% hydrochloric acid. The solid product formed was collected by filtration.

2(1-Benzoyl-5-hydrazino-1,2,4-Triazol-3-yl)axetonitrile (18a) and 2(1-Benzoyl-5-phenylhydrazino-1,2,4-triazol-3-yl) axetonitrile (18b)

To a solution of triazole 16 (1.22 g, 0.005 mol) in ethanol (25 mL), either hydrazine hydrate (0.25 g, 0.005 mol) or phenylhydrazine (0.54 g, 0.49 mL, 0.005 mol) was added. The reaction mixture, in each case, was heated under reflux for 4 h, and then poured onto an ice/water mixture containing few drops of hydrochloric acid. The solid product formed in each case was collected by filtration.

5-Amino-7-phenyl-2-(phenylamino)-7H-[1,3,4]thiadiazolo[3,2a]pyridine-6,8-dicarbonitrile (14a), Ethyl 5-Amino-8-cyano-7-phenyl-2-(phenylamino)-7H-[1,3,4]thiadiazolo[3,2-a]pyridine-6-carboxylate (14b), N-(5-Amino-6,8-dicyano-7-phenyl-7H-[1,3,4]thiadiazolo[3,2-a]pyridine-2-yl0benzamide (14c), and Ethyl 5-Amino-2-(benzamido)-8-cyano-7-phenyl-7H-[1,3,4] thiadiazolo[3,2-a]pyridine-6-carboxylate (14d)

To a solution of either the thiadiazol4a (1.2 g, 0.005 mol) or 4b (1.08, 0.005 mol) in DMF (30 mL) containing triethylamine (0.5 g, 0.7 ml, 1.0 equiv.), either benzalmalononitrile 13a (0.77 g, 0.005 mol) or

benzalethyl cyanoacetate 13b (1.0 g, 0.005 mol) was added separately. The resulting reaction mixture in each case was heated under reflux for 4 h then poured onto an ice/water mixture and the pH was made acidic using hydrochloric acid. The formed solid product, in each case, was collected by filtration.

REFERENCES

- [1] M. M. Suni, V. A. Nair, and C. P. Joshua, Tetrahedron, 57, 2003 (2001).
- [2] D. Limal, V. Grand, R. Vanderesse, M. Marraud, and A. Aubry, *Tetrahedron Lett.*, 35, 3711 (1994).
- [3] S. Kabashima, T. Okawara, T. Yamasaki, and M. Furukawa, J. Heterocycl. Chem., 28, 1957 (1991).
- [4] G. G. Marian and K. Schulze, J. Heterocycl. Chem., 32, 275 (1995).
- [5] P. L. Meo, R. Noto, and G. Werber, J. Heterocycl. Chem., 30, 765 (1993).
- [6] R. Noto, M. Gruttadauria, P. L. Meo, V. Frenna, and G. Werber, J. Heterocycl. Chem., 32, 1277 (1995).
- [7] R. Noto, P. L. Meo, M. Gruttadauria, and G. Werber, J. Heterocycl. Chem., 33, 863 (1996).
- [8] H. Gruttadauria, P. L. Meo, R. Noto, and G. Werber, Gazz. Chim. Ital., 127, 277 (1997).
- [9] R. Noto, P. L. Meo, M. Gruttadauria, and G. Werber, J. Heterocycl. Chem., 36, 667 (1999).
- [10] A. A. Hassan, Bull. Soc. Chim. Fr., 131, 424 (1994).
- [11] A. A. Hassan, Y. R. Ibrahim, A. A. Semida, and A. E. Mourad, *Liebigs Ann. Chem.*, 989 (1994).
- [12] A. A. Hassan, Phosphorus, Sulfur, and Silicon, 101, 189 (1995).
- [13] A. A. Hassan, Y. R. Ibrahim, E. H. El-Tamany, A. A. Semida, and A. E. Mourad, *Phosphorus, Sulfur, and Silicon*, **106**, 167 (1995).
- [14] A. A. Hassan, N. K. Mohamed, A. A. Aly, and A. E. Mourad, *Monatsh. Chem.*, 128, 61 (1997).
- [15] R. A. Mekheimer and R. M. Shaker, J. Chem. Res. (S), 76 (1999); J. Chem. Res. (M), 445 (1999).
- [16] H. Z. Shams, R. M. Mohareb, M. H. Helal, and A. E. Mahmoud, *Phosphorus, Sulfur, and Silicon.*, 182, 237 (2007).
- [17] D. I. Brahmbhatt, S. Shashibala, and K. C. Patel, Eur. Polym. J. 35, 317 (1999).
- [18] N. K. Chodankar and S. Seshadri, Dyes Pigm., 6, 331 (1984).
- [19] M. Zhang and R. W. Harper, Biorg. Med. Chem. Lett., 7, 1629 (1997).
- [20] G. M. Castanedo and D. P. Sutherlin, Tetrahedron Lett., 42, 7181 (2001).
- [21] P. M. Hawkey and A. A. Lewis, *Medical Bacteriology—A Practical Approach*, (Oxford University Press, Oxford, UK, 1994), 2nd ed., pp. 181–194.
- [22] N. Rameshkumar, M. Ashokkumar, E. H. Subramanian, E. R. Ilavarasan, and S. K. Sidhar, Eur. J. Med. Chem., 38, 1001 (2003).