Dalton Transactions

An international journal of inorganic chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Q. Liang, J. H. Lin, J. C. DeMuth, M. L. Neidig and D. Song, *Dalton Trans.*, 2020, DOI: 10.1039/D0DT02087G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the Information for Authors.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/dalton

View Article Online

View Journal

ARTICLE

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Syntheses and Characterizations of Iron Complexes of Bulky *o*-Phenylenediamide ligand

Qiuming Liang, ^a Jack H. Lin, ^a Joshua C. DeMuth, ^b Michael L. Neidig ^b and Datong Song *^a

We report the syntheses of a family of tetrahedral iron complexes bearing a bulky redox active *o*-phenylenediamide ligand. The electronic structures of these complexes have been investigated by Mössbauer spectroscopy, magnetic susceptibility measurements, and X-ray crystallography.

Introduction

Published on 17 August 2020. Downloaded on 8/18/2020 1:41:39 AM

Within the last decade, redox-active ligands have attracted considerable attention, owing to their unusual and intriguing electronic structures.¹ Redox-active ligands have more energetically accessible levels for reduction or oxidation.¹ Their coordination to metal centres induces radical reactivity and electron reservoir behaviour, which is often utilized to develop exciting catalytic activities.1-3 The oxidation level of the redoxactive ligands in complexes can be determined by high-quality X-ray crystallography, a broad set of spectroscopic methods, and density functional theory (DFT) calculations. A combination of a redox active metal, such as iron, with redox-active ligands can lead to unique chemical and magnetochemical properties. In these cases, the determination of the oxidation state of the transition-metal ion is not straightforward. For instance, a ferric ion in such species may adopt a high-, intermediate-, or low-spin electronic configuration ($S_{Fe} = 5/2$, 3/2, or 1/2, respectively; S is the total spin quantum number), whereas a ferrous ion may also have high-, intermediate-, and low-spin electronic configurations (S_{Fe} = 2, 1, or 0, respectively). Strong intramolecular antiferromagnetic coupling of a radical ligand or a second metal centre may yield a large variety of different spin ground states.

 α -Diimines and *o*-phenylenediamides (pda) are classic examples of redox-active ligands that form five-membered chelate rings with a large variety of transition and main-group elements.⁴⁻⁶ It has been clearly established that α -diimines and pda ligands each exist in three different redox forms in coordination compounds, including a closed-shell dianion (*S* = 0), an open-shell π radical monoanion (S = 1/2), and a closedshell neutral (S = 0) form (Scheme 1).⁴ Iron complexes based on α -diimine ligands including 1,4-diaza-1,3-butadiene (dad) and bis(imino)acenaphthene (BIAN) were widely investigated.⁷⁻⁹ Furthermore, iron complexes of α -diimine ligands are efficient catalysts for the hydrogenation of alkenes,^{8a,9a} hydrovinylation of dienes,^{8b} (cyclo)dimerization of dienes,^{8c-e} cyclotrimerization of alkynes,^{8f} Diels–Alder type cycloaddition of dienes with alkynes,^{8f} dehydrogenation of amine–boranes,^{8g} and hydrosilylation of carbonyls.^{9b,c}



Scheme 1. Different redox states of α -diimine and *o*-phenylenediamide (pda) derived ligands.

In contrast to iron α -diimine complexes, iron pda complexes are relatively less studied. Wieghardt and co-workers have reported several homoleptic pda iron complexes and dad-pda heteroleptic iron complexes, which displayed sophisticated electronic structures.⁷ Hernán-Gómez and Costas have recently reported the reactivity of $[Fe(^{C6F5}pda)(THF)]_2$ ($^{C6F5}pda = N,N'$ bis(pentafluorophenyl)-o-phenylenediamido-) toward the intramolecular C(sp³)-H activation of diazoesters in the presence of LiAI[OC(CF₃)₃]₄ under mild reaction conditions, yielding a variety of carbocyclic cyclopentanes as well as bicyclic spiro and fused ring compounds.^{7f} Our group has reported the [Fe^I(^{Dipp}pda⁻⁻)(toluene)] (^{Dipp}pda = N,N'-bis(2,6-diisopropylphenyl)-o-phenylenediamido-) and [Fe⁰(^{Dipp}pda⁰)(CO)₃] complexes.^{7g} We herein report the synthesis of a series of iron compounds supported by the Dipppda ligand platform at various oxidation

^{a.} Davenport Chemical Research Laboratories, Department of Chemistry, University of Toronto, 80 St. George Street, Toronto, Ontario, Canada, M5S 3H6.

^{b.} Department of Chemistry, University of Rochester, Rochester, New York, USA,

^{14627.}

E-mail: d.song@utoronto.ca

⁺ Electronic supplementary information (ESI) available: NMR and Mössbauer spectra and X-ray crystallographic experimental details. CCDC 2009196–2009200, 2009218 and 2022767–2022770. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/x0xx00000x.

Journal Name

states. Spectroscopic data as well as X-ray crystallographic analysis are used to characterize this series of new complexes.



Scheme 2. Synthesis of compound 2.



Figure 1. Molecular structure of 2 (30% probability thermal ellipsoids). All hydrogen atoms are omitted for clarity.

Results and Discussion

Published on 17 August 2020. Downloaded on 8/18/2020 1:41:39 AM

Reacting DipppdaH₂ (1) with [FeCl(HMDS)₂(THF)] (HMDS = hexamethyldisilazide) in toluene at 120 °C gives complex 2 in good yields (Scheme 2). The use of bulky Dipp substituents is important, as smaller pda ligands tend to form [Fe(pda)₂].⁷ The 1H NMR spectrum of complex $\boldsymbol{2}$ in C_6D_6 at 25 $^\circ C$ shows paramagnetically shifted and broadened resonances between +159 and -15 ppm. X-ray crystallographic analysis of 2 revealed a dinuclear structure with a crystallographically imposed inversion centre (Figure 1). The bond lengths within the Dipppda ligands reveal a quinoid-type distortion. The C(1)-N(1) and C(6)–N(2) bond lengths are 1.341(2) and 1.346(2) Å (Table 1), respectively, in between the typical C-N single and double bond lengths. The C_{α} - $C_{\alpha'}$ and C_{γ} - $C_{\gamma'}$ bond lengths are 1.452(2) and 1.405(3) Å, respectively, while the other four bonds of the phenylene backbone feature the long C_{α} -- C_{β} and $C_{\alpha'}$ -- $C_{\beta'}$ bonds (1.419(2) and 1.421(2) Å) and short $C_{\beta}-C_{\gamma}$ and $C_{\beta'}-C_{\gamma'}$ bonds (1.363(3) and 1.361(3) Å). The pattern in these metric parameters suggests that the pda ligands are open-shell $\boldsymbol{\pi}$ radical anions, i.e., $^{Dipp}pda^{-}$ (S = 1/2).⁴ As such, the iron centres are Fe(II) to balance the charge. The Fe(1)–N(1) and Fe(1)–N(2) bond lengths are 1.979(2) and 1.977(1) Å, respectively, suggesting that the Fe^{II} centres are high-spin, i.e., $S_{Fe} = 2$. The intramolecular Fe–Fe distance of 3.0255(4) Å in 2 is considerably longer than the sum of the covalent radii (2.50 Å), making it outside of the range of Fe-Fe bonds.¹⁰ The Fe-Fe distance is slightly longer than that found in $[Fe(\mu_2-CI)(^{Dipp}dad)]_2$ (2.9764(6) Å, Dippdad = N,N'-bis-(2,6-diisopropylphenyl)-2,3-dimethyl-1,4diaza-1,3-butadiene),^{7a,b} suggesting an even weaker exchange

coupling between the two iron centres. The solution magnetic moment of **2** is 5.4 μ_B in C₆D₆ at 25 °C (Table 2), while [Fe(μ_2 -Cl)(^{Dipp}dad)]₂ was reported to have solution^{7a} magnetic moment of 3.86 μ_B . The greater solution magnetic moment of compound **2** compared to that of [Fe(μ_2 -Cl)(^{Dipp}dad)]₂ is likely due to a weaker antiferromagnetic coupling between the two iron centres in **2**.

Zero-field ⁵⁷Fe Mössbauer spectroscopy was used to examine the oxidation state and spin state of the iron centres of complex **2** (Table 2). The Mössbauer spectrum of complex **2** at 80 K features one doublet with an isomer shift (δ) of 0.84 mm/s and a quadruple splitting ($|\Delta E_Q|$) of 3.64 mm/s (Figure S10), typical for high-spin Fe^{II} species.^{7c,11} Such a spectrum suggests that the two high-spin ferrous centres are equivalent, consistent with the crystallographic data. Overall, the experimental data are consistent with an electronic structure with strong antiferromagnetic coupling within each ^{Dipp}pda⁺⁻⁻ Fe^{II} pair and weak coupling between the two iron centres across the bridging chloride ligands.¹²

With complex 2 in hand, we set to reduce the metal centre to low valence, hoping for dinitrogen binding. Compound 2 can be reduced with two equivalents of 0.5% sodium amalgam to afford the diamagnetic complex 3 (Scheme 3), featuring a lowspin Fe(I) centre and a Dipppda⁻⁻ ligand with strong antiferromagnetic coupling in between.^{7g} Attempts to reduce complex 2 in non-arene solvents were not fruitful. The reduced product is able to capture the trace amount of arene vapour in the glovebox atmosphere to give [Fe(^{Dipp}pda)(n⁶-arene)], which prevents dinitrogen from binding. The Mössbauer spectrum of 3 (Table 2) features a doublet with an isomer shift of 0.37 mm/s, typical for a low spin ($S_{Fe} = 1/2$) Fe(I) centre,¹¹ and a quadrupole splitting of 0.79 mm/s (Figure S11). While the isomer shift is similar to the analogous $Fe(^{Xyl}dad)(n^6-toluene)$ ($\delta = 0.44$ mm/s, $|\Delta E_Q| = 0.41 \text{ mm/s})^{8h}$ and Fe(^{Dipp}BIAN)(η^6 -toluene) ($\delta = 0.45$ mm/s, $|\Delta E_Q| = 0.41$ mm/s),⁹ the quadruple splitting is slightly larger.

Previously we showed that replacing the η^{6} -toluene ligand in $\boldsymbol{3}$ with three stronger charge-neutral $\pi\text{-acceptor}$ CO ligands resulted in the formation of the [Fe(pda)(CO)₃] where pda is in its charge neutral form and oxidation state of the iron centre is zero, i.e., shift from (Dipppda*-)FeI to (Dipppda0)Fe0.7g We reason that if we replace the η^6 -toluene ligand with charge-neutral σ donor ligands instead, we may be able to achieve the opposite, i.e., shift from (Dipppda*-)FeI to (Dipppda2-)FeII. The addition of two or more equivalents of pyridine to a THF solution of 3 affords a four-coordinate complex 4 (Scheme 3). The X-ray crystal structure of 4 shows a distorted tetrahedral coordination geometry of the iron centre (Figure 2). The C(1)–N(1) and C(6)– N(2) bond lengths are 1.392(3) and 1.399(3) Å (Table 1), respectively, which are close to a typical C–N single bond length. The C_{α} – $C_{\alpha'}$ and C_{ν} – $C_{\nu'}$ bond lengths are 1.430(3) and 1.367(4) Å, respectively, while the other bond lengths of the phenylene backbone are more even, i.e., within the range of 1.388(4)-1.397(4) Å. Although a crystal structure with a higher resolution is preferable (i.e., to see whether the $C_v - C_{v'}$ bond length is significantly different from the $C_{\beta}-C_{\gamma}$ and $C_{\beta'}-C_{\gamma'}$ bonds), the current metric parameters suggest that the N,N-chelating ligand

ARTICLE



Figure 2. Molecular structure of 4, 5 and 7 (30% probability thermal ellipsoids). All hydrogen atoms are omitted for clarity

is a closed-shell dianion $^{\text{Dipp}}\text{pda}^{2-}$ (*S* = 0). To balance the overall charge, the iron centre is a ferrous ion. The solution magnetic moment of **4** is 5.2 μ_{B} in THF-*d*₈ at 25 °C (Table 2), suggesting a high-spin Fe^{II}. Complex **4** exhibits isomer shift of 0.72 mm/s and large quadrupole splitting of 4.02 mm/s (Table 2, Figure S12). These values are characteristic for high-spin ferrous species (*S*_{Fe} = 2),^{7c,11} corroborating with the crystallographic data and solution magnetic moment. The dissolution of **4** in arene solvents results the immediate formation of [Fe(^{Dipp}pda)(n⁶-arene)] complexes accompanied by a colour change from blue–purple to intense red–purple. The nature of the auxiliary ligand on iron clearly plays a key role on the electronic structure of the pda-Fe moiety.

In comparison, we examined the ligand substitution reactions of **2**, which has a (^{Dipp}pda^{•-})Fe(II) moiety to find out whether we could achieve the electron shift between the pda

ligand and the iron centre by using different auxiliary ligands. The reaction of 2.5 equivalents of pyridine with **2** in toluene affords the four-coordinate complex **5** (Scheme 3). The X-ray crystal structure of **5** revels a distorted tetrahedral coordination geometry at the iron centre. The Fe(1)–N(1) and Fe(1)–N(2) bond lengths of 1.995(3) and 2.018(3) Å (Table 1) suggest a high-spin Fe^{II} ($S_{Fe} = 2$) centre. The C–C and C–N bond lengths of the pda ligand of **5** and the Mössbauer parameters ($\delta = 0.81$ mm/s, $|\Delta E_Q| = 2.05$ mm/s, Table 2) are similar to those of **2**, suggesting a π radical anion ^{Dipp}pda^{•-} (S = 1/2). The solution magnetic moment of **5** is 4.2 μ_B in C₆D₆ at 25 °C (Table 2), suggesting strong antiferromagnetic coupling between the high-spin ferrous centre ($S_{Fe} = 2$) and the π radical anionic ligand (S = 1/2).

Unable to push an electron from Fe(II) into Dipppda^{•-} with the charge-neutral pyridine ligand, we turned to the anionic π -donor chloride ligand. The addition of excess tetrabutylammonium chloride

Journal Name

ARTICLE

Published on 17 August 2020. Downloaded on 8/18/2020 1:41:39 AM

(5 equiv.) to a toluene solution of **2** at room temperature affords the four-coordinate ferrous salt **6** (Scheme 3). In the solid-state structure of **6**, the C–C and C–N bond lengths of the pda ligand are similar to those of **2**, i.e., a π -radical anion ^{Dipp}pda^{•-} (S = 1/2). Therefore, the iron centre is assigned as Fe^{II} to balance the charge. The long Fe–N bonds (2.018(2) and 2.003(2) Å) suggest that the Fe^{II} centre in **6** is high-spin. The Mössbauer parameters ($\delta = 0.80 \text{ mm/s}$, $|\Delta E_Q| = 3.04 \text{ mm/s}$, Table 2) are similar to those of **2** and **5**. The solution magnetic moment of **6** is 4.1 μ_B in C₆D₆ at 25 °C (Table 2), suggesting strong antiferromagnetic coupling between the ^{Dipp}pda^{•-} ligand and high-spin Fe^{II} centre.

The anionic π -donor chloride ligand is unable to push an electron from Fe(II) into Dipppda^{•-}, presumably because the Dipppda^{•-} ligand is not reducing enough. We are curious to know what product we will get if we oxidize complex 6 by 1e⁻, i.e., [Fe^{II}(Dipppda⁰)Cl₂] or [Fe^{III}(Dipppda^{•-})Cl₂]. The reaction of 6 with one equivalent of silver tetrafluoroborate at room temperature in toluene results in a colour change from dark blue-purple to dark green. Compound 7 can be isolated from the reaction mixture as the major product (Scheme 3). The solid-state structure of 7 reveals a C_s symmetry and a distorted tetrahedral coordination geometry at the iron centre (Figure 2). The metric parameters of the pda ligand in 7 are similar to those in 2, 5, and 6, i.e., Dipppda⁻⁻, whereas the Fe-N bonds (1.967(2) Å) in 7 (Table 1) are slightly shorter than those in 2, 5, and 6. The solution magneticmoment of **7** is 5.1 μ_B in C₆D₆ at 25 °C (Table 2). The electronic structure of **7** can be described as [Fe^{III}Cl₂(^{Dipp}pda^{•-})], with strong antiferromagnetic coupling between the π -radical anion Dipp pda⁻⁻ (S = 1/2) ligand and high-spin ferric ion (S = 5/2). The isomer shift and the quadrupole splitting of 7 are 0.39 mm/s and 2.12 mm/s (Table 2, Figure S15), respectively. The reduced isomer shift is consistent with a more oxidized iron centre. In contrast, $[FeCl_2(dad)]^{7a,b}$ and $[Fe(^{Dipp}BIAN)X_2]^{9j,k}$ were found to contain Fe^{II} centres and neutral N,N-chelate ligands. It appears that the Dipppda*is less reducing than the corresponding ^{Dipp}BIAN^{•-} and dad^{•-} ligands. The minor impurity in the sample of 7 displays similar isomer shift and quadruple splitting to those of 3, whose formation from the reaction of 6 and AgBF₄ in toluene could be rationalized as the initial chloride abstraction from 6 by Ag⁺ followed by toluene coordination to the iron centre.

As mentioned above, the bulky Dipp groups are crucial to the synthesis of **2** in minimizing the formation of $[Fe(^{Dipp}pda)_2]$. In order to prepare the homoleptic complex [Fe(^{Dipp}dpa)₂], 8, we adopted Wieghardt's synthetic procedures (Scheme 4),^{7c} i.e., reacting the doubly deprotonated 1 with FeCl₂ followed by oxidation with FeCl₃. The molecular structure of 8 consists of two chelating Dipppda ligands bound to a distorted tetrahedral iron centre (Figure 3). The metric parameters of the pda ligands in 8 (Table 1) are in between those of the Dipppda⁻⁻ and Dipppda²⁻ ligands. For example, the C(1)–N(1) and C(6)–N(2) bond lengths are 1.372(3) and 1.376(3) Å, respectively, whereas the corresponding C-N bond lengths in the lithium salts of the ^{Dipp}pda⁻⁻ and ^{Dipp}pda²⁻ are 1.341(3) and 1.395(2) Å, respectively. The solution magnetic moment of **8** is 2.7 μ_B in C₆D₆ at 25 °C (Table 2). The crystallographically imposed 2-fold axis relating the two Dipppda ligands suggests two identical ligands within one molecule of 8. The electronic structure of a closely related complex [Fe(C6F5pda)₂] was studied in detail by Wieghardt and

co-workers and reported as [Fe^{III}(^{C6F5}pda^{•-})(^{C6F5}pda²⁻)]_{ri}with ap intermediate-spin Fe^{III} (S_{Fe} = 3/2) antiferPornaghetically-coupled to a ligand-based radical.^{7c} Based on the literature precedence and the available data, we tentatively rationalize the overall electronic structure of 8 using the following major resonance [Fe^{III}(^{Dipp}pda^{•-})(^{Dipp}pda²⁻)] structures [Fe^{III}(Dipppda²⁻)(Dipppda⁻⁻)] in a 1:1 ratio, i.e., a fully delocalized radical over two chelate ligands is antiferromagnetically coupled to an intermediate-spin ferric centre (S = 3/2). The Fe-N bond lengths in 8 (1.950(2) and 1.973(2) Å) are longer than those in [Fe^{III}(^{C6F5}pda^{•–})(^{C6F5}pda^{2–})] (avg. 1.897(3) Å), which could be due to the significantly increased steric congestion associated with the two Dipppda ligands. Unfortunately, the Mössbauer data for **8** (δ = 0.42 mm/s, $|\Delta E_Q|$ = 2.51 mm/s, Table 2, Figure S16) fall in a range for which an unambiguous oxidation state and spin state assignment for the iron centre is not possible.11







Figure 3. Molecular structure of 8 (30% probability thermal ellipsoids). All hydrogen atoms are omitted for clarity.

The structure of **8** was also examined with broken symmetry DFT calculations. The full structure of **8** was optimized into the same triplet ground state starting from various broken symmetry guesses, using PBEPBE¹³ functionals and TZVP¹⁴ basis set with Gaussian 16 software.¹⁵ The optimized geometry closely resembles that of the crystal structure, as evidenced by the bond length data shown in Table 1. As shown in Figure 4, the spin density on Fe is 2.442, slightly lower than the expected value of an intermediate spin Fe(III), whereas the spin density of -0.442 is fully delocalized over the two dpa ligands with significant spin density on the nitrogen donor atoms. Such a spin density distribution is consistent with the proposed electronic structure above, i.e., an intermediate spin Fe(III) with significant

Journal Name

covalency and antiferromagnetic coupling between the metal centre and the two pda ligands.

TDDFT calculation was performed using PBEPB functionals and def2svp¹⁶ basis set on the optimized structure from broken symmetry DFT calculation using the same functionals and basis set. The TDDFT simulated UV-Vis-NIR spectrum matches well with the experimental spectrum in terms of the wavelength of the absorption but not the intensity (Figure S21), which is expected. The frontier orbitals (Figures S22 and S23) and electronic transitions responsible for the low energy absorption bands are listed in Table S3. Due to the involvement of multiple transitions in each excited state and the mixed metal and ligand contribution to each frontier orbital, it is difficult to classify the nature of the low-energy absorption bands.



Figure 4. Computed spin density map for complex **8**. The Dipp groups on the dpa ligands are omitted from this plot for clarity, although they were included in the calculation. Colour key: orange for Fe, blue for N, grey for C, and white for H.



Scheme 5. Syntheses of complexes 9–12. The electronic structures of 9, 11, and 12 are ambiguous with available data; the tentative drawings of these compounds are just to show that the two [Dipppda–Fe] fragments within each molecule are different.



Figure 5. Molecular structure of 9–12 (30% probability thermal ellipsoids). All hydrogen atoms are omitted for clarity.

alton Transactions Accepted Manuscr

View Article Online DOI: 10.1039/D0DT02087G

ARTICLE

In an effort to prepare mononuclear low coordinate iron complexes for small molecule activation, we set to substitute the bridging chloride ligand in 2 with a bulky anionic ligand and break the dimeric structure of 2. The addition of two equivalents of KO^tBu to a toluene solution of 2 at room temperature affords the dinuclear complex 9 with two bridging tert-butoxide ligands (Scheme 5). The observed solution magnetic moment 6.1 μ_B at 25 °C in C₆D₆ (Table 2). Interestingly, the metric parameters (Table 1) of the two Dipppda ligands are different in the X-ray structure of 9 (Figure 5). One of the $^{\text{Dipp}}\text{pda}$ ligands resembles that in complex $\boldsymbol{8}$ (C_{\alpha}\text{-}N bond lengths: 1.370(3) and 1.374(3) Å), while the other resembles that in complex **4** (C_α–N bond lengths: 1.394(3) and 1.395(3) Å). These metric parameters suggest that one of the two Dipppda ligands is in a slightly higher oxidation state than the other. The ligand at a higher oxidation state makes longer Fe-N bonds (1.984(2) and 1.985(2) Å) than the other (1.958(2) and 1.959(2) Å). The Mössbauer spectrum of a solid sample of **9** at 80 K displays two doublets in a 1:1 ratio (δ = 0.66 and 0.54 mm/s; $|\Delta E_q| = 2.81$ and 2.59 mm/s, respectively, Table 2, Figure S17) with large quadruple splitting values. Such Mössbauer data suggest that one iron centre is in a slightly higher oxidation state than the other and the valence is not fully delocalized on Mössbauer time

Table 1. Selected bond distances of complexes.



| | 2 | 4 | 5 | 6 | 7 | 8 | 8 (DFT) | 9 | 10 | 11 | 12 |
|---|----------------------|----------------------|----------------------|----------------------|----------|--------------------------|---|----------------|----------|-----------|-----------|
| Fe-N | 1.979(2) 1.977(1) | 1.934(2) 1.941(2) | 1.995(3) 2.018(3) | 2.018(2) 2.003(2) | 1.967(2) | 1.950(2) 1 1.973(2) 1 | | 1.984(2) | 1.995(2) | 1.992(2) | 2.015(3) |
| | | | | | | | 1.931 | 1.931 1.985(2) | | 1.982(2) | 2.016(3) |
| | | | | | | | 1.960 | 1.958(2) | | 1.961(2) | 1.955(3) |
| | | | | | | | | 1.959(2) | | 1.957(2) | 1.952(3) |
| N–C _a | 1.341(2) 1.346(2) | 1.392(3) 1.399(3) | 1.345(5) 1.333(5) | 1.351(3) 1.355(3) | 1.328(3) | 1.372(3) 1.376(3) | 1.377 1.380 | 1.370(3) | 1.341(3) | 1.368(3) | 1.338(5) |
| | | | | | | | | 1.374(3) | | 1.364(4) | 1.335(5) |
| | | | | | | | | 1.394(3) | | 1.393(4) | 1.379(5) |
| | | | | | | | | 1.395(3) | | 1.375(3) | 1.378(5) |
| C_{α} – C_{α} ' | 1 452(2) | 1.430(3) | 1.442(5) | 1.445(3) | 1.464(3) | 1.437(3) | 1.439 | 1.443(4) | 1.453(4) | 1.448(4) | 1.457(6) |
| | 1.452(2) | | | | | | | 1.430(4) | | 1.435(4) | 1.433(6) |
| C_{α} – C_{β} | 1.419(2) 1.421(2) | 1.388(4) 1.389(3) | 1.425(6) 1.431(6) | 1.419(3) 1.413(3) | 1.419(3) | | | 1.414(3) | 1.416(4) | 1.413(4) | 1.414(6) |
| | | | | | | 1.406(3) 1.398(3) | 1.416 1.413 | 1.411(4) | | 1.400(4) | 1.423(6) |
| | | | | | | | | 1.407(4) | | 1.390(4) | 1.397(6) |
| | | | | | | | | 1.406(3) | | 1.409(4) | 1.391(5) |
| C _β – C _γ | 1.363(3) 1.361(3) | 1.397(4) 1.393(4) | 1.367(6) 1.355(6) | 1.366(3) 1.367(4) | 1.356(3) | | | 1.373(4) | 1.356(4) | 1.362(4) | 1.366(6) |
| | | | | | | 1.373(4) | .373(4) 1.389 1.378 .373(3) 1.389 1.381 1.375 | 1.378(4) | | 1.369(4) | 1.342(7) |
| | | | | | | 1.373(3) | | 1.381(3) | | 1.383(4) | 1.375(6) |
| | | | | | | | | 1.375(4) | | 1.377(4) | 1.381(6) |
| $C_{\gamma} - C_{\gamma}'$ | 1.405(3) | 1.367(4) | 1.401(7) | 1.408(4) | 1.430(3) | 1.398(4) | 1.408 | 1.401(4) | 1.414(6) | 1.403(4) | 1.403(7) |
| | | | | | | | | 1.391(4) | | 1.398(4) | 1.395(7) |
| Fe–Fe | 3.0255(4) | - | - | - | - | - | - | 2.9729(6) | - | 2.9246(5) | 3.2620(8) |
| | | | | | | | | | | | |

scale, consistent with the crystallographic data. Unfortunately, the Mössbauer data are in the range that does not allow for an unambiguous assignment of the oxidation state and spin state of the iron centres.¹¹

Obtaining a dinuclear structure with four-coordinate iron centres when reacting complex 2 with tert-butoxide, we turned to a bulkier anionic HMDS ligand. The addition of one equivalent of LiHMDS to 2 in a toluene solution at room temperature affords the three-coordinate complex 10 (Scheme 5). The solidstate molecular structure has a C_2 symmetry, with a trigonal planar geometry at iron (Figure 5). The metric parameters (Table 1) suggest a Dipppda*- ligand coordinated to the iron centre. The zero-field Mössbauer spectrum of a solid sample of **10** shows an unsymmetrical doublet with $\delta = 0.48$ mm/s and $|\Delta E_Q| = 1.25$ mm/s (Table 2, Figure S18). The isomer shift and quadruple splitting are typical for a three-coordinate high-spin ferrous center.¹⁷ The unsymmetrical broadening is presumably due to intermediate rates of spin relaxation.11 The observed solution magnetic moment of **10** is 4.1 $\mu_{\rm B}$ at 25 °C in C₆D₆ (Table 2), suggesting a strong antiferromagnetic coupling between the radical ligand and a high-spin Fe^{II} centre.

ARTICLE

Exposing a toluene solution of **10** to 1 atm of CO₂ at room temperature results in the formation of TMS–NCO in a quantitative spectroscopic yield (with respect to complex **10**) within 30 minutes along with a mixture of new paramagnetic species, from which complex **11** can be isolated via recrystallization (Scheme 5). Metal silylamides have been reported to react with CO₂ to give isocyanates,^{18–20} carbodiimides²¹ and metal isocyanato complexes²² along with the corresponding silyl ethers, metal siloxides or unidentified species. However, only a few reports show high selectivity towards the isocyanate under mild conditions,¹⁹ as the resulting metal silyloxide species can further react with CO₂ or isocyanate product (isoelectronic with CO₂) to carbodiimide or metal isocyanate and silyl ether.

In the solid-state, complex **11** has a dinuclear structure with two bridging trimethylsiloxide ligands between two ^{Dipp}pda–Fe fragments (Figure 5), resembling the structure of **9**. One of the ^{Dipp}pda ligands in **11** displays two C_a–N distances of **1.368(3)** and **1.364(4)** Å, respectively, whereas the other displays two C_a–N distances of **1.393(4)** and **1.375(3)** Å, respectively (Table 1). The trend in the Fe–N bond lengths of **11** is also similar to that of **9**. The observed solution magnetic moment 5.9 μ_B at 25 °C in C₆D₆ (Table 2). The Mössbauer spectrum of **11** at 80 K displays similar pattern as that of **9** (Table 2, Figure S19), suggesting a slight difference in the oxidation states of the two iron centres. Once again, the Mössbauer data for **11** are in the range that does not allow for an unambiguous assignment of the oxidation state and spin state of the iron centres.¹¹

When a saturated hexanes solution of 10 is exposed to 1 atm. of CO₂ at -35 °C, a few crystals of complex **12** were obtained (Scheme 5). The molecular structure of 12 features two iron centres bridged by a trimethylsilyl carbonate and a trimethylsiloxide. One Dipppda ligand in complex 12 displays two C_{α} -N distances of 1.338(5) and 1.335(5) Å, respectively, whereas the other 1.379(5) and 1.378(5) Å, respectively. The Fe-N bond lengths in 12 also have the same pattern as in 9 and 11. A better quality X-ray structure is needed to elucidate the oxidation state of the Dipppda ligand. Complex 12 has the longest intramolecular Fe-Fe distance (3.2620(8) Å) among all the dinuclear complexes in this work. Unfortunately, complex 12 readily loses CO₂ to form complex 11 under vacuum, which hampers the isolation and further characterizations. The formation of 12 can be rationalized as the insertion of CO₂ into the Fe–O bond of the in situ formed 11. Although isocyanates tend to show similar reactivity as CO₂, we observed no reaction between TMS-NCO and complex 11 produced in the reaction of 10 and CO₂. The lack of reactivity is presumably due to the steric congestion around the iron centres.

| able 2. Sol | ution Magnet | ic Moment | at 298 K and N | 1össbauer data at 8 | 30 K. |
|-------------|--------------|-----------------------|----------------|-------------------------|-------|
| | Complex | $\mu_{	ext{eff}}{}^a$ | δ, mm/s | $ \Delta E_{Q} $, mm/s | _ |
| | 2 | 5.4 | 0.84 | 3.64 | |
| | 3 | - | 0.37 | 0.79 | |
| | 4 | 5.2 | 0.72 | 4.02 | |
| | 5 | 4.2 | 0.81 | 2.05 | |
| | 6 | 4.1 | 0.80 | 3.04 | |
| | 7 | 5.1 | 0.39 | 2.12 | |
| | 8 | 2.7 | 0.42 | 2.51 | |
| | ٥ | 6 1 | 0.66 | 2.81 | |
| | 9 | 0.1 | 0.54 | 2.59 | |

0.55

0.76

0.60

0.79

2.28

2.61

^a Solution magnetic moment by Evans method.

4.1

5.9

10

11

Conclusions

In summary, a series of iron o-phenylenediamide complexes were synthesized. Their electronic structures were investigated through spectroscopic methods, and X-ray crystallography. The dimeric complex 2 contains two high-spin ferrous ions and two ^{Dipp}pda•⁻ π -radical monoanionic ligands with strong antiferromagnetic coupling within each Fe-Dippdpa pair. Complex **3** features a low-spin Fe(I) centre and a ^{Dipp}pda⁻⁻ ligand, where the electronic structure of the Fe^{I_Dipp}dpa⁻⁻ moiety can be manipulated with different auxiliary ligands on iron. On the other hand, the Fe^{II_Dipp}dpa⁻⁻ moiety in complexes **2**, **5** and **6** is persistent with various combinations of chloride and pyridine ligand. Such a phenomenon can be attributed to the low reducing power of the Dippdpa⁻⁻ ligand. This hypothesis is consistent with the fact that 1e⁻ oxidation of complex 6 gives Fe^{III_Dipp}dpa^{•-} (rather than Fe^{II_Dipp}dpa⁰) moiety in **7**. Complex **8** with two bulky Dipppda ligands on one iron centre is tentatively assigned as [Fe^{III}(^{Dipp}pda^{•-})(^{Dipp}pda²⁻)] \leftrightarrow [Fe^{III}(^{Dipp}pda²⁻)(^{Dipp}pda^{•-})], where the is metal antiferromagnetically coupled to the fully delocalized ligandbased radical based on literature precedence and available data. Using a bulky anionic ligand HMDS to replace the bridging chloride in 2, we were able to obtain a mononuclear threecoordinate iron complex 9, which shows reactivity toward CO₂ to give the trimethylsiloxide-bridge dimeric complex 11 and TMS-NCO selectively. Complex 11 can selectively react with CO₂ over the bulkier TMS-NCO in a reversible fashion to give complex 12. Further investigation of the potential catalytic activities of these iron complexes are underway in our laboratories.

ransactions Accepted Manusc

Journal Name

at -35 °C. ¹H NMR (600 MHz, C₆D₆) δ 158.25 (33.53,4H)₂₆67,20 (36.53, 4H), 2.02 (13.48, 24H, Dipp-CH₃), ±59.173(207), 12,28H); -8.44 (8.85, 4H), -8.46 (7.39, 4H), -11.35 (9.73, 4H), -14.63 (28.63, 24H, Dipp-CH₃). Evans method (298 K, C₆D₆): μ_{eff} = 5.4 μ_B. Anal. Calcd for C₆₀H₇₆N₄Cl₂Fe₂·(C₄H₁₀O): C, 69.25; H, 7.81; N, 5.05. Found: C, 68.85; H, 7.48; N, 5.41.

3: To a 20 mL vial containing Hg (1.45 g) and 5 mL of toluene was added sodium metal (7.2 mg, 0.31 mmol). After stirring the resulting amalgam for 30 min, a solution of 2 (147.5 mg, 0.14 mmol) in 5 mL of toluene was added. The reaction mixture was stirred for 24 hours, then decanted and filtered through Celite. The filtrate was concentrated to dryness under vacuum. The solid residue was recrystallized from hexanes to afford metallic green crystals of 3 (132.7 mg, 82%). NMR data obtained are identical to those previously reported.7g

4: To a stirring solution of 3 (287.3 mg, 0.50 mmol) in 10 mL of THF was added pyridine (89 µL, 1.10 mmol). The reaction mixture was allowed to stir at room temperature overnight, and then filtered over Celite. The filtrate was concentrated to ~1 mL, top-layered with 5 mL of pentane and cooled to -35 °C overnight to yield dark blue purple crystals. The supernatant was decanted off, and the solid was washed with cold pentane $(3 \times 1 \text{ mL})$ and then dried under vacuum (212.3 mg, 66%). Crystals suitable for X-ray crystallography were obtained from cooling a concentrated diethyl ether solution at –35 °C. ¹H NMR (600 MHz, THF-d₈) δ 118.39 (3921.27, 2H), 42.13 (31.48, 4H), 29.07 (238.34, 4H), 14.76 (156.00, 4H), 14.05 (710.18, 2H), 5.29 (37.32, 12H, Dipp-CH₃), 0.28 (114.16, 12H, Dipp-CH₃), −17.90 (40.82, 2H), -32.56 (123.98, 2H), -36.47 (26.66, 2H). Evans method (298 K, THF- d_8): μ_{eff} = 5.2 μ B. Anal. Calcd for C₄₀H₄₈N₄Fe·(C₄H₈O): C, 74.14; H, 7.92; N, 7.86. Found: C, 74.52; H, 7.70; N, 8.13.

5: To a stirring solution of 2 (259.0 mg, 0.25 mmol) in 10 mL of toluene was added pyridine (50 µL, 0.63 mmol). The reaction mixture was allowed to stir at room temperature overnight, and then concentrated to dryness under vacuum. The residue was extracted into diethyl ether, filtered over Celite and slowly concentrated to \sim 2 mL. Top-layering with 5 mL of pentane and cooling to -35 °C overnight yielded green microcrystalline solid. The supernatant was decanted off, and the solid was washed with cold pentane $(3 \times 1 \text{ mL})$ and then dried under vacuum (218.4 mg, 73%). ¹H NMR (600 MHz, C₆D₆) δ 162.94 (1558.36, 1H), 104.68 (1394.75, 1H), 59.36 (1711.85, 1H), 34.19 (148.48, 2H), 2.63 (637.74, 12H, Dipp-CH₃), -8.42 (96.38, 1H), -15.24 (290.19, 12H, Dipp-CH₃), -28.15 (684.49, 2H). Evans method (298 K, C_6D_6): μ_{eff} = 4.2 μ_B . Anal. Calcd for $C_{35}H_{43}N_3FeCl$: C, 70.41; H, 7.26; N, 7.04. Found: C, 70.07; H, 7.61; N, 7.20.

6: To a solid mixture of 2 (259.0 mg, 0.25 mmol) and tetrabutylammonium chloride (347.4 mg, 1.25 mmol) was added 10 mL of toluene. The reaction mixture was allowed to stir at room temperature overnight, and then filter through Celite. The filtrate was concentrated to ~1 mL, top-layered with 5 mL of hexanes and cooled to -35 °C overnight to afford blue

Experimental Section

General considerations

All reactions were carried out in a dinitrogen-filled glovebox or using the standard Schlenk techniques under dinitrogen. Glassware was dried in a 180 °C oven overnight. Diethyl ether, hexanes, pentane, and toluene solvents were dried by a Grubbs-type solvent purification system manufactured by Innovative Technology and degassed prior to use. THF solvent was dried by refluxing and distilling over sodium benzophenone ketyl under dinitrogen. Pyridine and hexamethyldisiloxane (HMDSO) were dried by refluxing and distilling over calcium hydride under dinitrogen. C₆D₆, and THF-d₈ were degassed through three consecutive freeze-pump-thaw cycles. All solvents were stored over 3 Å molecular sieves prior to use. Unless otherwise noted, all NMR spectra were recorded on an Agilent DD2 600 MHz spectrometer at 25 °C. Chemical shifts are referenced to the solvent signals. Solution magnetic moments were measured at 25 °C by the method originally described by Evans with stock and experimental solutions containing a known amount of a cyclohexane standard.²³ Elemental analyses were carried out by ANALEST at the University of Toronto. Electronic spectra were recorded with a Perkin-Elmer doublebeam UV-VIS-NIR spectrometer Lambda 1050. All commercially available chemicals were used as received. Fe(HMDS)₂Cl(THF),²⁴ and 1²⁵ were prepared according to literature procedures. The ¹H NMR data are reported as chemical shift with the peak width at half-height in Hz, integration value, and partial assignment based on integration given in parentheses sequentially.

57Fe Mössbauer Spectroscopy. All measurements for 57Fe Mössbauer spectroscopy were performed using non-enriched solids of the as-isolated complexes. All samples were prepared in an inert atmosphere glovebox equipped with a liquid nitrogen fill port to enable sample freezing to 77 K within the glovebox. Each sample was loaded into a Delrin Mössbauer sample cup for measurements and loaded under liquid nitrogen. Low temperature ⁵⁷Fe Mössbauer measurements were performed using a See Co. MS4 Mössbauer spectrometer integrated with a Janis SVT-400T He/N₂ cryostat for measurements at 80 K. Isomer shifts were determined relative to α -Fe at 298 K. All Mössbauer spectra were fit using the program WMoss (SeeCo). Errors of the fit analyses were the following: $\delta \pm 0.02$ mm/s and $\Delta E_Q \pm 3\%$. For multicomponent fits, the quantitation errors were ± 3% (e.g., 50 ± 3%).

2: To the mixture of 1 (857.3 mg, 2.00 mmol) and Fe(HMDS)₂Cl(THF) (968.3 mg, 2.00 mmol) was added 10 mL of toluene. The reaction mixture was allowed to stir at 120 °C overnight. The reaction mixture changed from dark red to dark green. After cooling to room temperature, the reaction mixture was filtered through Celite. Volatiles were removed under vacuum, leaving a dark green solid. The solid was stirred with 10 mL of hexanes for 1 hour then collected on a frit, which was washed with hexanes $(3 \times 1 \text{ mL})$ and dried under high vacuum (882.4 mg, 85%). Crystals suitable for X-ray crystallography were obtained by cooling a concentrated diethyl ether solution

Journal Name

purple crystals. The supernatant was decanted off, and the solid was washed with hexanes (3 × 1 mL) and then dried under vacuum (319.2 mg, 80%). Crystals suitable for X-ray crystallography were obtained from vapor diffusion of pentane into its toluene solution at room temperature. ¹H NMR (600 MHz, C₆D₆) δ 161.99 (165.36, 1H), 70.63 (138.71, 1H), 12.70 (103.26, 8H), 7.37 (57.54, 8H), 2.44 (43.08, 8H), 1.14 (30.45, 12H), 0.58 (20.98, 12H, Dipp-CH₃), 0.05 (34.41, 8H), -7.76 (69.52, 12H, Dipp-CH₃), -25.24 (22.36, 2H). Evans method (298 K, C₆D₆): μ_{eff} = 4.1 μ_{B} . Anal. Calcd for C₄₆H₇₄N₃Cl₂Fe: C, 69.42; H, 9.37; N, 5.28. Found: C, 69.82; H, 9.62; N, 5.17.

7: In the absence of light, silver tetrafluoroborate (53.5 mg, 0.28 mmol) was added to a stirring solution of 6 (199.0 mg, 0.25 mmol) in 10 mL of toluene. The reaction mixture was allowed to stir at room temperature overnight, and then filter through Celite. The filtrate was concentrated to dryness. The residue was dissolved in toluene and filtered through Celite again, then concentrated \sim 1 mL, top-layered with 5 mL of pentane and cooled to -35 °C overnight to afford dark green microcrystalline solid. The supernatant was decanted off, and the solid was washed with pentane $(3 \times 1 \text{ mL})$ and then dried under vacuum (99.2 mg, 72%). Crystals suitable for X-ray crystallography were obtained from cooling a concentrated diethyl ether solution at -35 °C. ¹H NMR (600 MHz, C₆D₆) δ 218.94 (1083.40, 1H), 17.12 (175.98, 4H), 6.85 (570.77, 2H, overlap with C₆D₆), 0.95 (566.66, 12H, Dipp-CH₃), -1.30 (212.26, 12H, Dipp-CH₃), -37.18 (176.00, 2H). Evans method (298 K, C₆D₆): μ_{eff} = 5.1 μ_{B} . Anal. Calcd for C₃₀H₃₈N₂Cl₂Fe·0.6(C₇H₈): C, 67.49; H, 7.09; N, 4.60. Found: C, 67.55; H, 7.10; N, 4.66.

8: To a solution of 1 (535.8 mg, 1.25 mmol) in 5 mL of THF was slowly added "BuLi (2.5 M in hexanes, 1.00 mL, 2.50 mmol) at -80 °C. The resulting mixture was allowed to warm to room temperature slowly and further stirred for 2 h. The resulting suspension was cooled to -80 °C and a pre-cooled (-80 °C) slurry of anhydrous FeCl₂ (79.2 mg, 0.63 mmol, in 10 mL of THF) was added. The resulting mixture was allowed to warm to room temperature slowly and further stirred for 2 h, during which time the colour changed to dark purple. The mixture was cooled to -35 °C and a pre-cooled (-35 °C) solution of FeCl₃ (202.8 mg, 1.25 mmol, in 10 mL of THF) was added. The reaction mixture was stirred overnight, with a colour changed from dark purple to dark green. All volatiles were removed under vacuum, the residue was extracted into pentane and filtered through Celite. The filtrated was concentrated to ~5 mL, top-layered with 3 mL of HMDSO and cooled to -35 °C overnight to yield X-ray quality dark green crystals. The supernatant was decanted off, and the solid was washed with HMDSO (3 \times 1 mL) and then dried under vacuum (445.6 mg, 78%). ^1H NMR (600 MHz, $C_6D_6)$ δ 66.12 (117.84, 4H), 22.44 (36.62, 4H), 14.40 (25.23, 4H), 9.21 (223.83, 4H), 6.15 (531.51, 2H), 4.27 (23.57, 12H, Dipp-CH₃), 3.92 (19.07, 12H, Dipp-CH₃), -0.63 (22.96, 12H, Dipp-CH₃), -2.82 (46.64, 12H, Dipp-CH₃), -3.70 (124,70, 4H), -12.45 (35.67, 4H). Evans method (298 K, C_6D_6): μ_{eff} = 2.7 μ_B . Anal. Calcd for C₆₀H₇₆N₄Fe·0.13(C₆H₁₈Si₂O): C, 78.48; H, 8.49; N, 6.02. Found: C, 78.89; H, 8.65; N, 5.61.

ARTICLE

View Article Online

9: To a stirring solution of **2** (259.0 mg, 0.25^{1} MMO) PPTORMER was added KO^tBu (56.1 mg, 0.50 mmol) in toluene. The reaction mixture was allowed to stir at room temperature overnight, and then filtered through Celite. The filtrate was concentrated to ~1 mL, top-layered with 3 mL of pentane and cooled to -35 °C overnight to yield X-ray quality dark blue purple crystals. which was washed with pentane (5 × 1 mL) and dried under high vacuum (162.2 mg, 58%). ¹H NMR (600 MHz, C₆D₆) δ 45.56 (277.82, 4H), 40.71 (268.15, 4H), 26.80 (522.61, 4H), 13.94 (282.00, 12H, Dipp-CH₃), 8.72 (442.06, 18H, OC(CH₃)₃), 3.54 (311.05, 12H), 1.26 (135.57, 12H, Dipp-CH₃), -4.40 (909.99, 2H), -9.30 (314.89, 12H, Dipp-CH₃), -19.68 (89.56, 4H), -31.60(50.38, 4H), -35.93 (136.64, 4H). Evans method (298 K, C₆D₆): $\mu_{eff} = 6.1 \ \mu_{B}$. Anal. Calcd for C₆₈H₉₄N₄O₂Fe₂·(C₇H₈): C, 74.86; H, 8.54; N, 4.66. Found: C, 74.74; H, 8.56; N, 4.56.

10: To a stirring solution of 2 (259.0 mg, 0.25 mmol) in toluene was added LiHMDS (83.7 mg, 0.5mmol) in toluene. The reaction mixture was allowed to stir at room temperature overnight, and then concentrated to dryness under vacuum. The residue was extracted into pentane, filtered over Celite and slowly concentrated to dryness to afford a green brown microcrystalline solid, which was washed with cold pentane (3 × 0.5 mL) and dried under high vacuum (266.0 mg, 83%). Crystals suitable for X-ray crystallography were obtained from top-layering the pentane solution with HMDSO at -35 °C. ¹H NMR (600 MHz, C₆D₆) δ 266.68 (450.23, 1H), 130.82 (406.15, 2H), 11.91 (403.82, 18H, HMDS-CH₃), 2.64 (105.35, 12H, Dipp-CH3), -20.20 (24.83, 2H), -27.94 (44.09, 4H), -38.61 (145.59, 12H), -88.02 (1824.32, 2H). Evans method (298 K, C_6D_6): μ_{eff} = 4.1 μ_B. Anal. Calcd for C₃₆H₅₆N₃Si₂Fe: C, 67.26; H, 8.78; N, 6.54. Found: C, 66.70; H, 8.81; N, 6.44.

11: A solution of 10 (160.7 mg, 0.25 mmol) in 10 mL of toluene was subjected to a freeze-pump-thaw cycle before 1 atm of CO₂ was introduced into the flask. The mixture was allowed to stir for 1 hour at room temperature, where the colour changed from green brown to dark blue. The reaction mixture was filtered through Celite. The filtrate was concentrated to ~1 mL, top-layered with 3 mL of pentane and cooled to -35 °C overnight to yield dark blue purple crystalline solid, which was washed with pentane $(5 \times 1 \text{ mL})$ and dried under high vacuum (89.8 mg, 63%). Crystals suitable for X-ray crystallography were obtained from cooling a concentrated diethyl ether solution at -35 °C. ¹H NMR (600 MHz, C₆D₆) δ 49.40 (279.92, 4H), 34.03 (43.46, 4H), 28.04 (42.69, 4H), 23.95 (69.80, 4H), 19.92 (43.02, 12H, Dipp-CH₃), -0.50 (47.15, 12H, Dipp-CH₃), -6.97 (7.63, 4H), -12.00 (390.53, 18H, OSi(CH₃)₃), -21.34 (74.98, 12H, Dipp-CH₃), -27.06 (30.93, 4H), -31.35 (283.53, 4H). Evans method (298 K, C_6D_6): $\mu_{eff} = 5.9 \,\mu_B$. Anal. Calcd for $C_{66}H_{94}N_4O_2Si_2Fe_2$: C, 69.33; H, 8.29; N, 4.90. Found: C, 69.44; H, 8.78; N, 4.63.

Conflicts of interest

There are no conflicts to declare.

Journal Name

Acknowledgements

We thank Natural Science and Engineering Research Council (NSERC) of Canada (grants to D.S.) and the National Science Foundation (CHE-1954480 to M.L.N.) for funding. Q.L. thanks the Ontario government for an Ontario Graduate Scholarship. J.H.L thanks NSERC of Canada for a USRA. We also acknowledge the Ontario Research Fund for funding the CSICOMP NMR lab at the University of Toronto enabling the purchase of several new NMR spectrometers.

Notes and references

- 1 (a) A. Chirila, B. G. Das, P. F. Kuijpers, V. Sinha and B. de Bruin, in Non-Noble Metal Catalysis: Molecular Approaches and Reactions, ed. R. J. M. K. Gebbink and M.-E. Moret, Wiley-VCH, Weinheim, Germany, 2019, p. 1-31. (b) B. de Bruin, P. Gualco and N. D. Paul, in Ligand Design in Metal Chemistry, ed. M. Stradiotto and R. J. Lundgren, John Wiley & Sons, Ltd, Chichester, UK, 2016, p. 176-204; (c) J. I. van der Vlugt Chem. Eur. J., 2019, 25, 2651; (d) J. Jacquet, M. Desage-El Murr and L. Fensterbank, ChemCatChem, 2016, 8, 3310; (e) A. I. Olivos Suarez, V. Lyaskovskyy, J. N. H. Reek, J. I. van der Vlugt and B. de Bruin, Angew. Chem. Int. Ed., 2013, 52, 12510; (f) O. R. Luca and R. H. Crabtree, Chem. Soc. Rev., 2013, 42, 1440; (h) R. F. Munhá, R. A. Zarkesh and A. F. Heyduk, Dalton Trans., 2013, 42, 3751; (o) V. Lyaskovskyy and B. de Bruin, ACS Catal., 2012, 2, 270; (j) K. G. Caulton, Eur. J. Inorg. Chem., 2012, 435. (k) V. K. K. Praneeth, M. R. Ringenberg and T. R. Ward, Angew. Chem. Int. Ed., 2012, 51, 10228; (I) W. Kaim, Eur. J. Inorg. Chem., 2012, 343; (m) P. J. Chirik, Inorg. Chem., 2011, 50, 9737; (n) P. J. Chirik and K. Wieghardt, Science, 2010, 327, 794.
- 2 (a) S. M. Rummelt, H. Zhong, I. Korobkov and P. J. Chirik J. Am. Chem. Soc., 2018, 140, 11589; (b) M. J. T. Wilding, D. A. Iovan, A. T. Wrobel, J. T. Lukens, S. N. MacMillan, K. M. Lancaster and T. A. Betley, J. Am. Chem. Soc., 2017, 139, 14757; (c) W. J. T. Wilding, D. A. Iovan and T. A. Betley, J. Am. Chem. Soc., 2017, 139. 12043.
- 3 (a) J. M. Hoyt, V. A. Schmidt, A. M. Tondreau and P. J. Chirik Science, 2015, 349, 960; (b) A. M. Tondreau, C. C. H. Atienza, K. J. Weller, S. A. Nye, K. M. Lewis, J. G. P. Delis and P. J. Chirik, Science, 2012, 335, 567. (c) D. A. Iovan, M. J. T. Wilding, Y. Baek, E. T. Hennessy and T. A. Betley, Angew. Chem. Int. Ed., 2017, 56, 15599; (d) E. T. Hennessy and T. A. Betley, Science, 2013, 340, 591.
- (a) C. G. Pierpont, Coord. Chem. Rev., 2001, 216-217, 99; (b) 4 D. L. J. Broere, R. Plessius and J. I. van der Vlugt, Chem. Soc. Rev., 2015, 44, 6886.
- (a) M. V. Joannou, M. J. Bezdek, K. Albahily, I. Korobkov and P. 5 J. Chirik Organometallics 2018, **37**, 3389; (b) M. J. Sgro and D. W. Stephan Dalton Trans., 2010, 39, 5786; (c) W. W. Kramer, L. A. Cameron, R. A. Zarkesh, J. W. Ziller and A. F. Heyduk Inorg. Chem. 2014, 53, 8825; (d) N. G. Léonard and P. J. Chirik ACS Catal., 2018, 8, 342; (e) W. N. Palmer and P. J. Chirik ACS Catal., 2017, 7, 5674; (f) L. Pei, F. Liu , H. Liao, J. Gao, L. Zhong, H. Gao and Q. Wu ACS Catal., 2018, 8, 1104; (g) X.-J. Yang, X. Fan, Y. Zhao, X. Wang, B. Liu, J.-H. Su, Q. Dong, M. Xu and B. Wu Organometallics, 2013, 32, 6945; (h) K. Y. Monakhov, J. van Leusen, P. Kögerler, E.-L. Zins, M. E. Alikhani, M. Tromp, A. A. Danopoulos and P. Braunstein Chem. Eur. J., 2017, 23, 6504; (i) M. van der Meer, Y. Rechkemmer, I. Peremykin, S. Hohloch, J. van Slageren and B. Sarkar Chem. Commun., 2014, 50, 1110; (j) H. Nishiyama, H. Ikeda, T. Saito, B. Kriegel, H. Tsurugi, J. Arnold and K. Mashima J. Am. Chem. Soc., 2017, 139, 6494; (k) J. Bendix and K. M. Clark, Angew. Chem. Int. Ed., 2016, 55, 2748; (I) T. Janes, M. Xu and D. Song, Dalton Trans., 2016, 45, 10672.

- 6 (a) D. A. Evans and A. H. Cowley, J. Am. Chem. Soc., 2012, 134, 15672; (b) D. A. Evans, I. Vargas-Baca and AoHo Cowley 2020 Am Chem. Soc., 2013, 135, 13939; (c) I. L. Fedushkin, V. A. Dodonov, A. A. Skatova, V. G. Sokolov, A. V. Piskunov and G. K. Fukin, Chem. Eur. J., 2018, 24, 1877; (d) K. V. Vasudevan, M. Findlater, I. Vargas-Baca and A. H. Cowley J. Am. Chem. Soc., 2012, 134, 176; (e) Y. Segawa, Y. Suzuki, M. Yamashita and K. Nozaki, J. Am. Chem. Soc., 2008, 130, 16069; (f) K. L. Bamford, L. E. Longobardi, L. Liu, S. Grimme and D. W. Stephan, Dalton Trans., 2017, 46, 5308; (g) W. Zhang, V. A. Dodonov, W. Chen, Y. Zhao, A. A. Skatova, I. L. Fedushkin, P. W. Roesky, B. Wu and X.-J. Yang, Chem. Eur. J., 2018, 24, 14994; (h) T. Janes, P. Zatsepin and D. Song, Chem. Commun., 2017, 53, 3090; (n) M. Ma, L. Shen, H. Wang, Y. Zhao, B. Wu and X.-J. Yang, Organometallics, 2020, 39, 1440; (m) Y. Liu, S. Li, X.-J, Yang, P. Yang and B. Wu, J. Am. Chem. Soc., 2009, 131, 4210.
- (a) S. C. Bart, E. J. Hawrelak, A. K. Schmisseur, E. Lobkovsky and P. J. Chirik, Organometallics, 2004, 23, 237; (b) N. Muresan, C. C. Lu, M. Ghosh, J. C. Peters, M. Abe, L. M. Henling, T. Weyhermöller, E. Bill and K. Wieghardt, Inorg. Chem., 2008, 47, 4579; (c) M. M. Khusniyarov, E. Bill, T. Weyhermüller, E. Bothe, K. Harms, J. Sundermeyer and K. Wieghardt, Chem. Eur. J., 2008, 14, 7608; (d) M. M. Khusniyarov, T. Weyhermüller, E. Bill and K. Wieghardt, J. Am. Chem. Soc., 2009, 131, 1208; (e) M. M. Khusniyarov, T. Weyhermüller, E. Bill and K. Wieghardt, Angew. Chem. Int. Ed., 2008, 47, 1228; (f) A. Hernán-Gámez, M. Rodríguez, T. Parella and M. Costas, Angew. Chem. Int. Ed., 2019, 58, 13904; (g) T. Janes, J. M. Rawson and D. Song, Dalton Trans., 2013, 42, 10640.
- 8 (a) S. C. Bart, E. J. Hawrelak, E. Lobkovsky and P. J. Chirik, Organometallics, 2005, 24, 5518; (b) V. A. Schmidt, C. R. Kennedy, M. J. Bezdek and P. J. Chirik, J. Am. Chem. Soc., 2018, 140, 3443; (c) H. tom Dieck and H. Bruder, J. Chem. Soc., Chem. Commun., 1977, 24; (d) H. tom Dieck and J. Dietrich, Angew. Chem. Int. Ed., 1985, 24, 781; (e) P. Le Floch, F. Knoch, F. Kremer, F. Mathey, J. Scholz, W. Scholz, K.-H. Thiele and U. Zenneck, Eur. J. Inorg. Chem., 1998, 119; (f) H. tom Dieck and R. Diercks, Angew. Chem. Int. Ed., 1983, 22, 778; (g) C. Lichtenberg, M. Adelhardt, T. L. Gianetti, K. Meyer, B. de Bruin and H. Grützmacher, ACS Catal., 2015, 5, 6230; (h) H. Lee, M. G. Campbell, R. H. Sánchez, J. Börgel, J. Raynaud, S. E. Parker and T. Ritter, Organometallics, 2016, 35, 2923; (i) R. K. O'Reilly, M. P. Shaver, V. C. Gibson and A. J. P. White, Macromolecules, 2007, 40, 7441; (j) C.-H. Ke, C.-H. Chen, M.-L. Tsai, H.-C. Wang, F.-T. Tsai, Y.-W. Chiang, W.-C. Shih, D. S. Bohle and W.-F. Liaw, J. Am. Chem. Soc., 2017, 139, 67. 9
 - (a) M. Villa, D. Miesel, A. Hildebrandt, F. Ra-gaini, D. Schaarschmidt and A. Jacobi von Wangelin, ChemCatChem 2017, 9, 3203; (b) A. Saini, C. R. Smith, F. S. Wekesa, A. K. Helms and M. Findlater Org. Biomol. Chem., 2018, 16, 9368; (c) X. Yu, F. Zhu, D. Bu and H. Lei RSC Adv., 2017, 7, 15321; (d) V. V. Khrizanforova, V. I. Morozov, M. N. Khrizanforov, A. N. Lukoyanov, O. N. Kataeva, I. L. Fedushkin and Yu. H. Budnikova, Polyhedron, 2018, 154, 77; (e) P. J. Larson, F. S. Wekesa, A. Singh, C. R. Smith, A. Rajput, G. P. McGovern, D. K. Unruh, A. F. Cozzolino and M. Findlater, Polyhedron, 2018, 159, 365; (f) A. Paulovicova, U. El-Ayaan, K. Umezawa, C. Vithana, Y. Ohashi and Y. Fukuda, Inorg. Chim. Acta., 2002, 339, 209; (g) D. A. Piryazev, M. A. Ogienko, A. V. Virovets, N. A. Pushkarevsky and S. N. Konchenko, Acta Crystallogr. Sect. C 2012, 68, m320; (h) I. L. Fedushkin, A. A. Skatova, N. M. Khvoinova, A. N. Lukoyanov, G. K. Fukin, S. Y. Ketkov, M. O. Maslov, A. S. Bogomyakov and V. M. Makarov, Russ. Chem. Bull., 2013, 62, 2122; (i) M. Schmitz, M. Seibel, H. Kelm, S. Demeshko, F. Meyer and H.-J. Krüger, Angew. Chem. Int. Ed., 2014, 53, 5988; (j) F. S. Wekesa, R. Arias-Ugarte, L. Kong, Z. Sumner, G. P. McGovern and M. Findlater, Organometallics,

10 | J. Name., 2012, 00, 1-3

This journal is C The Royal Society of Chemistry 20xx

Published on 17 August 2020. Downloaded on 8/18/2020 1:41:39 AM

Journal Name

2015, **34**, 5051; (k) M. J. Supej, A. Volkov, L. Darko, R. A. West, J. M. Darmon, C. E. Schulz, K. A. Wheeler and H. M. Hoyt, *Polyhedron*, 2016, **114**, 403; (l) L. A. Brown, F. S. Wekesa, D. K. Unruh, M. Findlater and B. K. J. Long, *Polym. Sci., Part A: Polym. Chem.*, 2017, **55**, 2824.

- W. P. Fehlhammer, H. Stolzenberg, in *Comprehensive* Organometallic Chemistry, ed. G. Wilkinson, F. G. A. Stone, F. W. Abel, Pergamon Press, Oxford, UK, 1982, p. 515-524.
- 11 P. Gütlich, E. Bill and A. X. Trautwein, *Mössbauer Spectroscopy* and *Transition Metal Chemistry. Fundamentals and Applications*, Springer, Berlin, Germany, 2011.
- 12 P. J. Hay, J. C. Thibeault, R. Hoffmann, J. Am. Chem. Soc. 1975, 97, 4884.
- 13 J. P. Perdew, K. Burke, M. Ernzerhof, *Phys. Rev. Lett.* 1996, **77**, 3865.
- 14 A. Schäfer, C. Huber, R. Ahlrichs, J. Chem. Phys. 1994, 100, 5829.
- 15 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2016.
- 16 F. Weigend, R. Ahlrichs, *Phys. Chem. Chem. Phys.* 2005, 7, 3297.
- 17 (a) H. Andres, E. L. Bominaar, J. M. Smith, N. A. Eckert, P. L. Holland, E. Münck, *J. Am. Chem. Soc.* 2002, *124*, 3012; (b) S. Yogendra, T. Weyhermüller, A. W. Hahn, S. DeBeer, *Inorg. Chem.* 2019, *58*, 9358.
- (a) M. Reiter, S. Vagin, A. Kronast, C. Jandl, B. Rieger, *Chem. Sci.* 2017, 8, 1876; (b) D. R. Moore, M. Cheng, E. B. Lobkovsky, G. W. Coates, *J. Am. Chem. Soc.* 2003, 125, 11911.
- (a) D. A. Dickie, K. B. Gislason, R. A. Kemp, *Inorg. Chem.* 2012, **51**, 1162; (b) A. M. Felix, B. J. Boro, D. A. Dickie, Y. Tang, J. A. Saria, B. Moasser, C. A. Stewart, B. J. Frost, R. A. Kemp, *Main Group Chem.* 2012, **11**, 13; (c) M. Xu, A. R. Jupp, D. W. Stephan, *Angew. Chem. Int. Ed.*, 2017, **56**, 14277; (d) D. L. J. Broere, B. Q. Mercado, P. L. Holland, *Angew. Chem. Int. Ed.*, 2018, **57**, 6507.
- 20 H. Yin, P. J. Carroll, E. J. Schelter, *Chem. Commun.* 2016, **52**, 9813.
- 21 L. R. Sita, J. R. Babcock, R. Xi, J. Am. Chem. Soc. 1996, **118**, 10912.
- 22 (a) C. Camp, L. Chatelain, C. E. Kefalidis, J. Pécaut, L. Maron, M. Mazzanti, *Chem. Commun.* 2015, **51**, 15454; (b) M. T. Whited, A. J. Kosanovich, D. E. Janzen, *Organometallics* 2014, **33**, 1416; (c) P. Arnold, Z. R. Turner, A. I. Germeroth, I. J. Casely, G. S. Nichol, R. Bellabarba, R. P. Tooze, *Dalton Trans.* 2013, **42**, 1333; (d) W. Sattler, G. Parkin, *J. Am. Chem. Soc.* 2011, **133**, 9708; (e) B. C. Fullmer, H. Fan, M. Pink, K. G. Caulton, *Inorg. Chem.* 2008, **47**, 1865; (f) H. Phull, D. Alberti, A. L. Korobkov, S. Gambarotta, P. H. M. Budzelaar, *Angew. Chem. Int. Ed.* 2006, **45**, 5331.
- 23 D. F. Evans, J. Chem. Soc., 1959, 2003.
- 24 J. S. Duncan, T. M. Nazif, A. K. Verma and S. C. Lee, *Inorg. Chem.*, 2003, **42**, 1211.

25 T. Wenderski, K. M. Light, D. Ogrin, S. G. Bott and C. J. Harlan, *Tetrahedron Lett.*, 2004, 45, 6851. DOI: 10.1039/D0DT02087G

J. Name., 2013, **00**, 1-3 | **11**



We report the reactivity of the iron complexes of a bulky phenylenediamide ligand.