



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcyc20>

Synthesis of N-Methyl Secondary Amines

Hephzibah J. Kumpaty ^a, John S. Williamson ^b & Sukanta Bhattacharyya ^c

^a Department of Chemistry, University of Wisconsin-Whitewater, Wisconsin, USA

^b Department of Medicinal Chemistry, School of Pharmacy, The University of Mississippi, Mississippi, USA

^c Argonaut Technologies, Foster City, California, USA

Published online: 15 Aug 2006.

To cite this article: Hephzibah J. Kumpaty, John S. Williamson & Sukanta Bhattacharyya (2003) Synthesis of N-Methyl Secondary Amines, *Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry*, 33:8, 1411-1416, DOI: [10.1081/SCC-120018703](https://doi.org/10.1081/SCC-120018703)

To link to this article: <http://dx.doi.org/10.1081/SCC-120018703>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>



SYNTHETIC COMMUNICATIONS®

Vol. 33, No. 8, pp. 1411–1416, 2003

Synthesis of *N*-Methyl Secondary Amines

Hephzibah J. Kumpaty,¹ John S. Williamson,² and
Sukanta Bhattacharyya^{3,*}

¹Department of Chemistry, University of
Wisconsin–Whitewater, Wisconsin, USA

²Department of Medicinal Chemistry,
School of Pharmacy, The University of Mississippi,
Mississippi, USA

³Argonaut Technologies, Foster City, California, USA

ABSTRACT

A diverse set of *N*-methyl secondary amines are obtained in high yields by an expedient reductive alkylation of commercially available methanolic methylamine.

Key Words: Reductive alkylation; Methylamine; Titanium(IV) isopropoxide; Sodium borohydride.


*Correspondence: Sukanta Bhattacharyya, Argonaut Technologies, 1101 Chess Drive, Foster City, CA 94404, USA; Fax: +650-655-4329; E-mail: sbhattacharyya@argotech.com.



Amines and their derivatives are proven synthetic targets because of their versatile applications.^[1] A recent report^[2] claims that about 25% of the registered drugs are based on amines. In connection with our ongoing studies^[3] on reductive amination reactions,^[4] we have earlier described^[5] a method for the preparation of *N*-methyl secondary amines by reductive amination of carbonyl compounds using a combination of titanium(IV) isopropoxide and sodium borohydride. A mixture of methylamine hydrochloride and triethylamine has been employed as the source of nucleophilic methylamine. The method conveniently avoids the use of gaseous methylamine, however, the overall reaction was slow.

As a sequel to this work, we reasoned that a more convenient transformation might be achieved using a commercially available solution of methylamine in methanol. We report herein the results of this investigation leading to a rapid, high-yielding protocol for the preparation of *N*-methyl secondary amines. The efficacy of the protocol has been evaluated on a varied set of carbonyl compounds and the results were compared with our reported^[5] data. The starting aldehydes and ketones were reacted with a commercially available methanol solution of methylamine (2 M) for 5–6 h in the presence of titanium(IV) isopropoxide, followed by the addition of solid sodium borohydride. The reaction mixture was allowed to stir for a further period of 2 h and quenched with water. The reaction is possibly proceeding through the formation of the titanium(IV) complex **1** (Sch. 1) as an intermediate,^[5] which is reduced either directly or via transient iminium species. Titanium(IV) isopropoxide has been utilized^[5,6] as a mild reagent compatible with a variety of potentially acid-sensitive functional groups that include acetal, lactam, acetonide and *tert*-butyldimethylsilyl ether.

The results are collated in the Table 1. The carbonyl substrates used in this study contained a number of other functional groups such as chloro, methoxy, cyano, nitro, and urethane. In the case of the *N*-methylamines derived from aldehydes, the pure products were isolated by simple diethyl ether extraction. For the products derived from ketonic substrates,



Scheme 1.

Synthesis of *N*-Methyl Secondary Amines

1413

Table 1. Synthesis of *N*-methylamine.

Entry	Carbonyl compound	Product amine ^a	% Yield ^b
1			85
2			88
3			70
4			95
5			82
6			85
7			80
8			88
9			92
10			90
11			92

(continued)



1414

Kumpaty, Williamson, and Bhattacharyya

Table 1. Continued.

Entry	Carbonyl compound	Product amine ^a	% Yield ^b
12			93
13			90
14			90
15			88
16			92

^aThe products were characterized by comparison of their ¹H, ¹³C NMR and GC-mass spectral data with the reported^[5] data.

^bYields are of isolated and pure products.

the crude *N*-methyl amines were extracted with hydrochloric acid (2 M) to separate the neutral materials. Unlike many reported reductive amination^[4] protocols, the present method works well with enolizable carbonyl compounds. The neutral nonaqueous reaction conditions, simple isolation of the products with high yields, and the convenient use of a commercially available solution of methylamine are the notable features of the present method. This protocol should provide an easy access to other amines bearing functionalized motifs due to the compatibility of titanium(IV) isopropoxide with a variety of acid-sensitive functional groups.

In summary, an expedient, high throughput access to various *N*-methyl secondary amines is reported via reductive amination of carbonyl compounds with a commercially available solution of methylamine in methanol using titanium(IV) isopropoxide and sodium borohydride.



EXPERIMENTAL SECTION

General Procedure for the Reductive Amination of Aldehydes with Methylamine

Titanium(IV) isopropoxide (2 mL, 6.6 mmol) was added to a commercially available solution of methylamine in methanol (2 M, 7.5 mL) followed by the addition of the starting aldehyde (5 mmol). The reaction mixture was stirred at ambient temperature for 5 h, after which sodium borohydride (0.2 g, 5 mmol) was added and the resulting mixture was further stirred for another period of 2 h. The reaction was then quenched by the addition of water (1 mL), the resulting inorganic precipitate was filtered and washed with diethyl ether (20 mL). The organic layer was separated and the aqueous part was further extracted with diethyl ether (20 mL \times 2). The combined ether extracts were dried (K_2CO_3) and concentrated in vacuo to give *N*-methyl secondary amines in high purity.

General Procedure for the Reductive Amination of Ketones with Methylamine

For the reductive amination of ketones, the same general procedure was used except that the combined diethyl ether extracts were next extracted with hydrochloric acid (2 M, 10 mL \times 2) to separate the neutral materials. The acidic aqueous solution containing the *N*-methylated amine hydrochloride salt was made alkaline (pH = 10) by slow addition of (10%, w/v) aqueous NaOH and extracted with diethyl ether (20 mL \times 2). The combined organic extracts were dried (K_2CO_3) and concentrated in vacuo to give pure *N*-methylated alkylamines.

REFERENCES

1. For some leading references, see: Statistical investigation into the structural complementarity of natural products and synthetic compounds: Henkel, T.; Brunne, R.M.; Mueller, H.; Reichel, F. *Angew. Chem. Int. Ed.* **1999**, 38, 643; Main, B.G.; Tucker, H. *Medicinal Chemistry*, 2nd Ed.; Genellin, C.R., Roberts, S.M., Ed.; Academic Press: New York, 1993; p. 187; Kukhar, V.P.; Svistunova, N. Yu.; Soloshonok, V.A.; Solodenko, V.A. *Russ. Chem. Rev. (Engl. Transl.)* **1993**, 62, 284; Kirschbaum, J. *Analytical Profiles of*



- Drug Substances*; Florey, K., Ed.; Academic Press: New York, 1983; Vol. 12, p. 1.
2. Brown, A.R.; Rees, D.C.; Rankovic, Z.; Morphy, J.R. *J. Am. Chem. Soc.* **1997**, *119*, 3288; see Ref. 10 cited therein.
 3. For example, see: Bhattacharyya, S. *Tetrahedron Lett.* **1994**, *35*, 2401; Bhattacharyya, S. *J. Org. Chem.* **1995**, *60*, 4928; Bhattacharyya, S.; Neidigh, K.A.; Avery, M.A.; Williamson, J.S. *Synlett* **1999**, 1781.
 4. For some recent reviews on reductive aminations see: (a) Whitesell, J.K. *Comprehensive Organic Synthesis*; Trost, B.M., Fleming, I., Eds.; Pergamon Press: Oxford, 1991; Vol. 6, p.724; (b) Hutchins, R.O.; Hutchins, M.K. *Comprehensive Organic Synthesis*; Trost, B.M., Fleming, I., Eds.; Pergamon Press: Oxford, 1991; Vol. 8, p. 25; (c) Gribble, G.W. *Reductions in Organic Synthesis*; Abdel-Majid, A.F., Ed.; ACS Symposium Series 641, American Chemical Society: Washington, DC, 1996; p. 167; (d) Abdel-Majid, A.F. *Reductions in Organic Synthesis*; ACS Symposium Series 641, American Chemical Society: Washington, DC, 1996; p. 201.
 5. Neidigh, K.A.; Avery, M.A.; Williamson, J.S.; Bhattacharyya, S.J. *Chem. Soc. Perkin Trans. 1* **1998**, 2527.
 6. Seebach, D.; Hungerbuhler, E.; Naef, R.; Schnurrenberger, P.; Weidmann, B.; Zueger, M. *Synthesis* **1982**, 138.

Received in the Netherlands June 15, 2002