

Phytochemistry, Vol. 49, No. 5, pp. 1375–1376, 1998 © 1998 Elsevier Science Ltd. All rights reserved Printed in Great Britain 0031–9422/98/5—see front matter

PII: S0031-9422(98)00091-0

AN ORSELLINIC ACID GLUCOSIDE FROM SYZYGIUM AROMATICA

REENA CHARLES, SHRI NIWAS GARG* and SUSHIL KUMAR

Department of Phytochemical Technology, Central Institute of Medicinal and Aromatic Plants, Lucknow–226015, India

(Received in revised form 2 February 1998)

Key Word Index—Syzygium aromatica; Myrtaceae; orsellinic acid glucoside.

Abstract—The buds of *Syzygium aromatica* yielded two triterpenoid acids, oleanolic acid and crategolic acid, and a new phenolic glucoside, orsellinic-2-O- β -D-glucopyranoside, from its methanolic extract. The structure was elucidated by spectroscopic methods and acid hydrolysis. © 1998 Elsevier Science Ltd. All rights reserved

INTRODUCTION

Syzygium aromatica (L.) Merr and Perry, a tree commonly known as clove, is indigenous to the Moluccas. In India it is cultivated in Kerala, Tamil Nadu, Karnataka, and Andaman and Nicobar Islands. Clove buds (whole or ground) are mainly used as a spice for domestic culinary purposes and for the industrial manufacture of sauces and pickles [1]. The essential oil obtained from distillation of the buds is of the highest quality and finds extensive use in the perfumery and flavouring industries and to a limited extent in the pharmaceutical industry. This oil shows antibacterial activity [2], while the acetone extract of buds shows anticholagouge activity [3]. Eugeniin, a compound isolated from the ethyl acetate extract of clove buds shows antiviral activity [4]. Most studies [5-7] have been concerned with the chemical constituents of the essential oil.

Despite its great importance in Indian perfumery, flavouring and pharmaceutical industries, *S. aromatica* had been the subject of limited investigation. This study has revealed the presence in the buds of oleanolic acid (which is already used as a drug in China [8] for the treatment of jaundice), crategolic acid and a new phenolic glucoside, orsellinic-2-O- β -D-glucopyranoside.

RESULTS AND DISCUSSION

The methanolic extract of *S. aromatica* buds yielded two triterpenoid acids, oleanolic acid and crategolic

acid [9], and a phenolic glucoside (1) orsellinic-2-O- β -D-glucopyranoside. Chemical analysis and the mass spectrum of compound 1 showed it to have the molecular formula $C_{14}H_{18}O_9$. IR absorption bands at 3490 and 1668 cm⁻¹ were indicative of hydroxy and carboxyl groups, respectively. The presence of a carbonyl group was further confirmed by the presence of a signal at δ 184.34 in its ¹³C NMR spectrum. The ¹H NMR spectrum revealed the presence of two meta coupled aromatic protons at δ 5.93 (d, J = 2 Hz) and 6.09 (d, J = 2 Hz) and a singlet at δ 2.25 (3H) indicative of an aromatic methyl group. Signals at δ 162.80, 166.85 and 143.91 were attributed to C-2 and C-4 of the benzene ring, and a ring methyl group at C-6, respectively [10–14]. Further signals in the ¹H NMR and ¹³C NMR spectra clearly indicated the presence of a glucose moiety. A one proton doublet at δ 4.78 (J = 7 Hz) was assigned to the β -anomeric proton which was further supported by the chemical shift in the ¹³C NMR spectrum at δ 104.70. Hydrolysis of compound 1 with methanolic HCl afforded two products; a sugar which was identified as glucose by paper chromatography and an aglycone identical to orsellinic acid (mp., mmp and Co-TLC). Thus, from the above spectroscopic and chemical data, compound 1 was concluded to be orsellinic-2-O- β -D-glucopyranoside.

EXPERIMENTAL

Mp: uncorr.; NMR: 200 MHz (¹H) and 50 MHz (¹³C) CD₃OD with TMS as int standard; MS: 70 eV (direct inlet); TLC: Merck silica gel 60 F_{254} plates, spots visualised by spraying with a 1% sol of vanillin in H₂SO₄ followed by heating at 110° for 15 min; CC: silica gel (60–120 mesh).

^{*}Author to whom correspondence should be addressed. Tel: 00 91 342676; Fax: 00 91 522342666; E-mail: root-@cimap.sirnetd.ernet.in. CIMAP Communication No. 97– 99 J





Plant material

S. aromatica was purchased from the local market and a specimen voucher has been deposited at the Institute herbarium.

Extraction and isolation

Powdered buds of *S. aromatica* (1.5 kg) were extracted with *n*-hexane, CHCl₃ and MeOH. The solvents were removed *in vacuo* to give three frs weighing 112.5 g, 100.5 g and 195 g, respectively. Fifty grams of the MeOH extract on CC on silica gel using hexane, hexane–EtOAc, EtOAc and EtOAc–MeOH, afforded oleanolic acid (1 g), crategolic acid (30 mg) and compound (1) (40 mg), $R_f 0.54$ in CHCl₃–MeOH (3 : 1).

Orsellinic-2-*O*-β-D-glucopyranoside (1). Colourless crystals (MeOH), mp 190–191°, $[\alpha]_{D}^{18} + 2.4°$ (*c* 1.0: MeOH). UV [λ]^{MeOH} nm : 255 and 295; IR [ν]^{MBT} cm⁻¹: 3490, 3279, 2921, 2912, 1668, 1615, 1542, 1457, 1421, 1386, 1320, 1268, 1220, 1028, 867, 840 and 735; ¹H NMR (200 MHz, CD₃OD) : δ 2.25 (*s*, 3H, CH₃), 3.30 to 3.38 (*m*, 2H), 3.58 (*dd*, *J* = 12,2 Hz, 1H), 3.72 (*dd*, *J* = 12,5 Hz, 1H), 3.97 (*dd*, *J* = 12,2 Hz, 1H), 4.78 (*d*, *J* = 7 Hz, 1H), 5.93 (*d*, *J* = 2 Hz, 1H), 6.09 (*d*, *J* = 2 Hz, 1H); ¹³C NMR (CD₃OD) : δ 20.98 (CH₃), 62.91 (C-6'), 72.86 (C-4'), 75.49 (C-2'), 78.90 (C-5'), 80.02 (C-3'), 100.72 (C-3), 104.70 (C-1'), 105.32 (C-1), 108.75 (C-5), 143.97 (C-6), 162.80 (C-4), 166.85 (C-2), 184.34 (C-8); EI-MS *m/z* (rel. int.): 330 [M]⁺ (5), 312

 $[M-H_2O]^+$ (8), 168 (90) (Found : C, 51.1 ; H, 50.38. C₁₄H₁₈O₉ requires C, 50.49 : H, 5.45%).

Acid hydrolysis of 1. Compound 1 (28 mg) in MeOH (2 ml) was refluxed with 5% HCl (2 ml) for 4 h in a water bath. After cooling, the reaction mixture was extracted with EtOAc. The EtOAc layer was dried over MgSO₄ and concd under reduced pressure. The extract was recrystallised from MeOH to give 2 (10 mg), mp 169–170°, which was identified as orsellinic acid by TLC, mp, and mmp. The aq. layer was processed for sugar. This sugar was identified as glucose by comparison with a standard sample on PC (BuOH–HOAc–H₂O, 4:1:5), using aniline hydrogen phthalate as the spray reagent.

Acknowledgements—We are grateful to the Head of the Phytochemical Technology Division for providing facilities and encouragement throughout the work. R.S.I.C. Lucknow is thanked for recording the 200 MHz ¹H NMR and 50 MHz ¹³C NMR and mass spectra. Dr G. N. Srivastava is gratefully thanked for help in procuring the genuine plant material.

REFERENCES

- 1. Anonymous, *Wealth of India*, Vol 10, CSIR, New Delhi, India, 1976, p. 93.
- Singh, B. S. and Singh, G., *Indian Drugs*, 1978, 15, 227.
- Husain, A., Virmoni, O. P., Popli, S. P., Misra, L. N., Gupta, M. M., Srivastava, G. N., Abraham, Z. and Singh, A. K., *Dictionary of Indian Medicinal Plants*, CIMAP, Lucknow, 1992.
- TaKechi, M. and Tnanaka, Y., *Plant Med.*, 1981, 42, 69–74.
- Lawrence, B. M., Major tropical spices-clove in essential oils 1976–77, Allured-Publ. Corp. Wheaton III, 1978.
- Masada, Y., Analysis of essential oil by gas chromatography and Mass Spectrometry, John Wiley and Sons Inc., New York, 1976.
- Muchalol, M. and Crouzet, J., *Minor volatile components of clove essential oil*, IX International Congress of essential oils at Singapor, 1983.
- 8. Liu, J., J. Ethnopharmacology, 1995, 49, 57-68.
- Narayanan, C. R. and Natu, A. A., *Phyto*chemistry, 1974, 13, 1999–2000.
- Donnelly, D. M. X. and Hutchinson, R. H., *Phy*tochemistry, 1990, **29**, 179–182.
- Suksamrarn, A., Eiamony, S., Piyachaturawat, P. and Byrni, L. T., *Phytochemistry*, 1997, 45, 103– 105.
- Elix, J. A., Jayanthi, V. K. and Wardlaw, J. H., Aust. J. of Chem., 1989, 42, 1423–1426.
- Elix, J. A. and Whitan, A. A., Aust. J. Chem., 1989, 42, 1961–1981.
- Xu, H. X., Kadota, S., Kurokawa, M., Shiraki, K., Matsunoto, T. and Namba, T., *Chem. Pharm. Bull.*, 1993, **41**, 1803–1806.
- Ghosal, S., Mittal, P., Kumar, Y. and Singh, S. K., *Phytochemistry*, 1989, 28, 3193–3196.