

Accepted Article

Title: Reconstitution of Iterative Thioamidation in Closthioamide Biosynthesis Reveals a Novel Nonribosomal Peptide Backbone-Tailoring Strategy

Authors: Kyle L. Dunbar, Maria Dell, Evelyn M. Molloy, Florian Kloss, and Christian Hertweck

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.201905992
Angew. Chem. 10.1002/ange.201905992

Link to VoR: <http://dx.doi.org/10.1002/anie.201905992>
<http://dx.doi.org/10.1002/ange.201905992>

Reconstitution of Iterative Thioamidation in Closthioamide Biosynthesis Reveals a Novel Nonribosomal Peptide Backbone-Tailoring Strategy

Kyle L. Dunbar,^[a] Maria Dell,^[a] Evelyn M. Molloy,^[a] Florian Kloss,^[b] and Christian Hertweck^{*[a,c]}

Abstract: Thioamide-containing nonribosomal peptides (NRPs) are exceedingly rare. Recently the biosynthetic gene cluster for the thioamidated NRP antibiotic closthioamide (CTA) was reported; however, the enzyme responsible for and the timing of thioamide formation remained enigmatic. Here, we use genome editing, biochemical assays and mutational studies to demonstrate that an Fe-S cluster-containing member of the adenine nucleotide α -hydrolase protein superfamily (CtaC) is responsible for sulfur incorporation during CTA biosynthesis. However, unlike all previously characterized members, CtaC functions in a thiotemplated manner. In addition to prompting a revision of the CTA biosynthetic pathway, the reconstitution of CtaC provides the first example of a NRP thioamide synthetase. Finally, using CtaC as a bioinformatic handle, we provide promising clues that thioamidated NRP biosynthetic gene clusters are more widespread than previously appreciated.

Thioamides are formed through the oxygen-by-sulfur substitution of amide bonds, a seemingly minor change that has major repercussions for the physicochemical properties of the compound.^[1] Accordingly, site-specific thioamidation has long been used by synthetic chemists to tune the properties of small molecules. In regard to natural products, only a modest number of thioamidated secondary metabolites are known.^[1c, 2] One notable example is closthioamide (CTA, Figure 1A), a symmetric nonribosomal peptide (NRP) produced by the anaerobic bacterium *Ruminiclostridium cellulolyticum* DSM 5812.^[3] This DNA gyrase inhibitor harbors six thioamide linkages that are essential for its antimicrobial activity and is the only bacterial thioamide-containing NRP identified to date.^[3a, 4]

Recently, we demonstrated that CTA is biosynthesized by an unusual thiotemplated, NRP synthetase (NRPS)-independent pathway (Figure 1B) and proposed a model for CTA maturation.^[5] However, the enzyme responsible for thioamide biosynthesis and the timing of thioamide formation remained unknown. Indeed, nothing is known about how these moieties are made in NRPs, in contrast to the great strides made towards understanding thioamide formation in both ribosomal peptides and nucleoside antimetabolites.^[1c, 6] Here, we identify and reconstitute the activity of a NRP thioamide synthetase (CtaC) involved in CTA maturation. Through phylogenetic analyses and biochemical assays, we demonstrate that CtaC is a novel

member of the adenine nucleotide α -hydrolase (AANH) protein superfamily that processes thiotemplated substrates. Through the use of substrate derivatives, we determine the timing of thioamidation and revise the CTA biosynthetic pathway. Finally, we leveraged the sequence of CtaC to demonstrate that diverse anaerobic bacteria have the potential to produce thioamidated NRPs.

Of the genes present in the CTA biosynthetic gene cluster, we predicted that the product of *ctaC* (Ccel_3258) would be responsible for thioamide biosynthesis as it is a member of the AANH protein superfamily (Figure 1C and Supplementary Information Figure S1).^[5] Members of this superfamily catalyze transformations ranging from cofactor and antibiotic biosynthesis to the posttranslational modification of proteins.^[7] Although these reactions are varied, all involve ATP-dependent adenylation of the substrate followed by nucleophilic attack at the activated position and elimination of AMP. Notably, some AANH proteins perform oxygen-by-sulfur substitutions on nucleotides in both primary and secondary metabolic pathways.^[6b, 7-8] However, no AANH protein is known to thioaminate a peptidic substrate.

While analysis of the metabolite profile of CtaC-deficient *R. cellulolyticum* indicated that CtaC is essential for CTA biosynthesis, no biosynthetic intermediates could be detected to clarify the role of this uncharacterized AANH protein.^[5] However, it is possible that the type II intron inserted in *ctaC* causes polar effects.^[9] We therefore used a CRISPR-Cas9 genome editing method to generate a new *R. cellulolyticum* Δ *ctaC* strain with a nonsense mutation in *ctaC* (Figure S2);^[5, 10] however, no oxygen isosteres of known CTA congeners were detected in culture extracts (Figure S3). We next turned to an *Escherichia coli* heterologous expression system for the production of polythioamidated peptides.^[5] A vector containing all of the CTA biosynthetic genes except *ctaC* was constructed (pCTA- Δ *ctaC*; Figure S4) and the metabolite profile of *E. coli* pCTA- Δ *ctaC* was checked for the accumulation of intermediates. Consistent with the *R. cellulolyticum* Δ *ctaC* results, no thioamide-free intermediates were detected (Figure S4).

In order to circumvent the difficulties in defining the role of CtaC by genetic approaches, we set out to characterize CtaC *in vitro*. Owing to solubility issues with His₆-tagged CtaC, the protein was isolated as an N-terminal maltose binding protein (MBP)-tagged fusion protein (Figure S5). During isolation we noted that aerobically purified MBP-CtaC solutions were slightly red, potentially indicating that CtaC binds iron. A recently identified tRNA-modifying subfamily (TtcA/TtuA/Ncs6) of the AANH superfamily was shown to require an iron-sulfur cluster for activity.^[11] Indeed, CtaC has the [4Fe-4S] cluster-binding motif found in TtcA/TtuA proteins (Figure S6).^[11a, 11c, 11d] However, the UV-visible spectrum of MBP-CtaC had only a weak absorbance at the characteristic wavelength for such metal centers (~410 nm,^[11a, 11c, 11d] Figure 2A) and purified with just 1.2 \pm 0.1 iron and 0.9 \pm 0.1 sulfide bound per protomer. To obtain MBP-CtaC with an appropriately assembled iron-sulfur cluster, the protein was treated with sulfide and ferrous iron in an anaerobic glovebox. The resultant MBP-CtaC solution adopted a brownish-red color, had an enhanced absorbance at 410 nm

[a] Dr. K. L. Dunbar, M. Dell, Dr. E. M. Molloy, Prof. Dr. C. Hertweck
Dept. of Biomolecular Chemistry, Leibniz Institute for Natural
Product Research and Infection Biology, HKI
Beutenbergstrasse 11a, 07745 Jena (Germany)
E-mail: christian.hertweck@leibniz-hki.de

[b] Dr. F. Kloss
Transfer Group Antiinfectives, Leibniz Institute for Natural Product
Research and Infection Biology, HKI
Beutenbergstrasse 11a, 07745 Jena (Germany)

[c] Prof. Dr. C. Hertweck
Chair of Natural Product Chemistry
Friedrich Schiller University Jena, 07743 (Germany)

Supporting information for this article is given via a link at the end of the document.




Figure 1. The CTA gene cluster contains an AANH homolog. (A) Structure of CTA. (B) The CTA biosynthetic gene cluster. AANH, adenine nucleotide α -hydrolase; PCP, peptidyl carrier protein. (C) Neighbor-joining phylogenetic tree of AANH proteins (1000 bootstrap replicates). Structures of the compounds biosynthesized by each AANH subclass are shown with the installed moiety colored red. See Supporting Information for a fully annotated cladogram.

(Figure 2A), and contained 4.5 ± 0.2 iron and 4.5 ± 0.1 sulfide per protomer.

Based on the thiotemplated nature of CTA biosynthesis, we previously hypothesized that thioamide formation would occur on substrates linked to one of the two peptidyl carrier proteins (PCPs; CtaE/CtaH) encoded in the biosynthetic gene cluster (Figure 1B).^[5] Thus, we purified CtaE and CtaH as N-terminal His₆-tagged fusions and used matrix-assisted laser desorption/ionization-time-of-flight-mass spectrometry (MALDI-TOF-MS) to verify that they were in their *apo* form (*i.e.* lacking the phosphopantetheinyl arm, Figure S7). As the timing of thioamide formation during CTA maturation was unknown, we synthesized coenzyme A (CoA)-conjugates differing in the presence or absence of 4-hydroxybenzoic acid (HBA) (1-CoA and 2-CoA; Figure 2B). These synthetic CoA-linked substrates were loaded on the *apo*-PCPs by the promiscuous phosphopantetheinyl transferase (PPTase) Sfp to generate the corresponding peptidyl-*holo*-PCPs (*e.g.* 1-*holo*-CtaE; Figure S7).^[12]

With appropriately loaded PCPs prepared, we tested CtaC for thioamidation activity. As all AANH superfamily members require ATP for activity,^[7] thioamidation reactions were supplemented with ATP and magnesium. Lithium sulfide acted as the sulfur donor since a sulfur-mobilization enzyme is not encoded in the CTA biosynthetic gene cluster and other sulfur-inserting AANH enzymes have been shown to accept this non-native sulfur donor *in vitro*.^[11b-d, 13] All thioamidation reactions

were performed in an anaerobic glovebox and progress was monitored by MALDI-TOF-MS. Although no processing of 1-*holo*-CtaH (Figure S8) or 2-*holo*-CtaE was observed (Figure S9), the mass of 1-*holo*-CtaE increased by approximately 32 Da in reactions compared to controls (Figure 2C). This shift is consistent with the installation of two thioamides and was dependent on the presence of CtaC, ATP and sulfide. To definitively identify the product of the CtaC reaction, the thioester linking 1 to *holo*-CtaE was hydrolyzed, and the released products were analyzed by LC-HR-MS. A peak with a retention time and mass consistent with di-thioamidated 1 was only observed in reactions containing all components (Figure 2D, Figure S10). Furthermore, MS/MS analysis of the product localized the sulfur atoms to the amide bonds (Figure S10).

In order to probe the substrate scope of CtaC, thioamidation reactions were performed on substrates not linked to CtaE (1, 1-CoA, 3-CoA, 4, and 5; Figure 2B). Despite using elevated enzyme concentrations and long reaction times, no turnover was observed by LC-HR-MS (Figure S11–S15). Together, these data demonstrate that CtaC is a thiotemplated thioamide synthetase that functions on CtaE-bound substrates and that sulfur insertion occurs after HBA is added to the poly- β -alanine backbone.

AANH enzymes adenylate their substrates and thus produce AMP as a byproduct of the reaction.^[7] When we measured the ATP hydrolysis products from thioamidation reactions with 1-*holo*-CtaE by HPLC, both AMP and ADP were detected (Figure S16). Moreover, neither the production of AMP nor ADP correlated with substrate processing by CtaC (Figure S16). Although members of the AANH superfamily are highly divergent in both primary sequence and function, they share a fully conserved ATP-binding motif that is used to present ATP for substrate adenylation.^[7] Thus, as an alternative approach to investigate the ATP-dependent activity of CtaC, we generated mutant proteins bearing single alanine substitutions in the ATP-binding pocket (CtaC_{D77A} and CtaC_{S78A}, Figure S5–S6) and assayed the mutants for their ability to thioaminate 1-*holo*-CtaE. As predicted, the activity of both mutants was abrogated (Figure 2E, Figure S17).

As mentioned above, members of the TtcA/TtuA family of sulfur-inserting AANH enzymes require an oxygen-labile [4Fe-4S] cluster coordinated by three conserved cysteine residues for activity.^[11a, 11c, 11d] Although the precise role is unclear, the prevailing hypothesis is that the cluster coordinates sulfide at the unique iron site and activates it for insertion into the substrate.^[11] To investigate the role of the Fe-S cluster in thioamide formation by CtaC, we generated mutant proteins bearing single alanine substitutions to residues predicted to be involved in Fe-S cluster-binding (CtaC_{C154A}, CtaC_{C157A}, and CtaC_{C284A}, Figure S5–S6). While all mutant proteins bound similar levels of iron and sulfide as wild-type CtaC (Figure S18), their thioamidation activity was diminished (Figure 2E, Figure S17). These results establish that although single mutations to the Fe-S cluster binding site are not sufficient to fully disrupt cluster coordination, the activity of CtaC is highly sensitive to perturbations to its metal binding site. Notably, this phenomenon has been observed for other Fe-S cluster dependent AANH enzymes.^[11a, 11d] In further support of the essential nature of the metal center for CtaC activity, processing of 1-*holo*-CtaE was completely lost in thioamidation reactions performed with CtaC under aerobic conditions or with CtaC stripped of the Fe-S cluster (Figure 2E, Figure S17).

Over the course of our studies we noted that regardless of the reaction time and concentration of CtaC used, tri-




Figure 2. CtaC is a PCP-dependent thioamide synthetase. (A) UV-visible absorbance spectra of as-isolated and reconstituted MBP-CtaC. (B) Structures of substrates used in this study. (C) Representative MALDI-TOF-MS spectral overlay of 6 h *in vitro* thioamidation reactions with 1-*holo*-CtaE. (D) LC-HR-MS profiles of thioamidation reactions with 1-*holo*-CtaE following thioester cleavage. Traces correspond to the extracted ion chromatogram of the $[M-H]^-$ ionic species for di-thioamidated **1** and are displayed with m/z values \pm 5 ppm from the calculated exact mass. (E) Representative MALDI-TOF-MS spectral overlay of 22 h *in vitro* thioamidation reactions with 1-*holo*-CtaE and mutant CtaC enzymes, CtaC under aerobic conditions (+ O₂), or *apo*-CtaC (- Fe-S). (F) Representative MALDI-TOF-MS spectral overlay of 22 h thioamidation reactions with 3-*holo*-CtaE. (G) Updated biosynthetic scheme for CTA maturation. (C, E, F) Guide lines denote the expected shifts (+ 16 Da) obtained from the substitution of oxygen by sulfur. Red line: heat-inactivated CtaC.

thioamidated 1-*holo*-CtaE was never detected (Figure S19), accounting for only four of the six thioamides present in CTA. The initial biosynthetic model for CTA maturation posited that CtaC installs the final two thioamides after the dimerization of di-thioamidated **1** through diaminopropane (DAP);^[5] however, the inability of CtaC to process free substrates casts doubt on this proposal. Indeed, products bearing three or four thioamidated β -alanine residues were detected from *R. cellulosyticum* and the *E. coli* heterologous expression system, although their biosynthetic relevance was unknown.^[3b, 5] We therefore reasoned that a CtaE-bound substrate containing three, rather than two, β -alanine residues is the on-pathway substrate for thioamidation by CtaC. Accordingly, 3-CoA (Figure 2B) was synthesized, loaded onto CtaE *in vitro* using Sfp, and the conjugate (3-*holo*-CtaE) was tested for processing by CtaC. The MALDI-TOF-MS spectra demonstrated that the mass of 3-*holo*-CtaE increased by approximately 48 Da in reactions compared to controls (Figure

2F). Furthermore, using a combination of LC-HR-MS and LC-HR-MS/MS, the product was assigned as tri-thioamidated **3** (Figure S20).

In light of these data, we propose a revised biosynthetic pathway for CTA maturation (Figure 2G). Briefly, the ATP-grasp (CtaD) and acetyl-CoA synthetase (CtaI) proteins are predicted to load β -alanine and HBA onto CtaE and CtaH, respectively. HBA is synthesized from chorismate by CtaA^[5] and β -alanine is either biosynthesized from aspartate by the decarboxylase CtaF or taken from primary metabolism. The peptidyl transferase CtaG would transfer HBA from CtaH to tri- β -alanine-CtaE to form 3-CtaE. Following the thioamidation of 3-CtaE by CtaC, DAP — which is potentially synthesized from aspartate by the sequential action of a reductase (CtaK), aminotransferase (CtaB), and decarboxylase (CtaF) — would be coupled to two molecules of tri-thioamidated **3** by the transglutaminase CtaJ to form CTA.^[5] We suggest that the terminal β -alanine remains on




Figure 3. Homologs of CtaC are found in diverse thiotemplated biosynthetic pathways. (A) Sequence similarity network of CtaC homologs. Nodes are colored based on the class of biosynthetic machinery found near the CtaC homolog, while their shape denotes the phylum of the organism harboring the gene cluster. NCT, non-canonical thio-template; PKS, polyketide synthase. (B) Representative biosynthetic gene clusters that encode a CtaC homolog (selected based on architecture diversity). Numbers correspond to the node numbering in panel A. rSAM, radical S-adenosylmethionine enzyme.

CtaE, thus serving as a priming residue that allows CtaC to concurrently install all thioamides in chemically equivalent positions (Figure 2G).

As CtaC is the first example of a NRP thioamide synthetase, we sought to determine if this strategy for thioamidated NRP biosynthesis exists outside CTA maturation. To this end, we retrieved 100 homologs of CtaC from GenBank and generated a sequence similarity network with the protein sequences (Figure 3A).^[14] Next, we surveyed the genomic neighborhood surrounding the *ctaC* homolog for genes encoding secondary metabolite biosynthetic enzymes (Figure 3A). Notably, genes encoding thiotemplated biosynthetic enzymes were found in approximately 75% of cases. In addition to a small number of CTA-like biosynthetic gene clusters, many CtaC homologs are encoded in the vicinity of NRPSs (Figure 3B), suggesting that this strategy for thioamidation is also used by canonical NRP biosynthetic pathways. Moreover, the majority of homologs were

found in anaerobic bacteria (Table S5), suggesting that anaerobes represent an untapped trove of thioamidated NRPs.

In summary, we have reconstituted the first instance of thioamide formation in a NRP. The data presented herein demonstrate that this unusual thioamide synthetase functions on thiotemplated substrates and represents a new subfamily of the Fe-S cluster dependent AANH enzymes. In addition to further clarifying the CTA biosynthetic pathway, the characterization of CtaC serves as a foundation for the discovery of additional thioamidated NRPs. Future studies will focus on unearthing the native sulfur source for this enzyme and the characterization of the remaining steps in CTA maturation.

Acknowledgements

We thank A. Perner, T. Kindel and M. García-Altare Pérez for MS measurements and H. Heinecke for NMR measurements. We are grateful to K. Ishida and J. Kumpfmüller for technical advice. K.L.D. and F.K. were supported by the Humboldt Research Fellowship for Postdoctoral Researchers and InfectControl 2020 (FKZ 03ZZ0803A), respectively. Financial support by the DFG (Leibniz Award to C.H.) is gratefully acknowledged.

Conflict of Interest

The authors declare no conflict of interest.


Keywords: antibiotics • biosynthesis • enzymes • natural products • thioamide

- [1] a) A. Choudhary, R. T. Raines, *ChemBioChem* **2011**, *12*, 1801-1807; b) E. J. Petersson, J. M. Goldberg, R. F. Wissner, *Phys. Chem. Chem. Phys.* **2014**, *16*, 6827-6837; c) N. Mahanta, D. M. Szantai-Kis, E. J. Petersson, D. A. Mitchell, *ACS Chem. Biol.* **2019**, *14*, 142-163.
- [2] a) K. L. Dunbar, D. H. Scharf, A. Litomska, C. Hertweck, *Chem. Rev.* **2017**, *117*, 5521-5577; b) S. Banala, R. D. Sussmuth, *ChemBioChem* **2010**, *11*, 1335-1337.
- [3] a) T. Lincke, S. Behnken, K. Ishida, M. Roth, C. Hertweck, *Angew. Chem. Int. Ed.* **2010**, *49*, 2011-2013; b) S. Behnken, T. Lincke, F. Kloss, K. Ishida, C. Hertweck, *Angew. Chem. Int. Ed.* **2012**, *51*, 2425-2428.
- [4] a) A. I. Chiriach, F. Kloss, J. Kramer, C. Vuong, C. Hertweck, H. G. Sahl, *J. Antimicrob. Chemother.* **2015**, *70*, 2576-2588; b) V. F. Miarí, P. Solanki, Y. Hleba, R. A. Stabler, J. T. Heap, *Antimicrob. Agents Chemother.* **2017**, *61*; c) F. Kloss, A. I. Chiriach, C. Hertweck, *Chem. Eur. J.* **2014**, *20*, 15451-15458.
- [5] K. L. Dunbar, H. Buttner, E. M. Molloy, M. Dell, J. Kumpfmüller, C. Hertweck, *Angew. Chem. Int. Ed.* **2018**, *57*, 14080-14084.
- [6] a) N. Mahanta, A. Liu, S. Dong, S. K. Nair, D. A. Mitchell, *Proc. Natl. Acad. Sci. U.S.A.* **2018**, *115*, 3030-3035; b) A. Litomska, K. Ishida, K. L. Dunbar, M. Boettger, S. Coyne, C. Hertweck, *Angew. Chem. Int. Ed.* **2018**, *57*, 11574-11578; c) G. E. Kenney, L. M. K. Dassama, M. E. Pandelia, A. S. Gizzi, R. J. Martinie, P. Gao, C. J. DeHart, L. F. Schachner, O. S. Skinner, S. Y. Ro, X. Zhu, M. Sadek, P. M. Thomas, S. C. Almo, J. M. Bollinger, Jr., C. Krebs, N. L. Kelleher, A. C. Rosenzweig, *Science* **2018**, *359*, 1411-1416; d) D. D. Nayak, N. Mahanta, D. A. Mitchell, W. W. Metcalf, *Elife* **2017**, *6*; e) J. Liu, Z. Lin, Y. Li, Q. Zheng, D. Chen, W. Liu, *Org. Biomol. Chem.* **2019**, *17*, 3727-3731; f) C. J. Schwalen, G. A. Hudson, B. Kille, D. A. Mitchell, *J. Am. Chem. Soc.* **2018**, *140*, 9494-9501.
- [7] M. Fellner, R. P. Hausinger, J. Hu, *Crit. Rev. Biochem. Mol. Biol.* **2018**, *53*, 607-622.
- [8] a) N. Shigi, *Front. Microbiol.* **2018**, *9*, 2679; b) S. Coyne, C. Chizzali, M. N. Khalil, A. Litomska, K. Richter, L. Beerhues, C. Hertweck, *Angew. Chem. Int. Ed.* **2013**, *52*, 10564-10568.
- [9] K. N. McAllister, L. Bouillaut, J. N. Kahn, W. T. Self, J. A. Sorg, *Sci. Rep.* **2017**, *7*, 14672.
- [10] T. Xu, Y. Li, Z. Shi, C. L. Hemme, Y. Li, Y. Zhu, J. D. Van Nostrand, Z. He, J. Zhou, *Appl. Environ. Microbiol.* **2015**, *81*, 4423-4431.
- [11] a) D. Bouvier, N. Labessan, M. Clemancey, J. M. Latour, J. L. Ravanat, M. Fontecave, M. Atta, *Nucleic Acids Res.* **2014**, *42*, 7960-7970; b) Y. Liu, D. J. Vinyard, M. E. Reesbeck, T. Suzuki, K. Manakongtreecheep, P.

- L. Holland, G. W. Brudvig, D. Soll, *Proc. Natl. Acad. Sci. U.S.A.* **2016**, *113*, 12703-12708; c) S. Arragain, O. Bimai, P. Legrand, S. Caillat, J. L. Ravanat, N. Touati, L. Binet, M. Atta, M. Fontecave, B. Golinelli-Pimpaneau, *Proc. Natl. Acad. Sci. U.S.A.* **2017**, *114*, 7355-7360; d) M. Chen, S. I. Asai, S. Narai, S. Nambu, N. Omura, Y. Sakaguchi, T. Suzuki, M. Ikeda-Saito, K. Watanabe, M. Yao, N. Shigi, Y. Tanaka, *Proc. Natl. Acad. Sci. U.S.A.* **2017**, *114*, 4954-4959.
- [12] L. E. Quadri, P. H. Weinreb, M. Lei, M. M. Nakano, P. Zuber, C. T. Walsh, *Biochemistry* **1998**, *37*, 1585-1595.
- [13] C. T. Lauhon, W. M. Erwin, G. N. Ton, *J. Biol. Chem.* **2004**, *279*, 23022-23029.
- [14] J. A. Gerlt, J. T. Bouvier, D. B. Davidson, H. J. Imker, B. Sadkhin, D. R. Slater, K. L. Whalen, *Biochim. Biophys. Acta* **2015**, *1854*, 1019-1037.

Entry for Table of Contents:

COMMUNICATION



K.L. Dunbar, M. Dell, E. M. Molloy, F. Kloss, C. Hertweck*

Page No. – Page No.

Reconstitution of Iterative Thioamidation in Closthioamide Biosynthesis Reveals a Novel Nonribosomal Peptide Backbone-Tailoring Strategy

Thiotemplated thioamidation: Closthioamide (CTA) is a thioamide-containing nonribosomal peptide (NRP). The thioamide synthetase responsible for sulfur insertion in CTA assembly was identified and characterized *in vitro*, shedding new light on the thiotemplated assembly of this unusual antibiotic. Bioinformatic analyses indicate that anaerobic bacteria may represent an untapped source for thioamidated NRPs.