Natural Products

Total Synthesis of the Antibiotic Erypoegin H and Cognates by a PtCl₂-Catalyzed Cycloisomerization Reaction**

Alois Fürstner,* Eike K. Heilmann, and Paul W. Davies

The growing number of methicillin-resistant *Staphylococcus aureus* (MRSA) strains that are also resistant to other antibiotics in clinical use, including vancomycin, constitutes a serious threat worldwide. Even though the urgent need for innovative therapeutic agents capable of combating such deadly bacteria has been widely recognized, the number of new lead structures fueling that quest remains alarmingly low.^[1]

A promising candidate in this regard is the pterocarpene nucleus (**G**, Scheme 1) which constitutes the core of a sizeable number of bioactive natural products, mainly derived from plants of the *Leguminosae* family. Amongst the recent publications highlighting the potential of this heterocyclic scaffold as a pharmacophore,^[2,3] a report on the secondary metabolites of *Erythrina poeppigiana* is particularly noteworthy.^[4] It showed that erypoegin H (1), the most active



[*] Prof. A. Fürstner, Dipl.-Chem. E. K. Heilmann Max-Planck-Institut für Kohlenforschung 45470 Mülheim an der Ruhr (Germany) Fax: (+ 49) 208-306-2994 E-mail: fuerstner@mpi-muelheim.mpg.de
Dr. P. W. Davies School of Chemistry University of Birmingham

Edgbaston, Birmingham, B152TT (UK)

[**] Generous financial support by the MPG and the Fonds der Chemischen Industrie is gratefully acknowledged. We thank Dr. R. Mynott and his team for expert NMR assistance and Umicore AG & Co KG, Hanau, for a gift of noble metal salts. isoflavonoid isolated from the roots of this ornamental plant, not only exhibits a broad spectrum of activity against Grampositive bacteria in general, but also exhibits a significant and uniform activity against a panel of 13 different MRSA strains and vancomycin-resistant enterococci (minimum inhibitory concentration (MIC) 12.5 μ gmL⁻¹).^[4]

Preliminary data suggest that **1** interferes with the nucleic acids and/or the metabolism of MRSA bacteria, whilst seemingly leaving their cell walls intact.^[4] To fully assess the potential of this lead compound, more detailed studies on its mode of action and toxicity are warranted, and exploration of pertinent structure/activity relationships (SAR) should allow the activity to be further improved. To this end, it was necessary to develop a concise and inherently flexible route that delivers meaningful amounts of **1** and congeners (for example, **2–6**).

We reasoned that the benzofuran synthesis recently developed by our group would lend itself to this goal (Scheme 1).^[5-7] Based on a PtCl₂-catalyzed carboalkoxylation



 $\it Scheme 1.$ Synthesis of the pterocarpene nucleus by a $PtCl_2-catalyzed$ carboalkoxylation reaction of an alkyne.

reaction, the underlying concept diverges broadly from the established construction modes of pterocarpenes.^[8] Activation of the alkyne moiety in a substrate of type **A** by the carbophilic π acid PtCl₂ engenders nucleophilic attack from an oxygen atom of the adjacent acetal, thereby resulting in a *trans*-alkoxyplatination ($\mathbf{B} \rightarrow \mathbf{C}$).^[5,9] The released oxocarbe-



© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

nium ion **E** then shifts to the most nucleophilic position in **D** to give product $\mathbf{F}_{,}^{[10]}$ which can be readily elaborated into the targeted scaffold **G** by conventional means.

The synthetic venture commenced with the di-iodination of resorcinol $7^{[11]}$ followed by consecutive attachment of a pivaloyl and a trimethylsilylethoxymethyl group (Scheme 2).



Scheme 2. a) KI, KIO₃, aq HCl, 61%; b) PivCl, Et₃N, CH₂Cl₂, 64%; c) SEMCl, Et₃N, DMAP (cat.), toluene, 94%; d) PivCl, Et₃N, CH₂Cl₂, 99%; e) Me₃SiC=CH, [PdCl₂(PPh₃)₂] (1.5 mol%), Cul (1.5 mol%), Et₃N, Ar/H₂ (1 atm), 93%; f) AgNO₃ (10 mol%), aq acetone, 99%; g) [PdCl₂(PPh₃)₂] (3 mol%), Cul (3 mol%), Et₃N, Ar/H₂ (1 atm), 74%; h) PtCl₂ (10 mol%), CO (1 atm), 4.Å MS, toluene, 80°C, 84%; i) (NH₄)₂Ce(NO₃)₆ (CAN), MeCN/H₂O, 82%; j) 1. Zn(*i*Pr)₂, [Li(acac)] (10 mol%), NMP; 2. CuCN·2 LiCl (10 mol%), prenyl bromide, 86%; k) LiAlH₄; THF; l) I₂, PPh₃, imidazole, Et₂O/MeCN, 70% (over both steps); Piv = pivaloyl, DMAP = 4-dimethylaminopyridine, SEM = trimethylsilylethoxymethyl, MS = molecular sieves, acac = acetylacetone, NMP = N-methylpyrrolidine.

This sequence takes advantage of the greatly different accessibility of the two phenolic OH groups in **8**. Steric reasons also explain the regioselective course of the subsequent Sonogashira reaction of di-iodide **9** with alkyne **12**, which derives from compound $10^{[12]}$ in a few high yielding reactions; this transformation had to be carried out under a reducing atmosphere to avoid extensive homocoupling of the alkyne partner.^[13] The resulting compound **13**, on exposure to catalytic amounts of PtCl₂ in toluene under a CO atmosphere,^[5,14] underwent a remarkably clean cycloisomerization with formation of the desired benzofuran derivative **14**. This reaction was best performed in the presence of powdered

molecular sieves to sequester traces of adventitious water that might protonate the putative organoplatinum intermediate of type **C** and/or **D** and hence reduce the efficiency of the $O \rightarrow C$ shift. Under these optimized conditions, the cycloisomerization of 13 proceeded exceedingly well and afforded 14 in 84 % yield on a multigram scale (see the Experimental Section). The efficiency and operational simplicity of this particular transformation highlights once again the remarkable application profile of noble-metal-catalyzed skeletal rearrangements in general^[9,15] and toward heterocycle syntheses in particular.^[9,16] The pronounced migratory aptitude of the SEM acetal in 13 stands in striking contrast to the behavior of the symmetry-related pivaloyl group occupying an ortho position on the substrate's other benzene ring, which does not engage at all in such a $O \rightarrow C$ transfer and hence merely serves as a nonparticipating protecting group. Moreover, the compatibility of the iodide substituent is particularly noteworthy, and highlights the orthogonal nature of platinum catalysis to established redox-based transition-metal catalysts.^[9] In synthetic terms, the intact iodide was not only instrumental for the successful completion of the synthesis of erypoegin H but also constitutes a valuable entry point into late-stage divergent modifications of the target, as required for future SAR studies.

Next, the benzylic OH group had to be unmasked. This step turned out to be more difficult than anticipated, with a host of fluoride and Lewis acid based reagents failing to induce effective cleavage of the trimethylsilylethyl ether in **14**. Gratifyingly, oxidative conditions (CAN, aq MeCN)^[17,18] were found to deliver the corresponding aldehyde **15** in excellent yield and purity.

With this compound in hand, the stage was set for the introduction of the prenyl side chain^[18] which was originally envisaged to be carried out by conventional cross-coupling methodology. However, despite considerable experimentation, the yields remained invariably low and the reactions were plagued by the formation of various side-products. Recourse to the advanced metalation technology developed by Knochel and co-workers provided a most effective solution to this problem.^[19] Thus, in the presence of catalytic amounts of [Li(acac)] as a nucleophilic promoter of the exchange process, the reaction of 15 with $Zn(iPr)_2$ in NMP afforded a highly functionalized organozinc compound which could be cleanly alkylated with excess prenyl bromide in the presence of a catalytic amount of CuCN·2LiCl. It is remarkable that neither the aldehyde function nor the three ester moieties in 15 interfere with this reaction to any noticeable extent,^[19] as can be deduced from the 86% yield of 16, which could be isolated on a gram scale. We have no doubt that this methodology is amenable to further scale-up and will allow a host of other substituents to be introduced onto this advanced synthetic intermediate at a later stage of the project. Treatment of 16 with LiAlH₄ reduced the aldehyde and removed the peripheral ester groups concurrently. The resulting crude product 17 was subjected to an intramolecular etherification under standard conditions to complete the tetracyclic framework of erypoegin H (1), which was obtained in a respectable 28% yield over the nine steps of the longest linear sequence. The spectroscopic data of the synthetic

Communications

samples were in excellent agreement with those reported in the literature (Table 1).^[4]

Cleavage of the pivaloyl groups under nonreducing conditions converted aldehyde 16 into the des-methyl ana-

Table 1: Reference data set of compounds 1-3.

1: ¹H NMR (400 MHz, CDCl₃): δ = 7.36 (d, *I* = 8.1 Hz, 1 H), 7.04 (d, J=8.3 Hz, 1 H), 6.77 (d, J=8.3 Hz, 1 H), 6.45 (dd, J=8.1, 2.4 Hz, 1 H), 6.43 (d, J=2.3 Hz, 1 H), 5.54 (s, 2 H), 5.39 (thept, J=7.3, 1.3 Hz, 1 H), 5.26 (brs, 1 H), 5.01 (brs, 1 H), 3.69 (d, J = 7.3 Hz, 2 H), 1.89–1.90 (m, 3 H), 1.78 ppm (d, J=1.2 Hz, 3 H); ¹³C NMR (100 MHz, CDCl₃): $\delta =$ 156.7, 155.1, 154.5, 152.0, 147.0, 135.3, 121.2, 120.9, 119.2, 116.1, 112.6, 111.2, 110.2, 108.4, 106.2, 103.9, 65.7, 25.8, 23.1, 17.9 ppm; UV (MeOH/H₂O 4:1): λ_{max}=350, 334, 242, 210, 204 nm; MS (EI): *m/z* (%): 322 (77) [M]⁺, 266 (100), 237 (6), 152 (5); HRMS (EI): m/z: calcd for C₂₀H₁₈O₄: 322.1205; found: 322.1207 **2**: ¹H NMR (400 MHz, $[D_6]$ acetone): $\delta = 7.92$ (d, J = 8.6 Hz, 1 H), 7.62 (d, J = 8.4 Hz, 1 H), 7.05 (d, J = 8.3 Hz, 1 H), 7.02 (dd, J = 8.6, 2.3 Hz, 1 H), 6.95 (d, /=2.2 Hz, 1 H), 5.43 (thept, /=7.4, 1.4 Hz 1 H), 3.70 (d, J = 7.4 Hz, 2 H), 1.91–1.92 (m, 3 H), 1.69 ppm (d, J = 1.1 Hz, 3 H); ¹³C NMR (100 MHz, [D₆]acetone): δ = 161.9, 160.6, 158.5, 156.1, 156.0, 155.0, 132.6, 123.5, 122.5, 118.9, 116.6, 114.5, 114.4, 113.4, 106.1, 104.1, 103.9, 25.8, 23.4, 18.0 ppm; UV (MeOH): λ_{max} = 346, 304, 253, 209 nm; MS (EI): m/z (%): 337 (13), 336 (54) [M]⁺, 281 (44), 280 (100), 252 (7); HRMS (ESI): m/z: calcd for $C_{20}H_{16}O_5 + Na$: 359.0890; found: 359.0894 **3**: ¹H NMR (400 MHz, [D₆]acetone): δ = 7.90 (d, J = 8.6 Hz, 1 H), 7.66 (d, J = 8.5 Hz, 1 H), 7.01 (dd, J = 8.6, 2.2 Hz, 1 H), 6.95 (d, J = 2.2 Hz, 1 H),6.88 (d, J = 8.5 Hz, 1 H), 3.09 (t, J = 6.8 Hz, 2 H), 1.96 (t, J = 6.8 Hz, 2 H), 1.39 ppm (s, 6 H); ¹³C NMR (75 MHz, [D₆]acetone): $\delta = 162.0$, 160.6, 158.5, 156.1, 155.1, 154.2, 123.5, 119.4, 116.2, 116.2, 114.4, 107.3, 106.0, 104.1, 103.9, 75.6, 32.1, 26.8, 17.3 ppm. MS (EI): m/z (%): 337 (26), 336 (75) [M]⁺, 319 (4), 280 (100), 252 (7); HRMS (ESI): m/z: calcd for C₂₀H₁₆O₅ + Na: 359.0890; found: 359.0887.

logue of erypoegin F **4**, another flavonoid from *E. poeppigiana* endowed with considerable antimicrobial activity.^[4b] Furthermore, oxidation of **16**, deprotection of the ester moieties, and treatment of the resulting acid **19** with $H_2SO_4/HOAc$ led to concomitant lactonization and addition of the phenolic OH group onto the adjacent prenyl side chain; sojagol (**3**) thus formed is a well known phytoalexin originally isolated from *Glycine max* (soybeans) and *Phaseolus aureus* (mung beans; Scheme 3).^[20] Uncoupling of the two cyclization



Scheme 3. a) NaClO₂, NaH₂PO₄, tBuOH/H₂O, 2-methylbutene, 93%; b) KOMe, MeOH; c) H₂SO₄, HOAc, 52% (over two steps); d) DCC, DMAP, MeCN, 77% (over two steps). DCC = N,N'-dicyclohexylcarbodiimide.

events was also possible by treatment of **19** with DCC, which led to the proposed structure of isosojagol (**2**).^[21] However, the spectroscopic data of synthetic **2** clearly differ from those

of the putative "isosojagol" but closely match the reported spectra of "phaseol", an isomeric coumestan for which structure **5** had been proposed.^[22] Since the constitution of our synthetic sample is unambiguous, we must conclude that the original structure assignments of isosojagol and phaseol are *both* incorrect. While our study now shows that "phaseol" is definitely represented by structure **2** rather than by **5**, the actual constitution of "isosojagol" remains to be elucidated.

Experimental Section

14: A Schlenk flask was charged with compound 13 (4.25 g, 5.66 mmol), 4-Å MS (1.77 g, powdered), and toluene (30 mL). After addition of PtCl₂ (150 mg, 0.56 mmol), the mixture was purged with CO through a canula for 7 min before it was vigorously stirred at 85 °C under an atmosphere of CO (1 atm) for 5.5 h. For work up, the mixture was filtered through a pad of silica, the filtrate was evaporated, and the residue purified by flash chromatography (hexanes/EtOAc, 10:1) to give benzofuran 14 as an off-white foam (3.56 g, 84 %). ¹H NMR $(400 \text{ MHz}, \text{ CDCl}_3)$: $\delta = 7.65 \text{ (d}, J = 8.3 \text{ Hz},$ 1H), 7.53 (d, J = 8.4 Hz, 1H), 7.09 (dd, J = 8.4, 2.3 Hz, 1H), 7.03 (d, J=2.3 Hz, 1 H), 6.99 (d, J=8.3 Hz, 1 H), 4.53 (s, 2 H), 3.51-3.55 (m, 2H), 1.45 (s, 9H), 1.38 (s, 9H), 1.17 (s, 9H), 0.92-0.96 (m, 2H), -0.01 ppm (s, 9H); ¹³C NMR (100 MHz, CDCl₃): $\delta = 176.4$ (2C), 176.2, 155.6, 152.8, 151.0, 150.3, 149.4, 132.3, 126.1, 120.5, 120.3, 118.9, 118.0, 117.0, 116.7, 71.9, 67.7, 62.7, 39.4, 39.2, 39.1, 27.4, 27.1 (2C), 18.3, -1.4 ppm; MS (EI): m/z (%): 750 (87) $[M]^+$, 722 (11), 666 (35), 638 (26), 633 (16), 553 (12), 548 (15), 464 (27), 380 (31), 85 (12), 73 (25), 57 (100); HRMS (ESI): m/z: calcd for $C_{35}H_{47}IO_8Si + Na$: 773.1977; found: 773.1969; elemental analysis calcd (%) for C₃₅H₄₇IO₈Si: C 56.00, H 6.31; found: C 55.95, H 6.35.

Received: February 28, 2007 Published online: May 10, 2007

Keywords: alkynes · antibiotics · heterocycles · natural products · platinum

- For an excellent review on the current status of research in the field of antibiotics, see F. von Nussbaum, M. Brands, B. Hinzen, S. Weigand, D. Häbich, *Angew. Chem.* 2006, 118, 5194–5254; *Angew. Chem. Int. Ed.* 2006, 45, 5072–5129, and references therein.
- [2] Leading references: a) G. Belofsky, R. Carreno, K. Lewis, A. Ball, G. Casadei, G. P. Tegos, J. Nat. Prod. 2006, 69, 261-264;
 b) T. Hatano, Y. Shintani, Y. Aga, S. Shiota, T. Tsuchiya, T. Yoshida, Chem. Pharm. Bull. 2000, 48, 1286-1292; c) C. Selvam, S. M. Jachak, R. G. Oli, R. Thilagavathi, A. K. Chakraborti, K. K. Bhutani, Tetrahedron Lett. 2004, 45, 4311-4314, and references therein.
- [3] For the remarkable activity of the prenylated pterocarpene 6 against *Streptococcus mutans*, the pathogene causing caries, see J. He, L. Chen, D. Heber, W. Shi, Q.-Y. Lu, *J. Nat. Prod.* 2006, 69, 121–124.
- [4] a) H. Tanaka, M. Sato, T. Oh-Uchi, R. Yamaguchi, H. Etoh, H. Shimizu, M. Sako, H. Takeuchi, *Phytomedicine* 2004, *11*, 331–337; b) H. Tanaka, T. Oh-Uchi, H. Etoh, M. Sako, M. Sato, T. Fukai, Y. Tateishi, *Phytochemistry* 2003, *63*, 597–602; c) H. Tanaka, M. Sato, S. Fujiwara, M. Hirata, H. Etoh, H. Takeuchi, *Lett. Appl. Microbiol.* 2002, *35*, 494–498.
- [5] A. Fürstner, P. W. Davies, J. Am. Chem. Soc. 2005, 127, 15024– 15025.
- [6] a) For a similar approach, see I. Nakamura, Y. Mizushima, Y. Yamamoto, J. Am. Chem. Soc. 2005, 127, 15022–15023; b) for

related benzofuran syntheses induced by other electrophiles, see the following for leading references and the references therein: D. Yue, T. Yao, R. C. Larock, *J. Org. Chem.* **2005**, *70*, 10292–10296; c) G. Balme, E. Bossharth, N. Monteiro, *Eur. J. Org. Chem.* **2003**, 4101–4111.

- [7] For recent syntheses of other bioactive heterocycles by our research group, see a) A. Fürstner, Angew. Chem. 2003, 115, 3706-3728; Angew. Chem. Int. Ed. 2003, 42, 3582-3603; b) A. Fürstner, K. Radkowski, H. Peters, Angew. Chem. 2005, 117, 2837-2841; Angew. Chem. Int. Ed. 2005, 44, 2777-2781; c) A. Fürstner, M. M. Domostoj, B. Scheiper, J. Am. Chem. Soc. 2006, 128, 8087-8094; d) A. Fürstner, K. Radkowski, H. Peters, G. Seidel, C. Wirtz, R. Mynott, C. W. Lehmann, Chem. Eur. J. 2007, 13, 1929-1945; e) A. Fürstner, F. Feyen, H. Prinz, H. Waldmann, Tetrahedron 2004, 60, 9543-9558, and references therein.
- [8] a) Review: W. Stadlbauer, T. Kappe, *Heterocycles* 1993, 35, 1425–1440; b) leading references on more recent syntheses: B. A. Chauder, A. V. Kalinin, N. J. Taylor, V. Snieckus, *Angew. Chem.* 1999, 111, 1528–1530; *Angew. Chem. Int. Ed.* 1999, 38, 1435–1438; c) C. C. Li, Z. X. Xie, Y. D. Zhang, J. H. Chen, Z. Yang, *J. Org. Chem.* 2003, 68, 8500–8504; d) T. Yao, D. Yue, R. C. Larock, *J. Org. Chem.* 2005, 70, 9985–9989, and references therein.
- [9] a) For a review on the use of Pt^{II}, Au^I, and related late-transitionmetal templates as π-acidic catalysts, see A. Fürstner, P. W. Davies, *Angew. Chem.* 2007, *119*, 3478–3519; *Angew. Chem. Int. Ed.* 2007, 46, 3410–3449; b) C. Nieto-Oberhuber, S. López, E. Jiménez-Núñez, A. M. Echavarren, *Chem. Eur. J.* 2006, *12*, 5916–5923.
- [10] Although the mechanism shown captures the key features of the transformation, the exact mechanism and the identity of the intermediates involved in these shift processes remains under study.
- [11] F. L. Weitl, J. Org. Chem. 1976, 41, 2044-2045.
- [12] K. Hiroya, N. Suzuki, A. Yasuhara, Y. Egawa, A. Kasano, T. Sakamoto, *Perkin 1* 2000, 4339–4346.
- [13] a) A. Elangovan, Y.-H. Wang, T.-I. Ho, Org. Lett. 2003, 5, 1841– 1844; b) review: K. Sonogashira in Metal-catalyzed Cross-

coupling Reactions (Eds.: F. Diederich, P. J. Stang), Wiley-VCH, Weinheim, **1998**, pp. 203-229.

- [14] For the accelerating effect of CO on PtCl₂-catalyzed reactions, see also a) A. Fürstner, P. W. Davies, T. Gress, *J. Am. Chem. Soc.* 2005, *127*, 8244–8245; b) A. Fürstner, C. Aïssa, *J. Am. Chem. Soc.* 2006, *128*, 6306–6307.
- [15] a) A. Fürstner, H. Szillat, B. Gabor, R. Mynott, J. Am. Chem. Soc. 1998, 120, 8305-8314; b) A. Fürstner, F. Stelzer, H. Szillat, J. Am. Chem. Soc. 2001, 123, 11863-11869; c) V. Mamane, T. Gress, H. Krause, A. Fürstner, J. Am. Chem. Soc. 2004, 126, 8654-8655; d) A. Fürstner, P. Hannen, Chem. Eur. J. 2006, 12, 3006-3019.
- [16] a) V. Mamane, P. Hannen, A. Fürstner, *Chem. Eur. J.* 2004, 10, 4556–4575; b) A. Fürstner, J. W. J. Kennedy, *Chem. Eur. J.* 2006, 12, 7398–7410; c) A. Fürstner, V. Mamane, *Chem. Commun.* 2003, 2112–2113.
- [17] G. A. Molander, Chem. Rev. 1992, 92, 29-68.
- [18] The phasing is critical: exploratory experiments suggest that the oxidative cleavage of the trimethylsilylethyl ether is incompatible with a prenyl group already in place, which partly converts into a conjugated diene in the presence of CAN in aqueous MeCN.
- [19] a) F. F. Kneisel, M. Dochnahl, P. Knochel, Angew. Chem. 2004, 116, 1032-1036; Angew. Chem. Int. Ed. 2004, 43, 1017-1021;
 b) L.-Z. Gong, P. Knochel, Synlett 2005, 267-270; c) F. F. Kneisel, H. Leuser, P. Knochel, Synthesis 2005, 2625-2629.
- [20] a) N. T. Keen, A. I. Zaki, J. J. Sims, *Phytochemistry* 1972, 11, 1031–1039; b) H. Zilg, H. Grisebach, *Phytochemistry* 1968, 7, 1765–1772.
- [21] a) M. J. O'Neill, S. A. Adesanya, M. F. Roberts, *Phytochemistry* 1984, 23, 2704–2705; b) D. Morandi, J. L. Le Quere, *New Phytol.* 1991, 117, 75–79.
- [22] a) M. J. O'Neill, Z. Naturforsch. C 1983, 38, 698-700; b) P. Caballero, C. M. Smith, F. R. Fronczek, N. H. Fischer, J. Nat. Prod. 1986, 49, 1126-1129; c) N. Abe, H. Sato, S. Sakamura, Agric. Biol. Chem. 1987, 51, 349-353.