

Synthetic Methods


Direct Transformation of Secondary Amides into Secondary Amines: Triflic Anhydride Activated Reductive Alkylation**

Kai-Jiong Xiao, Ai-E Wang, and Pei-Qiang Huang*

Amines are an important class of compounds that constitute the major body of bioactive natural products (alkaloids) and pharmaceuticals.^[1] Amides are a class of easily available and highly stable compounds.^[2] Consequently, the synthesis of amines by reductive alkylation of amides is of high relevance in organic synthesis,^[3] and has been the focus of recent research.^[4–6] The transformation of tertiary amides into amines has been achieved by indirect methods via thioamides.^[4] Very recently, the direct transformation of tertiary amides into amines by reductive alkylation has been reported.^[5,6] However, although the reductive mono- or dialkylation of secondary lactams/amides^[7,8] has been reported, a general method for the synthesis of secondary amines, with a variety of substituent patterns, has not yet been described. In addition, the secondary amide also serves as a powerful directing group in C–H activation.^[9] Consequently, the development of a general and direct method for the transformation of secondary amides into secondary amines is highly desirable in synthetic organic chemistry.

As a continuation of our endeavour to develop step-economical syntheses,^[10] we recently reported a direct method for the reductive alkylation of lactams/amides with Grignard and organolithium reagents (Scheme 1).^[5] Herein, we report


a one-pot synthesis of secondary amines by the reductive alkylation of secondary amides with organocerium reagents^[11,12] using trifluoromethanesulfonic anhydride (Tf₂O)^[13,14] as the activation reagent (Scheme 2).



Scheme 2. One-pot transformation of secondary amides into secondary amines.


The conversion of *N*-cyclohexylbenzamide **1a** into amine **2a** was chosen as a prototype reaction (Table 1). A CH₂Cl₂ solution of amide **1a** and 2-fluoropyridine^[13] (2-F-Py, 1.2 equiv) was successively treated with 1.1 molar equivalents of Tf₂O (–78 °C, then 0 °C), 3.0 molar equivalents of RM/

Table 1: Influence of the reducing agent on the reductive alkylation of secondary amides.



Entry	Reductant [H]	Equiv	Yield[%] ^[a]
1	Et ₃ SiH	3.0	no product
2	LiAlH ₄	3.0	86
3	NaBH ₄	3.0	75
4	NaBH ₄ , MeOH	3.0	87
5	NaBH ₄ , MeOH	2.0	88

[a] Yield of the isolated product. 2-F-Py = 2-fluoropyridine; cHex = cyclohexyl.



Scheme 1. One-pot transformation of tertiary amides into amines. Tf₂O = trifluoromethanesulfonic anhydride; DTBMP = 2,6-di-*tert*-butyl-4-methylpyridine.

[*] K.-J. Xiao, Dr. A.-E. Wang, Prof. Dr. P.-Q. Huang
Department of Chemistry and
Fujian Provincial Key Laboratory for Chemical Biology
College of Chemistry and Chemical Engineering
Xiamen University, Xiamen, Fujian 361005 (P.R. China)
E-mail: pqhuang@xmu.edu.cn

[**] The authors are grateful to the National Basic Research Program (973 Program) of China (Grant No. 2010CB833200), the NSF of China (21072160; 20832005), and the Fundamental Research Funds for the Central Universities of China (Grant No. 201112G001) for financial support, and for a Scholarship Award for Excellent Doctoral Student granted by Ministry of Education of China (2010). We are grateful to Prof. Dr. G. M. Blackburn for valuable discussions and to Y.-H. for the preparation of some starting materials.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201204098>.

CeCl₃ (–78 °C), and a reducing agent. It was found that triethylsilane was inactive in this reaction (Table 1, entry 1), while lithium aluminum hydride (Table 1, entry 2) and sodium borohydride (Table 1, entry 3) worked well to provide the desired amine in good yield. NaBH₄ was selected as the reducing agent as it is cheaper, less hazardous, and more easily handled than LiAlH₄. Moreover, the addition of methanol was found to accelerate the reaction (Table 1, entries 4 and 5). Thus, the optimized protocol for the direct conversion of secondary amides into amines was identified as successive treatment of a dichloromethane solution of amide and 2-fluoropyridine (1.2 equiv) with 1.1 molar equivalents of Tf₂O (–78 °C, then 0 °C), and 3.0 molar equivalents of RM/

CeCl₃ (−78°C), followed by reduction with 2.0 molar equivalents of NaBH₄ in methanol.

Under these optimized reaction conditions, the scope of this transformation was studied. As can be seen from Table 2, a variety of secondary amides, including aryl (Table 2,

entries 1–12), alkanoyl (Table 2, entries 13–19), and alkenoyl (Table 2, entry 20) amides, were transformed into the corresponding secondary amines in good to excellent yields. The substituents on the nitrogen atom of the secondary amides did not have much influence on the reaction efficiency, regardless

Table 2: Direct transformation of secondary amides **1** into amines **2**.


$ \begin{array}{c} \text{O} \\ \parallel \\ \text{R}^1-\text{C}-\text{N}-\text{R}^2 \\ \text{H} \\ \mathbf{1} \end{array} \xrightarrow[\begin{array}{c} \text{2) RM/ CeCl}_3 \text{ (3.0 equiv)} \\ \text{3) NaBH}_4, \text{ MeOH} \end{array}]{ \begin{array}{c} \text{1) Tf}_2\text{O (1.1 equiv)} \\ \text{2-F-Py (1.2 equiv), CH}_2\text{Cl}_2 \end{array} } \begin{array}{c} \text{R} \\ \\ \text{R}^1-\text{C}-\text{N}-\text{R}^2 \\ \text{H} \\ \mathbf{2} \end{array} $							
Entry	Substrate	RM	Product (Yield [%]) ^[a]	Entry	Substrate	RM	Product (Yield [%]) ^[a]
1		<i>n</i> BuLi		11		<i>n</i> BuMgBr	
	1 a		2 a (88)		1 g		2 j (84)
2		<i>i</i> PrMgBr		12		<i>n</i> BuMgBr	
	1 a		2 b (84)		1 h		2 k (79)
3		PhMgBr		13		<i>n</i> BuMgBr	
	1 a		2 c (82)		1 i		2 l (89)
4		Ph≡Li		14		<i>n</i> BuMgBr	
	1 a		2 d (79)		1 j		2 m (90)
5		<i>n</i> BuLi		15		<i>n</i> BuMgBr	
	1 b		2 e (86)		1 k		2 n (64)
6		<i>n</i> BuMgBr		16		<i>n</i> BuLi	
	1 b		2 e (91)		1 l		2 o (82)
7		<i>n</i> BuMgBr		17		BnMgBr	
	1 c		2 f (85)		1 l		2 p (84)
8		<i>n</i> BuLi		18		<i>n</i> BuMgBr	
	1 d		2 g (75)		1 m		2 q (76)
9		<i>n</i> BuLi		19		BnMgBr	
	1 e		2 h (68)		1 n		2 r (81)
10		<i>n</i> BuMgBr		20		<i>n</i> BuMgBr	
	1 f		2 i (81)		1 o		2 s (75)

[a] Reaction conditions: 1) amide (1.0 mmol), 2-fluoropyridine (1.2 mmol), Tf₂O (1.1 mmol), CH₂Cl₂ (4 mL), −78°C, then 0°C, 10 min; 2) RM/ CeCl₃ (3.0 mmol), −78°C, 1 h; 3) NaBH₄ (2.0 mmol), MeOH (5 mL), RT, 3 h. [b] Yield of the isolated product. Bn = benzyl.

of their identity: *n*-alkyl (Table 2, entries 5–7, 11–13, 15, 18–20), *sec*-alkyl (Table 2, Table 2, entries 1–4, 8–10, 16–17), or aryl (Table 2, entry 14). The reaction also went smoothly with hindered amides, such as **1m** (Table 2, entry 18). Functional groups such as an ether (Table 2, entry 9), an aromatic bromide (Table 2, entry 10), thiophene (Table 2, entry 12), a terminal C=C double bond (Table 2, entry 19), and a conjugated C=C double bond (Table 2, entry 20) were well tolerated. It should be noted that the reaction of α,β -unsaturated amides led to a highly selective 1,2-addition (Table 2, entry 20).

With regard to the organocerium reagents, alkyl (Table 2, entries 1, 2, 5–16, 18, 20), benzyl (Table 2, entries 17 and 19), aryl (Table 2, entry 3), and alkynyl (Table 2, entry 4) cerium reagents, generated from either organolithiums or Grignard reagents, worked well for the direct conversion of secondary amides into secondary amines.

A plausible mechanism for the direct transformation of secondary amides **1** into secondary amines **2** is shown in Scheme 3. The reaction may involve the formation of the highly electrophilic nitrilium ion intermediate **A**^[11] upon the action of Tf₂O/2-fluoropyridine (2-F-Py). Trapping of the nitrilium ion **A** with an organocerium reagent (RM/CeCl₃) forms ketimine **3**,^[15] which upon further reduction gives the corresponding secondary amine.



Scheme 3. Proposed reaction mechanism.

In summary, the first general method for the direct transformation of secondary amides into secondary amines with different substituent patterns has been developed. The significance and advantages of the method are: 1) the method is general, allowing the introduction of a variety of substituents; 2) organocerium reagents with high nucleophilicity yet attenuated basicity are used as the alkylating reagents, and are generated in situ from either Grignard reagents and organolithium reagents; 3) both LiAlH₄ and NaBH₄ can be used as the reducing reagent; 4) the ketimines **3** can also be isolated.^[15] Considering the broad spectrum of bioactivities exhibited by amines, along with the high stability, easy availability, and versatile uses of the secondary amide reactants, this versatile and mild method should find application in the synthesis of *N*-containing bioactive molecules and medicinal agents.

Experimental Section

General procedure for the direct transformation of secondary amides **1** into secondary amines **2**. Tf₂O (185 μ L, 1.1 mmol, 1.1 equiv) was added dropwise to a cooled (-78°C) solution of amide **1** (1.0 mmol, 1.0 equiv) and 2-fluoropyridine (103 μ L, 1.2 mmol, 1.2 equiv) in dichloromethane (4 mL). The reaction was warmed to 0°C in an ice bath and stirred for 10 min. The mixture was then cannulated into freshly prepared organocerium reagent (3.0 mmol, 3.0 equiv) in THF (15 mL) at -78°C and stirred for 1 h. Methanol (5 mL) and sodium borohydride (76 mg, 2.0 mmol) were added. The reaction mixture was warmed to RT and stirred for 3 h. The reaction was quenched with a saturated aqueous NaHCO₃ solution and extracted with dichloromethane (3×10 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel (eluting with 5–10% ethyl acetate in *n*-hexane) to afford the desired amine **2**.

Received: May 26, 2012

Published online: ■ ■ ■ ■ ■, ■ ■ ■ ■ ■

Keywords: amide activation · amines · C–C bond formation · cerium · synthetic methods

- [1] a) S. A. Lawrence, *Amines: Synthesis Properties, and Application*, Cambridge University Press, Cambridge, **2004**; for the synthesis of secondary amines, see: b) R. N. Salvatore, C. H. Yoon, K. W. Jung, *Tetrahedron* **2001**, *57*, 7785–7811.
- [2] W.-J. Yoo, C.-J. Li, *J. Am. Chem. Soc.* **2006**, *128*, 13064–13065, and references cited therein.
- [3] a) M. Schiess, Diss. ETH Nr. 7935, ETH Zürich, **1986**; b) D. Seebach, *Angew. Chem.* **2011**, *123*, 99–105; *Angew. Chem. Int. Ed.* **2011**, *50*, 96–101.
- [4] a) T. Murai, Y. Mutoh, Y. Ohta, M. Murakami, *J. Am. Chem. Soc.* **2004**, *126*, 5968–5969; b) T. Murai, F. Asai, *J. Am. Chem. Soc.* **2007**, *129*, 780–781; c) T. Murai, F. Asai, *J. Org. Chem.* **2008**, *73*, 9518–9521; d) T. Murai, K. Ui, Narengerile, *J. Org. Chem.* **2009**, *74*, 5703–5706; e) A. Agosti, S. Britto, P. Renaud, *Org. Lett.* **2008**, *10*, 1417–1420.
- [5] a) K.-J. Xiao, J.-M. Luo, K.-Y. Ye, Y. Wang, P.-Q. Huang, *Angew. Chem.* **2010**, *122*, 3101–3104; *Angew. Chem. Int. Ed.* **2010**, *49*, 3037–3040; b) K.-J. Xiao, Y. Wang, K.-Y. Ye, P.-Q. Huang, *Chem. Eur. J.* **2010**, *16*, 12792–12796.
- [6] a) K. Shirokane, Y. Kurosaki, T. Sato, N. Chida, *Angew. Chem.* **2010**, *122*, 6513–6516; *Angew. Chem. Int. Ed.* **2010**, *49*, 6369–6372; b) G. Vincent, R. Guillot, C. Kouklovsky, *Angew. Chem.* **2011**, *123*, 1386–1389; *Angew. Chem. Int. Ed.* **2011**, *50*, 1350–1353.
- [7] During the course of the preparation of this manuscript, Chida et al. reported a direct method for the reductive allylation of amides, see: Y. Oda, T. Sato, N. Chida, *Org. Lett.* **2012**, *14*, 950–953.
- [8] a) Y. N. Bubnov, F. V. Pastukhov, I. V. Yampolsky, A. V. Ignatenko, *Eur. J. Org. Chem.* **2000**, 1503–1505; b) Z. Li, Y. Zhang, *Tetrahedron Lett.* **2001**, *42*, 8507–8510.
- [9] a) M. Wasa, K. M. Engle, D. W. Lin, E. J. Yoo, J.-Q. Yu, *J. Am. Chem. Soc.* **2011**, *133*, 19598–19601; b) X. Wang, D. Leow, J.-Q. Yu, *J. Am. Chem. Soc.* **2011**, *133*, 13864–13867; c) S. L. Zultanski, G. C. Fu, *J. Am. Chem. Soc.* **2011**, *133*, 15362–15364; d) Q. Chen, L. Ilies, E. Nakamura, *J. Am. Chem. Soc.* **2011**, *133*, 428–429.
- [10] a) C.-P. Xu, Z.-H. Xiao, B.-Q. Zhuo, Y.-H. Wang, P.-Q. Huang, *Chem. Commun.* **2010**, *46*, 7834–7836; b) R.-F. Yang, P.-Q. Huang, *Chem. Eur. J.* **2010**, *16*, 10319–10322; for an account, see: c) P.-Q. Huang, *Synlett* **2006**, 1133–1149.

- [11] For the selection of organocerium reagent and application to ketone synthesis, see: K.-J. Xiao, A.-E. Wang, Y.-H. Huang, P.-Q. Huang, submitted.
- [12] For reviews on the chemistry of organocerium reagents, see: a) H.-J. Liu, K.-S. Shia, X. Shang, B.-Y. Zhu, *Tetrahedron* **1999**, 55, 3803–3830; b) G. Bartoli, E. Marcantoni, M. Marcolini, L. Sambri, *Chem. Rev.* **2010**, 110, 6104–6143.
- [13] For reviews on the chemistry of triflic acid and its derivatives, see: a) P. J. Stang, M. R. White, *Aldrichimica Acta* **1983**, 16, 15–22; b) I. L. Baraznenok, V. G. Nenajdenko, E. S. Balenkova, *Tetrahedron* **2000**, 56, 3077–3119; for selected examples on the use of TiF_3O as a powerful amide activator for a variety of transformations, see: c) A. B. Charette, P. Chua, *J. Org. Chem.* **1998**, 63, 908–909; d) G. E. Keck, M. D. McLaws, T. T. Wager, *Tetrahedron* **2000**, 56, 9875–9883; e) C. E. McDonald, A. M. Galka, A. I. Green, J. M. Keane, J. E. Kowalchick, C. M. Micklitsch, D. D. Wisnoski, *Tetrahedron Lett.* **2001**, 42, 163–166; f) A. B. Charette, M. Grenon, *Can. J. Chem.* **2001**, 79, 1694–1703; g) A. B. Charette, M. Grenon, A. Lemire, M. Pourashraf, J. Martel, *J. Am. Chem. Soc.* **2001**, 123, 11829–11830; h) M. Movassaghi, M. D. Hill, *J. Am. Chem. Soc.* **2006**, 128, 4592–4593; i) G. Barbe, A. B. Charette, *J. Am. Chem. Soc.* **2008**, 130, 18–19; j) J. W. Medley, M. Movassaghi, *J. Org. Chem.* **2009**, 74, 1341–1344; k) G. Pelletier, W. S. Bechara, A. B. Charette, *J. Am. Chem. Soc.* **2010**, 132, 12817–12819; l) W. S. Bechara, G. Pelletier, A. B. Charette, *Nat. Chem.* **2012**, 4, 228–234.
- [14] a) G. Bélanger, R. Larouche-Gauthier, F. Ménard, M. Nantel, F. Barabé, *J. Org. Chem.* **2006**, 71, 704–712; b) M. Movassaghi, M. D. Hill, *J. Am. Chem. Soc.* **2006**, 128, 14254–14255; c) M. Movassaghi, M. D. Hill, O. K. Ahmad, *J. Am. Chem. Soc.* **2007**, 129, 10096–10097; d) R. Larouche-Gauthier, G. Bélanger, *Org. Lett.* **2008**, 10, 4501–4504; e) F. Lévesque, G. Bélanger, *Org. Lett.* **2008**, 10, 4939–4942; f) S.-L. Cui, J. Wang, Y.-G. Wang, *J. Am. Chem. Soc.* **2008**, 130, 13526–13527; g) G. Bélanger, J. Boudreault, F. Lévesque, *Org. Lett.* **2011**, 13, 6204–6207; h) G. Bélanger, G. O'Brien, R. Larouche-Gauthier, *Org. Lett.* **2011**, 13, 4268–4271; i) G. Bélanger, M. Dupuis, R. Larouche-Gauthier, *J. Org. Chem.* **2012**, 77, 3215–3221.
- [15] Ketimines **3a–3d** were isolated and characterized (see the Supporting information for details).



Communications




Synthetic Methods

K.-J. Xiao, A.-E. Wang,

P.-Q. Huang*    

Direct Transformation of Secondary
Amides into Secondary Amines: Triflic
Anhydride Activated Reductive Alkylation



Versatile and mild: The first general method for the title transformation has been developed (see scheme; 2-F-Py = 2-fluoropyridine; Tf = trifluorosulfonyl). The amines are synthesized in good yields

and the ketimine intermediates can be isolated before the reduction. This method should find applications in the synthesis of nitrogen-containing bio-active molecules and medicinal agents.