

Mechanism of Skeletal Isomerization of *n*-Butane Using 1, 4-¹³C₂-*n*-Butane on Solid Strong Acids

Tetsuo Suzuki and Toshio Okuhara*

Graduate School of Environmental Earth Science, Hokkaido University, Sapporo 060-0810

(Received February 3, 2000; CL-000117)

Mechanism of skeletal isomerization of *n*-butane over Cs_{2.5}H_{0.5}PW₁₂O₄₀ as well as sulfated ZrO₂ was revealed at a temperature range 393 - 523 K by using 1,4-¹³C₂-*n*-butane. ¹³C-Distributions of isobutane at 393 K were close to binomial distribution, indicating a bimolecular pathway. On the other hand, an intramolecular (monomolecular) rearrangement became significant over Cs_{2.5}H_{0.5}PW₁₂O₄₀ at 523 K.


Skeletal isomerization of *n*-butane is practically important, because the product isobutane is a raw material for alkylation with butenes to form clean gasoline (C8 branched alkanes), and the dehydrogenation product, isobutylene can be transformed into methyl *tert*-butyl ether (MTBE). There are many reports about the catalysts active and selective for this reaction. Liquid acids like HF are effective,¹ but these have problems of environmental protection, etc. As solid acids, sulfated ZrO₂,^{2,4} a Cs hydrogen salt of H₃PW₁₂O₄₀, Cs_{2.5}H_{0.5}PW₁₂O₄₀,⁵⁻⁷ and their Pt-promoted catalysts^{8,9} have been demonstrated to be promising catalysts.

Since the selectivity to isobutane is closely related to the reaction mechanism, elucidation of mechanism is required for development of prominent catalysts. Matsushashi et al.¹⁰ proposed that monomolecular and bimolecular mechanisms operated on sulfated ZrO₂ at the initial and latter stage of the reaction, respectively. Recently, isotopic studies have intently been performed on sulfated ZrO₂,^{11,12} but there is a controversy on the mechanism. Garin et al.¹¹ claimed that this reaction occurred through monomolecular mechanism at 523 K and Adeeva et al.¹² inferred that it proceeded through bimolecular mechanism at 353 K, while these discussions are qualitative because of strong influence of fragmentation in mass analysis.

In the present study, we chose Cs_{2.5}H_{0.5}PW₁₂O₄₀ as well as sulfated ZrO₂ as solid acids, since the former possesses pure and very strong protonic acids available for this reaction,⁵ and the latter is exceptionally active at low temperatures for this reaction.^{2,4} Zeolites are active, but are non-selective.⁶

The isotopic composition was analyzed by Field-Ionization Mass Spectrometry. By this, we obtained the parent peak pattern which makes possible to discuss quantitatively. Here we report first the reaction mechanism of *n*-butane isomerization over Cs_{2.5}H_{0.5}PW₁₂O₄₀ as well as sulfated ZrO₂.

In the monomolecular mechanism, the isomerization would involve a protonated cyclopropane and a sequent primary carbenium ion as intermediates to form isobutane with 100%-selectivity.¹³ Reaction paths via the monomolecular mechanism are shown in Scheme 1, when 1,4-¹³C₂-*n*-butane was used as a reactant. ¹³C₂-Isobutane would be produced exclusively (intramolecular rearrangement), together with 1,3-¹³C₂-*n*-butane as an isotopomer by self-isomerization. On the other



Scheme 1. Reaction pathways for skeletal isomerization of 1,4-¹³C₂-*n*-butane.

hand, the bimolecular mechanism (Scheme 1) is possible if butenes and *sec*-butyl carbenium ion are formed on the catalyst surface to form octyl cation.¹³ The β -scission of octyl cation would give C₄ moieties, together with C₃ and C₅ hydrocarbons as byproducts. In this case, the intermolecular isotopic scrambling in isobutane is expected. Thus we can distinguish two mechanisms by the ¹³C isotopic distribution, intramolecular or intermolecular pattern.

The reaction was performed in a closed circulation system (300 cm³) equipped with an on-line GC at 393–523 K. After Cs_{2.5}H_{0.5}PW₁₂O₄₀ (110 m² g⁻¹, abbreviated as Cs2.5) and sulfated ZrO₂ (90 m² g⁻¹, abbreviated as SZ) were pretreated in a vacuum at 573 and 673 K, respectively, 40 torr of 1,4-¹³C₂-*n*-butane (Isotec Inc., ¹³C: 99%) was introduced. The product isobutane and reactant *n*-butane were separated to be analyzed by Field Ionization Mass Spectrometry (FI-MASS, JEOL JMS-SX102A) for ¹³C-distribution.

The selectivity to isobutane for Cs2.5 was changed from 91% to 87% when the reaction temperature was raised from 393 K to 523 K, while it decreased from 92% to 78% over SZ. Figure 1 shows the isotopic distributions of isobutane over Cs2.5, where the binomial patterns were optimized to fit the




Figure 1. ^{13}C -Distributions of isobutane in the isomerization of 1,4- $^{13}\text{C}_2$ - n -butane over $\text{Cs}_{2.5}\text{H}_{0.5}\text{PW}_{12}\text{O}_{40}$. A: 393 K (Conv. =10%) and B: 523 K (Conv. =11%).

observed values of $x = 0, 1, 3$, and 4 in $^{13}\text{C}_x^{12}\text{C}_{4-x}\text{H}_{10}$. As was not shown here, n -butane consisted mainly of $^{13}\text{C}_2$ - n -butane. It was observed that the isotopic distribution in isobutane at 393 K was close to a binomial distribution, while the fraction of $^{13}\text{C}_2$ -isobutane was slightly higher than that of the binomial distribution (Figure 1A). This indicates that at the low temperature, the isomerization on $\text{Cs}_{2.5}$ is the intermolecular process (bimolecular pathway), which is in accordance with the results with SZ.¹² On the other hand, as shown in Figure 1B, the fraction of $^{13}\text{C}_2$ -isobutane was much greater than that of the binomial one at 523 K. The ^{13}C -distributions in isobutane over SZ were essentially the same as those over $\text{Cs}_{2.5}$, while the patterns of ^{13}C -distribution were more close to the corresponding binomial distributions.

In Figure 2, the ratios (R) of contribution of monomolecular pathway to that of bimolecular pathway are plotted against the reaction temperature, where the ratio was estimated from the fraction of $^{13}\text{C}_2$ -isobutane which exceeded that of the binomial distribution. The ratios were 0.13 and 0.04 for $\text{Cs}_{2.5}$ and SZ at 393 K, respectively, demonstrating that the reaction at 393 K occurred exclusively via bimolecular mechanism. The ratio increased greatly as the reaction temperature increased and reached about 0.8 and 0.4 at 523 K over $\text{Cs}_{2.5}$ and SZ, respectively, indicating that the contribution of the monomolecular mechanism became significant on $\text{Cs}_{2.5}$ at the high temperature. This temperature dependence explains the controversy of the previous results for SZ.^{11,12} It should be noted that the contribution of monomolecular mechanism was always higher over $\text{Cs}_{2.5}$ than over SZ. This is accordance with the presence of




Figure 2. Ratio (R) of contribution of monomolecular pathway to that of bimolecular pathway. $\text{Cs}_{2.5}\text{H}_{0.5}\text{PW}_{12}\text{O}_{40}$ ($\circ$) and $\text{SO}_4^{2-}/\text{ZrO}_2$ ($\bullet$). The conversions were around 10%.

the larger amount of strong acid sites on $\text{Cs}_{2.5}$ than on SZ.¹⁴ The trend that SZ preferred bimolecular mechanism is probably due to the oxidative dehydrogenation ability of SZ,³ by which n -butane is dehydrogenated to butenes over SZ to change readily to butyl carbenium ion, and then it forms octyl cation intermediate of bimolecular mechanism.

References

- 1 G. A. Olah, G. K. S. Parakash, and J. Sommer, "Superacids", Wiley-Interscience, New York (1985).
- 2 M. Hino and K. Arata, *J. Am. Chem. Soc.*, **101**, 6439 (1979).
- 3 K. T. Wan, C. B. Khouw, and M. E. Davis, *J. Catal.*, **158**, 311 (1996).
- 4 C.-Y. Hsu, C. R. Heimbach, C. T. Armes, and B. C. Gates, *J. Chem. Soc., Chem. Commun.*, **1992**, 1645.
- 5 T. Okuhara, N. Mizuno, and M. Misono, *Advan. Catal.*, **41**, 141 (1995).
- 6 K. Na, T. Okuhara, and M. Misono, *J. Chem. Soc., Faraday Trans. 1*, **91**, 367 (1995).
- 7 N. Essayem, G. Coudurier, M. Fournier, and J. C. Vedrine, *Catal. Lett.*, **34**, 223 (1995).
- 8 K. Ebitani, J. Konishi, and H. Hattori, *J. Catal.*, **131**, 257 (1991).
- 9 K. Na, T. Okuhara, and M. Misono, *J. Catal.*, **170**, 96 (1997).
- 10 H. Matsushashi, H. Shibata, H. Nakamura, and K. Arata, *Appl. Catal. A: General*, **187**, 99 (1999).
- 11 F. Garin, L. Seyfried, P. Gitard, G. Maire, A. Abdulsamad, and J. Sommer, *J. Catal.*, **151**, 26 (1995).
- 12 V. Adeeva, G. D. Lei, and W. M. H. Sachtler, *Catal. Lett.*, **33**, 135 (1995).
- 13 B. C. Gates, "Catalytic Chemistry," John Wiley & Sons, Inc., New York (1992).
- 14 T. Okuhara, T. Nishimura, and M. Misono, *Stud. Surf. Sci. Catal.*, **101**, 581 (1996).