1963, *2*, 683; b) less basic organocerium reagents also exhibit azaenolization: D. Enders, E. Diez, R. Fernandez, E. Martin-Zamora, J. M. Munoz, R. R. Pappalardo, J. M. Lassaleta, *J. Org. Chem.* **1999**, *64*, 6329.

- [3] For selected recent examples, see: a) review of asymmetric Streckertype reactions: L. Yet, Angew. Chem. 2001, 113, 900; Angew. Chem. Int. Ed. 2001, 40, 875; b) for asymmetric Mannich reactions, see: W. Notz, K. Sakthivel, T. Bui, G. Zhong, C. F. Barbas III, Tetrahedron Lett. 2001, 42, 199; B. List, J. Am. Chem. Soc. 2000, 122, 9336; S. Saito, K. Hatanaka, H. Yamamoto, Org. Lett. 2000, 2, 1891; K. Miura, K. Tamaki, T. Nakagawa, A. Hosomi, Angew. Chem. 2000, 112, 2034; Angew. Chem. Int. Ed. 2000, 39, 1958.
- [4] a) G. K. Friestad, J. Qin, J. Am. Chem. Soc. 2000, 122, 8329; b) G. K. Friestad, J. Qin, J. Am. Chem. Soc. 2001, 123, 9922.
- [5] For reviews of allyl organometallic addition to C=N bonds, see: Y. Yamamoto, N. Asao, Chem. Rev. 1993, 93, 2207; E. F. Kleinman, R. A. Volkmann in Comprehensive Organic Synthesis, Vol. 2 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, 1991, pp. 975-1006; see also ref. [1]. For selected recent examples, see:a) allyltin: H. Nakamura, K. Aoyagi, J.-G. Shim, Y. Yamamoto, J. Am. Chem. Soc. 2001, 123, 372; F. M. Guerra, M. R. Mish, E. M. Carreira, Org. Lett. 2000, 2, 4265; J. A. Marshall, K. Gill, B. M. Seletsky, Angew. Chem. 2000, 112, 983; Angew. Chem. Int. Ed. 2000, 39, 953; b) allylboron: K. Watanabe, S. Kuroda, A. Yokoi, K. Ito, S. Itsuno, J. Organomet. Chem. 1999, 581, 103; c) allyltitanium: X. Teng, Y. Takayama, S. Okamoto, F. Sato, J. Am. Chem. Soc. 1999, 121, 11916; d) allylmagnesium: S. Maiorana, C. Baldoliu, P. D. Buttero, E. Licandro, A. Papagni, M. Lanfranchi, A. Tiripicchio, J. Organomet. Chem. 2000, 593-594, 380; A. G. Steinig, D. M. Spero, J. Org. Chem. 1999, 64, 2406; e) allylsamarium: R. Yanada, M. Okaniwa, A. Kaieda, T. Ibuka, Y. Takemoto, J. Org. Chem. 2001, 66, 1283; f) allylindium: H. M. S. Kumar, S. Anjaneyulu, E. J. Reddy, J. S. Yadav, Tetrahedron Lett. 2000, 41, 9311; T. H. Chan, W. Lu, Tetrahedron Lett. 1998, 39, 8605; g) allylgermanium: T. Akiyama, J. Iwai, Y. Onuma, H. Kagoshima, Chem. Commun. 1999, 2191.
- [6] a) X. Fang, M. Johannsen, S. Yao, N. Gathergood, R. G. Hazell, K. A. Jorgensen, J. Org. Chem. 1999, 64, 4844; b) D.-K. Wang, Y.-G. Zhou, Y. Tang, X.-L. Hou, L.-X. Dai, J. Org. Chem. 1999, 64, 4233; c) A. S. Pilcher, P. DeShong, J. Org. Chem. 1996, 61, 6901.
- [7] a) Sluggish reactivity of allylsilanes toward neutral imines has been noted: see refs. [1] and [5a]; b) although allylsilane addition to iminium ions is effective, a strong Lewis acid such as TiCl₄ or BF₃. Et₂O is often required; for a review, see: I. Fleming, J. Dunogus, R. Smithers, *Org. React.* 1989, *37*, 57; recent examples: J. V. Schaus, N. Jain, J. S. Panek, *Tetrahedron* 2000, *56*, 10263; T. Akiyama, M. Sugano, H. Kagoshima, *Tetrahedron Lett.* 2001, *42*, 3889.
- [8] A. S. Pilcher, H. L. Ammon, P. DeShong, J. Am. Chem. Soc. 1995, 117, 5166; C. J. Handy, Y.-F. Lam, P. DeShong, J. Org. Chem. 2000, 65, 3542.
- [9] The optical rotation of $4([\alpha]_{20}^{20} = -2.0 (c = 3.4 \text{ in CHCl}_3))$ was within experimental error of an authentic sample $([\alpha]_{20}^{20} = -2.2 (c = 3.4 \text{ in CHCl}_3))$ prepared from the known homoallylic amine: T. Basile, A. Bocoum, D. Savoia, A. Umani-Ronchi, *J. Org. Chem.* **1994**, *59*, 7766.
- [10] Hydrogenation of 3g gives 6, identical to that from *n*-propyl radical addition to 2g [Eq. (2)]. See ref. [4b].



- [12] Similar explanations have been advanced regarding the enhanced reactivity of bis(allyl)silanes. A. Shibato, Y. Itagaki, E. Tayama, Y. Hokke, N. Asao, K. Maruoka, *Tetrahedron* 2000, 56, 5373.

Molecular Insight Into Surface Organometallic Chemistry Through the Combined Use of 2D HETCOR Solid-State NMR Spectroscopy and Silsesquioxane Analogues**

Mathieu Chabanas, Elsje Alessandra Quadrelli, Bernard Fenet, Christophe Copéret,* Jean Thivolle-Cazat, Jean-Marie Basset,* Anne Lesage, and Lyndon Emsley*

The structural characterization of heterogeneous catalysts remains difficult. In the late eighties, Feher and later others showed that polyhedral oligomeric silsesquioxanes (POSS) can be used to model the surface structure of partially dehydroxylated silica and therefore infer information about the coordination sphere of metal complexes directly supported on oxides such as silica.^[1] This approach can also provide mechanistic insights on surface chemistry processes. However, no direct comparison between these models and their related surface complexes has been reported.

We have recently shown that the reaction of tris(neopentyl)neopentylidene tantalum, $[Ta(=CHtBu)(CH_2tBu)_3]$ (1) with SiO₂₋₍₅₀₀₎ gives a mixture of mono- $[\equiv$ SiO-Ta(=CHtBu)-(CH₂tBu)₂] and bisgrafted surface complexes $[(\equiv$ SiO)₂Ta(=CHtBu)(CH₂tBu)],^[2a] while its reaction with a SiO₂₋₍₇₀₀₎ provides the monografted species $[\equiv$ SiO-Ta-(=CHtBu)(CH₂tBu)₂] (2_s) as the sole surface species. The characterization was based on elemental analysis, the evolution of neopentane during grafting of 1, and solvolysis of 2_s as well as infrared (IR) spectroscopy.^[2b]

We report here the use of high-resolution solid-state onedimensional (1D) and two-dimensional (2D) NMR spectroscopy to study $\mathbf{2}_{s}$ and the use of high-resolution solution-state 1D and 2D NMR spectroscopy of its molecular analogue $[(c-C_{5}H_{9})_{7}Si_{7}O_{12}Si-O-Ta(=CHtBu)(CH_{2}tBu)_{2}](\mathbf{2}_{m})$ to define the structure of $\mathbf{2}_{s}$ at a molecular level and to investigate the reaction pathway leading to the grafted species.

[*] Dr. C. Copéret, Dr. J.-M. Basset, M. Chabanas, Dr. J. Thivolle-Cazat Laboratoire de Chimie Organométallique de Surface UMR-9986 CNRS - ESCPE Lyon, 43 blvd du 11 Novembre 1918 69626 Villeurbanne Cedex (France) Fax: (+33) 4-7243-1795 Dr. E. A. Quadrelli Dipartimento di Chimica e Chimica Industriale Via Risorgimento 35, 56100 Pisa (Italy) B. Fenet Centre de RMN UCB Lyon 1 - ESCPE Lyon 43 bd du 11 Novembre 1918, 69626 Villeurbanne Cedex (France) Dr. L. Emsley, Dr. A. Lesage Laboratoire de Stéréochimie et des Interactions Moléculaires UMR-117 CNRS - ENS Lyon, Ecole Normale Supérieure de Lyon 69364 Lyon Cedex (France) [**] We are also indebted to the CNRS, ENS Lyon, and ESCPE Lyon for financial support. M.C. is grateful to the French ministry of education,

- research, and technology (MENRT) for a pre-doctoral fellowship. E.A.Q. gratefully acknowledges Università di Pisa and S.N.A.M. for financial support. 2D HETCOR = two-dimensional heteronuclear correlation.
- Supporting information for this article is available on the WWW under http://www.angewandte.com or from the author.

Angew. Chem. Int. Ed. 2001, 40, No. 23 © WILEY-VCH Verlag GmbH, D-69451 Weinheim, 2001 1433-7851/01/4023-4493 \$ 17.50+.50/0

COMMUNICATIONS

As previously described, the surface complex 2_s , was prepared by treating a solution of $[Ta(=CH_tBu)(CH_2tBu)_3]$ in pentane (1.1 equivalents with respect to surface silanols; 0.7-0.8 accessible OH nm⁻²) with SiO₂₋₍₇₀₀₎.^[2b] After workup, 2_s was obtained as a yellow powder typically containing 4.2 % wt of Ta. Noteworthy is the rapid (within 1 min) coloration of silica and discoloration of the supernatant solution upon addition of the pentane solution of **1** to SiO₂₋₍₇₀₀₎. All the observations and chemical analyses are consistent with the proposed structure of 2_s [Eq. (1)], but further corroboration at the molecular level has been sought using both solid-state NMR spectroscopy of the grafted species and solution NMR spectroscopy of molecular model species.

$$1 + \equiv \text{SiOH} \longrightarrow [\equiv \text{SiO}-\text{Ta}(\equiv \text{CH}_{t}\text{Bu})(\text{CH}_{2}t\text{Bu})_{2}] + \text{NpH}$$
(1)
SiO₂₋₍₇₀₀₎ 2_s

To improve the signal-to-noise ratio in the solid-state NMR spectra of **2**, the complex was partially ¹³C enriched at the *a*-position by reaction of $[Ta(=C*HtBu)(C*H_2tBu)_3]$ (**1***) with SiO₂₋₍₇₀₀₎, which yielded $[\equiv$ SiO-Ta(=C*HtBu)(C*H_2tBu)_2] (**2***). The solid-state ¹H NMR spectrum of **2*** shows two broad resonance signals at $\delta = 4.2$ and 1.0 (see the Supporting Information),^[3] and its ¹³C solid-state cross polarization/magic angle spinning (CP/MAS) NMR spectrum displays strong signals at $\delta = 246$, 96.5, and 33.5 (Figure 1 a). Moreover, the 2D HETCOR solid-state NMR spectra provided an unequivocal assignment for **2*** (see Supporting Information for experimental details). The 2D NMR spectrum, recorded with



a contact time of 2 ms (Figure 1 c), shows clear correlations between the signals at $\delta = 4.2$ and 1.0 in the ¹H dimension (F₁) with those at $\delta = 246$ and 33.5, respectively, in the carbon dimension (F₂). The signal at $\delta = 96.5$ also correlates with the peak centered at $\delta = 1.0$, strongly suggesting that the methylene protons are buried under this massive signal (the chemical shifts of the corresponding protons for the surface model **2**_m are indeed around 1 ppm, see Table 1 and below).

A reaction intermediate can be detected if the preparation of 2_s^* involves a rather short (5 min or less) or a cold (below 15–20 °C) drying step; under such conditions, an extra signal appears in the solid-state NMR spectrum at $\delta = 110$ (Figure 1b), which readily disappears upon treating the sample under vacuum or under heat. During this step, neopentane is evolved. All these observations suggest that the intermediate observed in the reaction of $[Ta(=CH_tBu)(CH_2tBu)_3]$ with SiO₂₋₍₇₀₀₎, is the tetraalkyl derivative $[\equiv SiO-Ta(CH_2tBu)_4]$ (**3**_s) which is formed under kinetic control. The intermediate **3**_s rapidly releases neopentane, to finally yield **2**_s as the sole surface complex.

To corroborate the assignments proposed above and to gain further insight in the reaction mechanism suggested by the detection of intermediate 3_s , we have prepared a molecular model of 2_m using the silsesquioxane $(c-C_5H_9)_7Si_7O_{12}SiOH$ (4) a Feher-cage analogue,^[1d,f] that can mimic an isolated silanol of a silica surface partially dehydroxylated at 700 °C. The thermodynamic product of the reaction between the monosilanol 4 and [Ta(=CHtBu)(CH₂tBu)₃] is 2_m (for characteristic data, see Table 1) alongside one equivalent of neopentane.^[4]

The NMR spectroscopy data obtained for 2_m are in good agreement with those of its solid-state counterpart, 2s. Based on NMR spectroscopy data, the reaction is under kinetic control and reaches completion after 48 h. The kinetic monitoring of the reaction shows that a colorless intermediate, [(c-C₅H₉)₇- $Si_7O_{12}SiOTa(CH_2tBu)_4$] (**3**_m) is formed immediately upon mixing of the reagents. The longlived intermediate $\mathbf{3}_{\mathbf{m}}$ ($\tau_{1/2} = 9$ h) has been characterized by 1D and 2D solution NMR spectroscopy, which yields data in good agreement with those obtained for its solid counterpart 3_s (Table 1). The decomposition path of complex $\mathbf{3}_{\mathbf{m}}$ consists of the elimination a neopentane unit by means of α -H abstraction to yield the orange complex $2_{\rm m}$.

The combination of the data obtained in the solid state for the synthesis of 2_s and in solution for 2_m allows a mechanistic pathway to be proposed for the reaction:

4494

¹³C NMR spectrum.

© WILEY-VCH Verlag GmbH, D-69451 Weinheim, 2001

spectra of 2_s^* (2 ms of contact time). d) Proton traces of the signals at $\delta = 246$, 96.5, and 33.5 in the 2D ¹H –

1433-7851/01/4023-4494 \$ 17.50+.50/0

Table 1.	¹ H and	¹³ C chemica	al shifts of	1, 2 _m ,	and 3m	in C	C_6D_6
----------	--------------------	-------------------------	--------------	---------------------	--------	------	----------

Com-	$^{1}H N$	[MR [δ]	13 C NMR [δ]			
plex	resonance ^[a]	assignement	resonance	assignement		
1	0.81	$CH_2C(CH_3)_3$	34.7	$=CHC(CH_3)_3$		
	1.14	$CH_2C(CH_3)_3$	34.95	$CH_2C(CH_3)_3$		
	1.45	$=CHC(CH_3)_3$	35.05	$CH_2C(CH_3)_3$		
	1.89	$=CHC(CH_3)_3$	47.0	$=CHC(CH_3)_3$		
			113.75	$CH_2C(CH_3)_3$		
			249.9	$=CHC(CH_3)_3$		
2 _m	$1.1 - 2.1(m)^{[b]}$	$(c-CH(CH_2)_4)$	22.6, 22.65	$(c-CH(CH_2)_4)$		
			27.4, 27.45, 27.85, 27.9	$(c-CH(CH_2)_4)$		
	1.03(d) ^[c]	$CH_{a}H_{b}C(CH_{3})_{3}$	33.95	$CH_2C(CH_3)_3$		
	1.20(d) ^[c]	$CH_aH_bC(CH_3)_3$	34.4	$=CHC(CH_3)_3$		
	1.20	$CH_2C(CH_3)_3$	34.75	$CH_2C(CH_3)_3$		
	1.41	$=CHC(CH_3)_3$	45.55	$=CHC(CH_3)_3$		
	4.77	$=CHC(CH_3)_3$	95.6	$CH_2C(CH_3)_3$		
			245.2	$=CHC(CH_3)_3$		
3 _m	$1.1 - 2.1(m)^{[b]}$	$(c-CH(CH_2)_4)$	22.7, 22.6,	c - $CH(CH_2)_4)$		
	1.34	$CH_2C(CH_3)_3$	22.55			
	1.87	$CH_2C(CH_3)_3$	27.45, 27.55, 27.75, 27.9	$(c-CH(CH_2)_4)$		
			_[d]	$CH_2C(CH_3)_3$		
			35.2	$CH_2C(CH_3)_3$		
			108.7 ^[b]	$CH_2C(CH_3)_3$		

[a] multiplicity in parenthesis; singlets are not specified. [b] broad signal. [c] J = 14.0 Hz. [d] Not detected.

addition of the silanol to the alkylidene moiety of $1 (1 \rightarrow 3_s)$ followed by an α -H abstraction $(3_s \rightarrow 2_s; \text{Scheme 1})$.^[5]

The reactivities of **4** and SiO₂₋₍₇₀₀₎ are relatively similar (immediate reaction of **1** with silanols and a subsequent slower α -H abstraction step), which is probably because of similar pKa values^[6] and steric crowding. Finally, 2D HET-COR solid-state NMR spectroscopy is an important tool to unequivocally assign ¹H and ¹³C chemical shifts for surface complexes, and will be important in the future development of surface organometallic chemistry. In conclusion, these data taken together show, besides the analogy between surface and solution chemistry, that the combined use of high-resolution solid state NMR and solution NMR spectroscopy to investigate molecular analogues, is a powerful tool in surface organometallic chemistry, and enables the identification of

end products, and determination of the grafting-reaction mechanism. Moreover, we have shown that the Feher cage analogue **4**, is a good model for $SiO_{2-(700)}$, a silica partially dehydroxylated at 700 °C, which contains mainly isolated silanols, and provides a good estimation of the chemical shifts for surface complexes and can help in understanding elementary steps of surface reactions.

Received: April 2, 2001 Revised: September 18, 2001 [Z16881]

- For some examples on the preparation of Feher cages or analogues (POSS) and their reactivity with organometallics, see: a) F. J. Feher, D. N. Newman, J. F. Walzer, J. Am. Chem. Soc. 1989, 111, 1741; F. J. Feher, D. N. Newman, J. Int. Biomed. Inf. Data 1990, 112, 1931; b) F. J. Feher, K. J. Weller, J. J. Schwab, Organometallics 1995, 14, 2009; c) F. J. Feher, T. A. Budzichowski, Polyhedron 1995, 14, 3239; d) R. Duchateau, H. C. L. Abbendhuis, R. A. Van Santen, S. K.-H. Thiele, M. F. H. Van Tol, Organometallics 1998, 17, 5222; e) R. Duchateau, H. C. L. Abbendhuis, R. A. Van Santen, A. Meetsma, S. K.-H. Thiele, M. F. H. Van Tol, Organometallics 1998, 17, 5663; f) R. Duchateau, U. Cremer, R. J. Harmsen, S. I. Mohamud, H. C. L. Abbendhuis, R. A. Van Santen, A. Meetsma, S. K.-H. Thiele, M. F. H. Van Tol, M. Kranenburg, Organometallics 1999, 18, 5447; g) for a review on various surface models other than POSS, see: R. Murugavel, A. Voigt, M. G. Walawalkar, H. W. Roesky, Chem. Rev. 1996, 96, 2205.
- [2] a) V. Dufaud, G. P. Niccolai, J. Thivolle-Cazat, J.-M. Basset, J. Am. Chem. Soc. 1995, 117, 4288–4294; b) L. Lefort, M. Chabanas, O. Maury, D. Meunier, C. Copéret, J. Thivolle-Cazat, J.-M. Basset, J. Organomet. Chem. 2000, 593–594, 96.
- [3] The natural abundance ¹³C NMR spectrum displays three signals at δ = 33.5 (C(CH₃)₃), 96.5 (CH₂C(CH₃)₃), and 246 (=CH-C(CH₃)₃), albeit with a poor signal/noise ratio, especially for the latter. Therefore, 2 was partially ¹³C enriched at the α-position to the metal, 2^{*}_s, by reaction of [Ta(=C*HtBu)(C*H₂tBu)₃] (1*) with SiO₂₋₍₇₀₀₎.
- [4] These values are different from those obtained by Wolczanski and coworkers for [(tBu)₃SiOTa(=CHtBu)(CH₂tBu)₂]:R. E. Lapointe, P. T. Wolczanski, G. D. Van Duyne, *Organometallics* 1985, 4, 1810. This illustrates the dramatic effects on the NMR data of substituents on Si and the necessity of the oxygen atoms around the Si centers.
- [5] This was in fact indirectly shown in the studies of the formation of the surface complexes, see ref. [2a]. Noteworthy is the reaction of HCl on



Scheme 1. Comparison of the reaction of silica partially dehydroxylated at 700 °C (SiO₂₍₋₇₀₀₎) versus (c-C₅H₉)₇Si₇O₁₂SiOH.

Angew. Chem. Int. Ed. 2001, 40, No. 23

© WILEY-VCH Verlag GmbH, D-69451 Weinheim, 2001

Weinheim, 2001 1433-7851/01/4023-4495 \$ 17.50+.50/0

COMMUNICATIONS

[Ta(=CHtBu)(CH₂tBu)₃] which yields Cl-Ta(CH₂tBu)₄: R. R. Schrock, J. D. Fellmann, *J. Am. Chem. Soc.* **1978**, *100*, 3359.

[6] a) For comprehensive information on the pK_a values of silicas, see: *The Surface Properties of Silicas* (Ed.: A. P. Legrand), Wiley, New York, 1998, chapter 4B; b) for related comments on 4, see ref. [1f].



Richard C. D. Brown* and John F. Keily

2,5-Bis(hydroxymethyl)-substituted tetrahydrofurans (THF diols) are an important structural unit found in many natural and synthetic, biologically active molecules.^[1] An elegant approach to the synthesis of THF diols is by the permanganate-promoted oxidative cyclization of 1,5-dienes, which leads to the stereospecific formation of up to four new stereocenters and the THF ring.^[2-6] The same transformation has been carried out using other metal-oxo reagents and catalysts.^[7, 8] In the permanganate-promoted oxidative cyclization reaction, control of absolute stereochemistry has been achieved with chiral auxiliaries;^[4a,b] however, preparation of enantiomerically enriched THF diols by oxidative cyclizations of achiral 1,5-dienes has not been reported. Herein we report the novel high-yielding phase-transfer permanganate-promoted oxidation of geranyl benzoate (1) to the corresponding THF diol 2 (Scheme 1), and a procedure for the asymmetric oxidative cyclization of phenone dienes to produce enantiomerically enriched THF diols.

The use of phase-transfer conditions in permanganatepromoted oxidations is well known,^[9] although we were not aware of application to the oxidative cyclization of 1,5-dienes. Therefore before attempting to develop an asymmetric process, we investigated the phase-transfer oxidative cyclization of a readily available substrate **1** using achiral tertiary ammonium salts (Scheme 1).^[10]

Oxidation of diene **1** with potassium permanganate (2 equiv) in diethyl ether buffered with ethanoic acid gave the best yield of THF diol **2** (70%) when Adogen 464 (0.1–1 equiv) was employed (entry 4, Table 1). Oxidation of enone **4** under the same conditions gave the desired THF diol **5**,

- [**] This work was supported by the EPSRC (J.F.K.) and the Royal Society (R.C.D.B.). We thank Dr. Barry Lygo (University of Nottingham) for helpful discussions, and Syngenta and Merck Sharp and Dohme for unrestricted grants.
- Supporting information for this article is available on the WWW under http://www.angewandte.com or from the author.

4496

© WILEY-VCH Verlag GmbH, D-69451 Weinheim, 2001



Scheme 1. Oxidative cyclization of dienes 1 and 4. a) $0.4 \text{ M} \text{ KMnO}_4$ (2 equiv), AcOH (4 equiv), Adogen 464 (0.4 equiv)/Et₂O. Bz = benzoyl.

Table 1. Oxidative cyclization of geranyl benzoate (1) (see Scheme 1).^[a]

Entry	Solvent	Phase-transfer agent	Yield of 2 [%]	Yield of 3 [%]
1	CH ₂ Cl ₂ ^[b]	TBAB	8 ^[c]	not recorded
2	CH_2Cl_2	TBAB	44 ^[d]	21 ^[d]
3	Et_2O	TBAB	40 ^[d]	7 ^[d]
4	Et_2O	Adogen 464	70 ^[d]	10 ^[d]

[[]a] Reactions were conducted on a 0.19 mmol scale. TBAB = tetrabutylammonium bromide. [b] No AcOH added to the reaction mixture. [c] Yield estimated by HPLC. [d] Yields represent analytically pure isolated material.

albeit in lower yield (Scheme 1). We then returned to our primary goal of developing an asymmetric oxidative cyclization.

Studies on the oxidation of cinnamate esters by quaternary ammonium permanganates led Lee et al. to conclude that the transition state was electron-rich, and they proposed an enolate-like structure **6** with stabilization due to interaction with the ammonium ion (Scheme 2).^[11] The similarity of **6** and the initial adduct **7** formed in the phase-transfer-catalyzed nucleophilic epoxidation of enones suggested that chiral catalysts developed for the epoxidation reaction might also be applied to the oxidative cyclization.^[12]



Scheme 2. Possible reaction pathways for the permanganate-promoted oxidation of electron-poor olefins (top), and the nucleophilic epoxidation of α , β -enones with CIO⁻ (bottom).

Oxidative cyclization of geranyl benzoate (1) in the presence of **8** in dichloromethane/water/ethanoic acid at

1433-7851/01/4023-4496 \$ 17.50+.50/0 Angew. Chem. Int. Ed. 2001, 40, No. 23

^[*] Dr. R. C. D. Brown, J. F. Keily Department of Chemistry University of Southampton Highfield, Southampton SO17 4BJ (UK) Fax: (+44)23-80596805 E-mail: rcb1@soton.ac.uk