

# Naphthalene formation by allylation of zirconaindenes in the $\text{ZnX}_2\text{-Pd(PPh}_3)_4$ system

Zheng Duan,<sup>a</sup> Kiyohiko Nakajima<sup>b</sup> and Tamotsu Takahashi<sup>\*a</sup>

<sup>a</sup> Catalysis Research Center and Graduate School of Pharmaceutical Sciences, Hokkaido University, and CREST, Science and Technology Corporation (JST), Sapporo 060-0811, Japan.

E-mail: tamotsu@cat.hokudai.ac.jp

<sup>b</sup> Department of Chemistry, Aichi University of Education, Igaya, Aichi 448-8542, Japan


Received (in Cambridge, UK) 24th April 2001, Accepted 10th July 2001

First published as an Advance Article on the web 16th August 2001


Zirconaindenes reacted with allyl halides in the presence of  $\text{ZnX}_2$  ( $\text{X} = \text{Cl}$  or  $\text{Br}$ ) and a catalytic amount of  $\text{Pd(PPh}_3)_4$  to give naphthalene derivatives in good yield.

It has been believed for a long time that zirconacyclopentadienes are inert toward C–C bond formation reactions. Recently, we found that transmetalation reactions of zirconacyclopentadienes to copper,<sup>1</sup> nickel<sup>2</sup> and lithium<sup>3</sup> could open a way to various carbon–carbon bond formation reactions. In this paper we would like to report the novel naphthalene formation reaction<sup>4</sup> of zirconaindenes in the presence of  $\text{ZnX}_2$  and a catalytic amount of  $\text{Pd(PPh}_3)_4$ .<sup>5</sup>

When we investigated novel transmetalation to Zn, we tried the allylation reaction of zirconacyclopentadienes, since double allylation of zirconacyclopentadienes<sup>1</sup> proceeded in the presence of either a catalytic or stoichiometric amount of  $\text{CuCl}$ .<sup>1</sup> When only the zinc salts were used, the reactions were very sluggish. However in the presence of a catalytic amount of  $\text{Pd(PPh}_3)_4$  the reaction dramatically changed. Monoallylation products were clearly formed. The formation of double allylation products was not observed [eqn. (1)].



This is in sharp contrast to the allylation in the presence of  $\text{CuCl}$ .<sup>1</sup> Then our attention was turned to the allylation reaction of zirconaindenes. It is interesting to note that, when zirconaindene<sup>6</sup> **3a** was used, substituted naphthalene **4a** was clearly formed as a single product [eqn. (2)].<sup>†</sup> The formation of



regioisomer **5a** was not observed. The use of only a catalytic amount of  $\text{Pd(PPh}_3)_4$  without zinc salt did not give a clean reaction, although **4a** was formed in 33% GC yield.

The structure of the compound **4a** was confirmed by X-ray analysis.<sup>‡</sup> It clearly shows that the allylation reaction proceeded selectively at the benzene-carbon attached to Zr. When  $\text{MeOC}_6\text{H}_4\text{-}$  and  $\text{MeC}_6\text{H}_4\text{-}$  were used as substituents instead of the Ph group, the similar perfect regiocontrol was observed as shown in Table 1. This perfect regioselectivity is attributed to the high reactivity of the benzene-carbon attached to Zr compared with the carbon of the  $\text{-C(Ph)=C(Ph)}$  group attached to Zr. It is interesting to note that when the nucleophilic property

Table 1 The formation of naphthalene derivatives


Zirconacyclopentadienes	Product	Yield (%) <sup>a</sup>
 <b>3a</b> R = Ph <b>3b</b> R = $\text{C}_6\text{H}_4\text{OMe}$ <b>3c</b> R = $\text{C}_6\text{H}_4\text{Me}$	 <b>4a</b> R = Ph <b>4b</b> R = $\text{C}_6\text{H}_4\text{OMe}$ <b>4c</b> R = $\text{C}_6\text{H}_4\text{Me}$	 85(59) 61(42) 68(40)
 <b>3d</b>	 <b>4d</b> + <b>5d</b>	 79(47) 1.9:1
 <b>3e</b>	 <b>4e</b> + <b>5e</b>	 77(51) 3:1

<sup>a</sup> GC yields. Isolated yields are given in parentheses.

of the carbon in the  $\text{-C=C}$  group attached to Zr was enhanced by introducing alkyl groups such as the Et group and the Pr group instead of the Ph groups, carbon–carbon bond formation with allyl halides at the  $\text{-C(R)=C(R)}$ , R = Et or Pr, was observed. The isomers **5d** and **5e** were obtained in the case of **3d** and **3e**, respectively.

A possible mechanism of the naphthalene formation involves (i) transmetalation of the Zr-phenyl carbon bond to Zn giving **6**, (ii) transmetalation of **6** with an allylpalladium halide formed by oxidative addition of allyl halide to a  $\text{Pd}^0$  complex, (iii) reductive elimination to give allylated intermediate **7**, (iv) transmetalation of the second Zr–C bonds to Pd forming **8**, (v) insertion of the vinyl group into the Pd–C bond (vi)  $\beta$ -hydrogen elimination to give *exo*-methylene derivative **10** that isomerizes to the corresponding naphthalene **4** (Scheme 1).

Further investigations on mechanism and selectivity are currently in progress.



**Scheme 1**

## Notes and references

† Typical procedure: to zirconaindene **3a** prepared according to the literature were added 2 eq. of ZnBr<sub>2</sub>, a catalytic amount of Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%) and 2 eq. allyl bromide. The mixture was stirred at 50 °C for 12 h. Only **4a** was formed in 85% GC yield. After isolation and purification by column chromatography, **4a** was obtained in 59% isolated yield.

‡ Crystallographic data of **4a**: colorless prisms, monoclinic, space group C2/c,  $a = 21.312(1)$ ,  $b = 10.6981(6)$ ,  $c = 16.2781(8)$  Å,  $\beta = 117.713(4)^\circ$ ,  $Z = 8$ ,  $R = 0.044$ , GOF = 1.84. CCDC 165561. See <http://www.rsc.org/suppdata/cc/b1/b103674m/> for crystallographic files in .cif or other electronic format.

- 1 T. Takahashi, M. Kotor, K. Kasai and N. Suzuki, *Organometallics*, 1994, **13**, 4183.
- 2 T. Takahashi, F.-Y. Tsai, Y. Li., K. Nakajima and M. Kotor, *J. Am. Chem. Soc.*, 1999, **121**, 11093.
- 3 T. Takahashi, S. Q. Huo, R. Hara, Y. Noguchi, K. Nakajima and W.-H. Sun, *J. Am. Chem. Soc.*, 1999, **121**, 1094.
- 4 For a recent review of naphthalene formation from benzyne complexes, see M. A. Bennett and E. Wenger, *Chem. Ber.*, 1997, **130**, 1029.
- 5 For recent examples of naphthalene formation using transition metal compounds, see D. Peña, D. Pérez, E. Guitián and L. Castedo, *J. Am. Chem. Soc.*, 1999, **121**, 5827; E. Yoshikawa and Y. Yamamoto, *Angew. Chem., Int. Ed.*, 2000, **39**, 173; E. Yoshikawa, K. V. Radhakrishnan and Y. Yamamoto, *J. Am. Chem. Soc.*, 2000, **122**, 7280; T. Takahashi, R. Hara, Y. Nishihara and M. Kotor, *J. Am. Chem. Soc.*, 1996, **118**, 5154; T. Takahashi, Z. F. Xi, A. Yamazaki, Y. H. Liu, K. Nakajima and M. Kotor, *J. Am. Chem. Soc.*, 1998, **120**, 1672; R. C. Larock, M. J. Doty, Q. P. Tian and J. M. Zenner, *J. Org. Chem.*, 1997, **62**, 7536; N. Iwasawa, M. Shido, K. Maeyama and H. Kusama, *J. Am. Chem. Soc.*, 2000, **122**, 10226.
- 6 For transmetalation of acyclic organozirconium to Zn, see E. Negishi, D. E. Van Horn, T. Yoshida and C. L. Rand, *Organometallics*, 1983, **2**, 563.
- 7 G. Erker, *J. Organomet. Chem.*, 1977, **134**, 189; K. Kropp and G. Erker, *Organometallics*, 1982, **1**, 1246; S. L. Buchwald and B. T. Watson, *J. Am. Chem. Soc.*, 1986, **108**, 7411.