



IDENTIFICATION AND STRUCTURE-ACTIVITY RELATIONSHIP STUDIES OF OSTHOL, A CYTOTOXIC PRINCIPLE FROM *CNIDIUM MONNIERI*

Yukio Hitotsuyanagi, Hiroshi Kojima, Hiroshi Ikuta, Koichi Takeya and Hideji Itokawa*


Tokyo University of Pharmacy and Life Science, Horinouchi, Hachioji, Tokyo 192-03, Japan

Abstract: Osthol (**1**) was isolated from the fruit of *Cnidium monnieri* as a cytotoxic principle. Its structure-activity relationship study reveals that the 3,4-olefinic bond is essential for its cytotoxic activity, and the prenyl (C₅) unit attached at the 8 position enhances the cytotoxicity. Analogues **7** and **8** that have a longer alkoxy unit at the 7 position showed ten times higher cytotoxicity than **1**. Copyright © 1996 Elsevier Science Ltd

During the course of identifying the cytotoxic substances from higher plants,¹ we found that the methanol extract of the fruit of *Cnidium monnieri* (Umbelliferae) showed weak cytotoxicity (IC₅₀ = 24 µg/mL) against V-79 Chinese hamster lung cells, and bioassay-guided fractionation/isolation yielded osthol (**1**)² as the active substance.³ The IC₅₀ values of **1** against V-79, KB, P-388 and MM46 cells⁴ are 14, 12, 3.1 and 8.6 µg/mL, respectively. Although coumarins are a rather popular class of compounds from higher plants, and some of them have been reported to possess cytotoxic/antitumor activity,^{2,5} their structure-activity relationship (SAR) is still unclear. Since osthol (**1**) is available from this plant in quantity, chemical modification of this compound was made in order to obtain information about the SAR of **1** and to obtain more potent analogue which may qualify the potential of this lead as an antineoplastic agent.

The modification was made on the 3,4-olefinic bond, the substituent at position 7 and the prenyl group at position 8, which yielded analogues **2**—**16** as shown in Scheme 1.⁶ Catalytic hydrogenation of **1** over palladium on carbon gave the dihydro-(**2**) and tetrahydro-(**3**) derivatives in a ratio of 1:3, and the latter was re-dehydrogenated to **2** in 72% yield. The des-*O*-methylation of **2** and **3** using boron tribromide gave phenols **4** and **5**, respectively. The dehydroxylation of **4** occurred through the reduction of the intermediary triflate using 1,1-bis(diphenylphosphino)ferrocene, Pd(OAc)₂ and formic acid⁷ to afford compound **6**. The *O*-alkylation of **4**

Scheme 1



Reagents (a) H₂, Pd/C, CH₂Cl₂-EtOH, (b) Pd/C, mesitylene, reflux, (c) BBr₃, CH₂Cl₂, -78°C, (d) (CF₃SO₂)₂O, Et₃N, CH₂Cl₂, 0°C; 1,1-bis(diphenylphosphino)ferrocene, Pd(OAc)₂, HCO₂H, DMF, 60°C, (e) ROH, Ph₃P, diethyl azodicarboxylate, PhMe, room temp., (f) RCOCl, Et₃N, CH₂Cl₂, room temp., (g) Br₂, CH₂Cl₂, 0°C, (h) BF₃•OEt₂, H₂O-CH₂Cl₂, room temp., (i) *m*-CPBA, CH₂Cl₂, 0°C, (j) H₂SO₄, THF-H₂O, room temp.

Table 1. Cytotoxicity of coumarins against V-79 cells

compound	IC ₅₀ (μM)	compound	IC ₅₀ (μM)
osthol (1)	57	11	60
2	58	12	58
3	>100	13	60
4	40	14	>100
5	>100	15	>100
6	60	16	>100
7	5.8	coumarin	>100
8	5.6	7-hydroxycoumarin	58
9	16	7-methoxycoumarin	>100
10	17		

under Mitsunobu conditions⁸ with the corresponding alcohols produced ethers **7**–**10**, and the *O*-acylation with the corresponding acyl chlorides gave esters **11** and **12**. The bromination of **1** yielded dibromide **13**, and the acid catalyzed hydration of **1** gave tertiary alcohol **14**. The epoxidation of **1** using *m*-chloroperbenzoic acid (*m*-CPBA) yielded epoxide **15**, which was subjected to acid hydrolysis affording diol **16**.

The prepared compounds were evaluated using V-79 cells, and the results are summarized in Table 1. The values are given in molar concentration (μM) for comparison. Coumarin, 7-hydroxycoumarin and 7-methoxycoumarin were also evaluated for comparison. The unsaturation of the prenyl group is not essential to the activity (**1** vs. **2**). Since compounds **2**, **4** and **6** showed more potent activity than 7-methoxycoumarin, 7-hydroxycoumarin and coumarin, respectively, the C₅ unit attached at the position 8 proved to enhance the activity. Contrary to dibromide **13** retaining activity, the introduction of a hydrophilic moiety (**14**, **15** and **16**) into the prenyl group resulted in a loss in activity. With regard to the substituent at position 7, although the substitution of the methoxy group of **2** by a hydrogen (**6**) did not alter the activity, a hydroxy group (**4**) slightly enhanced the activity. The substitution of the methyl group of the 7-methoxy group by a longer alkyl chain (C₂–C₇) resulted in the enhancement of the activity (**7**–**10**), and the ethoxy (**7**) and the *n*-propoxy (**8**) analogues showed the most potent activity. The substitution by an acyloxy group did not alter the activity (**11** and **12**) in spite of the close similarity in length (**7** vs. **11**). Since the dihydrocoumarins **3** and **5** showed no activity, the 3,4-unsaturated bond is essential for such activity.

Since analogues **7** and **8** showed ten times more potent cytotoxicity than **1**, we currently intend to evaluate these analogues using other cell lines and prepare more potent analogues through these lines of modification.

Acknowledgement: This work was partly supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan.

References and notes

1. Itokawa, H.; Qiao, Z.-S.; Hirobe, C.; Takeya, K. *Chem. Pharm. Bull.* **1995**, *43*, 1171; Itokawa, H.; Nishimura, K.; Hitotsuyanagi, Y.; Takeya, K.; *Bioorg. Med. Chem. Lett.* **1995**, *5*, 821; Hirobe, C.; Palevitch, D.; Takeya, K.; Itokawa, H. *Natural Medicines* **1994**, *48*, 168.
2. The structure of **1** was confirmed by the comparison of its spectral data with the reported values. Gawron, A.; Giowwiak, K. *Planta Med.* **1987**, *48*, 526.
3. During the isolation process of **1**, we also isolated furocumarins xanthotoxin,^a bergapten,^{a,b} isopimpinellin,^{a,b} *O*-acetylcolumbianetin^c and edultin.^d Some of these compounds showed weak cytotoxicity against V-79 cells. a) Lee, K.-H.; Soine, T. O.; *J. Pharm. Sci.* **1969**, *58*, 681. b) Bergenthal, D.; Rózsa, Z.; Mester, I.; Reisch, J. *Arch. Pharm.* **1978**, *311*, 1026; Sin, H.-D.; Ding, J.-K., Lin, Z.-W.; Yi, Y.-F.; Fu, J. *Yaoxue Xuebao* **1982**, *17*, 835. c) Willette, R. E.; Soine, T. O. *J. Pharm. Sci.* **1964**, *53*, 275; Hata, K.; Kozawa, M.; Baba, K.; Yen, K.-Y. *Chem. Pharm. Bull.* **1971**, *19*, 640. d) Mitsuhashi, H.; Ito, T. *Chem. Pharm. Bull.* **1962**, *10*, 511; Nakazaki, M.; Yirose, Y.; Ikematsu, K. *Tetrahedron Lett.* **1966**, 4735; Savina, A. A.; Vandyshev, V. V.; Perelson, M. E.; Pimenov, M. G. *Khim. Prir. Soedin.* **1971**, 116.
4. KB: human epidermoid carcinoma of nasopharynx; P-388: mouse lymphocytic leukemia; MM46: mouse mammary carcinoma.
5. Rosskopf, F.; Kraus, J.; Franz, G. *Pharmazie* **1992**, *47*, 139; Gunatilaka, A. A. L.; Kingston, D. G. I.; Wijeratne, E. M. K.; Bandara, B. M. R.; Hofmann, G. A.; Johnson, R. K. *J. Nat. Prod.* **1994**, *57*, 518; Maucher, A.; von Angerer, E. *J. Cancer Res. Clin. Oncol.* **1994**, *120*, 502.
6. All new compounds gave satisfactory spectral (300 or 400 MHz ¹H NMR, IR, MS and HR-MS) and/or elemental analysis data.
7. Cacchi, S.; Ciattini, P. G.; Morera, E.; Ortari, G.; *Tetrahedron Lett.* **1986**, *27*, 5541.
8. Mitsunobu, O. *Synthesis* **1981**, 1.

(Received in Japan 13 May 1996; accepted 1 July 1996)