Bioorganic & Medicinal Chemistry Letters 20 (2010) 6840-6844

Contents lists available at ScienceDirect



Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl



Synthesis and pharmacological activity of 1,3,6-trisubstituted-4-oxo-1,4-dihydroquinoline-2-carboxylic acids as selective ET_A antagonists

Hardik J. Patel^{a,†}, Nicole Olgun^a, István Lengyel^b, Sandra Reznik^a, Ralph A. Stephani^{a,b,*}

^a Department of Pharmaceutical Sciences, St. John's University, 8000 Utopia Parkway, Jamaica, NY 11439, USA
^b Department of Chemistry, St. John's University, 8000 Utopia Parkway, Jamaica, NY 11439, USA

ARTICLE INFO

Article history: Received 16 July 2010 Accepted 16 August 2010

Keywords: Endothelin ETA antagonists ETAR selectivity versus ETBR 4-Oxo-1,4-dihydroquinoline-2-carboxylic acids

ABSTRACT

A series of 1,3,6-trisubsituted-4-oxo-1,4-dihyroquinoline-2-carboxylic acid analogs (**2a-m**) were designed and synthesized and their pharmacological activity determined, with the objective to better understand their SAR as potential ET_A selective inhibitors. Most of the compounds displayed significant ET_A antagonist activity having IC_{50} for inhibition of binding of the [¹²⁵I]ET-1 to ET_A receptor <10 nM, with good selectivity for ET_A antagonism over ET_B receptor. Based on the in vitro results, SAR of this series of compounds requires an alkoxy substituent at the 6-position to be a straight and saturated chain up to three carbons long, since substitution of unsaturated and branched alkyloxy groups results in decrease in ET_A antagonist activity. In this series, compound **2c** (6-*O*-*n*-propyl analog) was found to be most potent ($IC_{50} = 0.11 \text{ nM}$) with ET_B/ET_A selectivity of 8303.

© 2010 Elsevier Ltd. All rights reserved.

Endothelins are divided into three structurally similar subtypes: endothelin-1, 2, and 3 (ET-1, ET-2, and ET-3), which are members of 21 amino acid peptide families.¹ Among them, ET-1 is the major isoform involved in human cardiovascular and non-cardiovascular physiology and pathophysiology. ET-1 is a very potent vasoconstrictor, 10 times more potent than angiotensin II and 100 times more potent than epinephrine and also has inotropic, pro-inflammatory, and mitogenic properties. It also influences the salt-water homeostasis, and stimulates the renin–angiotensin–aldosterone and sympathetic nervous systems.^{2,3} Increased level of the plasma concentration of ET-1 has been implicated in several diseases such as hypertension, pulmonary hypertension, acute myocardial infarction, congestive heart failure, renal failure, atherosclerosis, pulmonary fibrosis, and cancer.⁴

ET-1 exerts its physiological action by binding to the specific receptors called endothelin receptor type A (ET_A) and B (ET_B), belonging to the heptahelical G-protein-coupled receptor superfamily.⁵ Endothelin receptors are widely expressed in most of the physiological systems, and interaction of ET_A expressed on VSMC with ET-1 results in vasoconstriction and cell proliferation.⁶ On the other hand, ET_B activation is mainly responsible for vasodilation depending on tissue location⁷ and clearance of endogenous endothelins.⁸

Endothelin receptor antagonists (ERAs), both selective ET_A antagonist and non-selective ET_A/ET_B antagonists, are novel thera-

peutics in clinical development for various diseases like pulmonary arterial hypertension, chronic heart failure, vascular remodeling (restenosis, atherosclerosis), renal failure, cancer, and lung fibrosis.⁹ Though theoretically it seems, selective ET_A receptor antagonists would be useful as therapeutic agents for the treatment of the aforementioned chronic pathological conditions. Advantages of a selective antagonist of the ET_A receptor¹⁰ are:

- Vasoconstriction in the human cardiovascular system by ET-1 appears to be predominantly mediated by the ET_A receptor.
- The mitogenic and pro-inflammatory effects induced by ET-1 are mediated by ET_A receptor stimulation.
- Blockade of the ET_B receptor would prevent the putative beneficial effects like vasodilation of ET-1-induced NO and PGI₂ release from the endothelium.
- Since the ET_B receptor in the lung is involved in clearance of ET-1, ET_B-antagonism will reduce the rate of clearance of exogenous ET-1 from the circulation.

Most non-peptidic endothelin receptor antagonists have the following common structural features: a carboxylic acid group or an acidic sulfonamide moiety between two aromatic or hetero-aromatic ring systems (Fig. 1).¹⁰ These pharmacophore groups are positioned so that they mimic amino acids of the ET-1 C-terminal, which is in a helical form. Different central core units like indane (enrasentan),¹¹ chromene,¹² cyclopentenopyridine (L-753037),^{13,14} and pyrrolidine (atrasentan)¹⁵ have been incorporated into endothelin antagonists currently in the clinical trials (Fig. 2).

^{*} Corresponding author. Tel.: +1 718 990 5215; fax: +1 718 990 1876.

E-mail address: stephanr@stjohns.edu (R.A. Stephani).

[†] Present address: Memorial Sloan-Kettering Cancer Center, New York, NY, USA.

⁰⁹⁶⁰⁻⁸⁹⁴X/\$ - see front matter \odot 2010 Elsevier Ltd. All rights reserved. doi:10.1016/j.bmcl.2010.08.074



Figure 1. General structure of endothelin receptor antagonists.¹⁰



Figure 2. Examples of endothelin receptor antagonists in clinical trials.

Earlier, when the indane ring of SB-209670 was superimposed on the quinolone ring, a good overlap was obtained with 1,3-disubstituted-2-carboxylic quinolones (1). Preliminary SAR resulted in compound 1 having potent ET_A antagonist activity and was 800 times more selective for ET_A .¹⁶ Also, it had been reported that introduction of an alkylamino or alkoxy was beneficial to potency and selectivity in analogs belonging to indane,¹¹ chromene,¹² and cyclopenteno[1,2*b*]pyridine^{13,14} carboxylic acid series. Therefore it was hypothesized that introducing an alkoxy substituent at the 6th position of the A ring of quinolone would lead to a new series of compounds with increased potency and selectivity towards ET_A receptor (Fig. 3). This hypothesis was validated by generating the CoMFA model of endothelin receptor antagonists having IC₅₀ values between 0.19 and 780 nM, which showed steric groups favorable at the 6 position of the quinolone ring.

In the synthesis of the target compounds 1,3,6-trisubstituted-4oxo-1,4-dihydro-quinoline-2-carboxylic acids (**2a-m**), compound **15** is the key intermediate. Alkylation of **15** with various alkyl halides led to analogs with different substituents at the 6-position of



Figure 3. Hypothesis for the synthesis of 1,3,6-trisubstituted-4-oxo-1,4-dihydroquinoline-2-carboxylic acid as selective ET_A antagonists.

the quinolone. As shown in Scheme 1, nitration of 3-hydroxyacetophenone with nitric acid in the presence of acetic acid resulted in the desired regioisomer 3-hydroxy-6-nitroacetophenone, **4**, which had the reported melting point.^{17,18} Compound **4** was condensed with 3-carboxybenzaldehyde to yield *E*-configuration chalcone **5**. Carboxylic acid **5** was converted into ester **6** by reaction with anhydrous ethanol saturated with HCl gas followed by MOM protection of the hydroxyl group resulting in compound **7**. Hydrogenation at 45 psi with PtO₂ catalyst for 45 minutes reduced both the nitro group and the double bond of chalcone **7** to give amine **8**.

For the synthesis of intermediate 6-ethylpiperonal, **11**, Scheme 2 was followed. Wolff–Kishner reduction of 3,4-methylenedioxyace-tophenone (**9**) resulted in crude product, which was purified by vacuum-distillation at 130 °C and 27 mm/Hg to yield 63.4% of pure **10**. Formylation of 3,4-methylenedioxy ethylbenzene (**10**) with α,α -dichloromethyl methyl ether as precursor of the formyl group, via alkylation catalyzed by titanium tetrachloride, resulted in 6-ethylpiperonal (**11**) in excellent yield.¹⁹

Reductive amination of 6-ethylpiperonal, **11** with aromatic amine **8** was then carried out by using hydride reducing agents such as sodium cyanoborohydride (NaBH₃CN),²⁰ or titanium (IV) isopropoxide (Ti(OiPr)₄)/sodium cyanoborohydride (NaBH₃CN),²¹ or sodium triacetoxyborohydride (NaBH(OAc)₃).²² Among these three methods, reductive amination with NaBH(OAc)₃ gave the best yield of product **12**. N-acylation of **12** with ethyl oxalyl chloride gave an excellent yield of compound **13**, which on cyclization by the hindered base 1,8-diazabicyclo-[5.4.0]undec-7-ene (DBU) in ethanol gave MOM-protected-4-quinolone diester **14**. Deprotection of MOM-group under acidic condition resulted in key intermediate **15** (Scheme 3).

Alkylation of **15** with various alkyl halides in presence of sodium hydride and potassium carbonate led to derivatives with different substituents at the 6-position of the quinolone **16a–k** (Scheme 4). The synthesis of the target compounds (**2a–m**) was accomplished by potassium hydroxide treatment of the 4-oxoquinoline diesters in moderate yields. The general synthetic scheme along with the physical properties is described in Supplementary data.

All compounds listed in Table 1 were evaluated for their ability to inhibit endothelin-A (ET_A) and endothelin-B (ET_B) receptors in vitro. The results are summarized in Table 1. For each compound, eight concentrations were tested in the presence of fixed concentration of radio-labeled ET-1([¹²⁵I] ET-1) on green monkey renal tubular (Vero) cells expressing ET_A receptors²³ and membrane expressing ET_B receptor. Besides thirteen proposed compounds, BQ-123 (selective ET_A antagonist) was taken as a positive control to validate the method. The IC₅₀ values, which are the concentrations that reduce the [¹²⁵I] ET-1 binding to the



Scheme 1. Preparation of intermediate **8**. Reagents and conditions: (a) HNO₃, glacial CH₃COOH, 70 ± 2 °C; (b) 3-carboxybenzaldehyde, 10 N NaOH (3 equiv), CH₃OH, reflux; (c) dry HCl gas, anhydrous C₂H₅OH; (d) MOMCl, NaH, dry K₂CO₃, dry DMF; (e) 45 psi H₂ gas, PtO₂, anhydrous CH₃OH.



Scheme 2. Synthesis of 6-ethylpiperonal, 11. Reagents and conditions (a) (i) $NH_2NH_2 \cdot H_2O$, ethanol, reflux, (ii) KOH, 160 °C; (b) TiCl₄, α, α -dichloromethyl methyl ether, dichloroethane, 0 °C to rt.



Scheme 3. Preparation of intermediate 15. Reagents: (a) NaBH(OAc)₃, DCE; (b) ethyl oxalyl chloride, NEt₃, dry DMF; (c) DBU, ethanol; (d) dry HCl gas, anhydrous C₂H₅OH.

receptors by 50%, were determined for each drug and are listed in Table 1.

On the basis of this study, the following structure–activity relationships are proposed for the 1,3,6-trisubstituted-4-oxo-1,4-dihydro-quinoline-2-carboxylic acid analogs, **2a–m** as an ET_A antagonists. Among the compounds tested, 6-0-ethyl (**2b**), 6-0-*n*propyl (**2c**), 6-0-*n*-butyl (**2g**), 6-0-*i*-butyl (**2h**), 6-0-sec-butyl (**2i**), 6-0-propan-3-ol (**2l**), 4-oxo-1,4-dihydroquinoline diacids displayed significant ET_A antagonist activity having IC₅₀ for inhibition of binding of the [¹²⁵1]ET-1 to ET_A receptor <10 nM. As shown in Table 1, when the 6-OH group of **2a** is replaced by a 6-alkoxy group of two carbons (**2b** IC_{50} : 6.6 nM), there is a sixfold increase in ET_A antagonist activity and a slight increase in selectivity for ET_A over ET_B , i.e., 473 for **2b** versus 399 for **2a**. Increase in chain length by one more carbon to 3 carbons (**2c**) resulted in 60-fold increase in ET_A (IC_{50} : 0.11 nM) and threefold increase in ET_B (IC_{50} : 913 nM) antagonist activity compared to **2b**. Although there is increase in binding affinity for both the receptors, a *n*-propoxy group at 6-position of quinoline is more beneficial for ET_A over ET_B . There is 10-fold decrease in ET_A antagonist activity compared to **2c** when the side chain length is increased by one carbon to butyl (**2g**;



Scheme 4. Synthesis of 6-O-substituted-4-oxo-quinoline-dicarboxylic acids (2a-m). Reagents and conditions: (a) alkyl halides, NaH, dry K₂CO₃, dry DMF; (b) 6 N KOH (6 equiv), C₂H₅OH, reflux.

Table 1

In vitro inhibitory binding assay (IC₅₀) of 1,3,6-trisubstituted-4-oxo-1,4-dihydroquinoline-2-carboxylic acids, **2a-m**



No.	R	Mp (°C)	$ET_A IC_{50}^a (nM)$	$ET_B IC_{50}^b (nM)$	ET_B/ET_A
2a	Н	198-199	38.9 ± 2.6	15561.5 ± 162.7	399
2b	C ₂ H ₅	213-215	6.6 ± 1.1	3119 ± 54.3	473
2c	$n-C_3H_7$	162–163	0.11 ± 0.02	913.4 ± 27.2	8304
2d	i-C ₃ H ₇	199-200	13.5 ± 1.5	677 ± 9.6	50
2e	$CH_2 = CH - CH_2$	165-166	38.1 ± 3.6	1131 ± 24.8	30
2f	CH ₃ -O-CH ₂	176–177	24.4 ± 2.9	1030 ± 17.3	43
2g	n-C ₄ H ₉	161-162	1.44 ± 0.6	548.2 ± 14.5	381
2h	i-C ₄ H ₉	229-231	6.3 ± 1.5	1983 ± 35.7	315
2i	sec-C ₄ H ₉	185-186	9.9 ± 1.7	_	-
2j	cyclo-C ₃ H ₅ -CH ₂	173–174	115.9 ± 11.4	_	-
2k	<i>n</i> -C ₅ H ₁₁	153-155	610 ± 54	_	-
21	HO-CH ₂ CH ₂ CH ₂	192–194	7.4 ± 0.9	4387 ± 63.7	593
2m	HOOC-CH ₂	177-180	284.5 ± 3.5	1453 ± 20.7	5
	BQ123	-	24.8 ± 1.3 ^c (25)	-	-

^a Monkey renal cells CCL-81 grown as monolayer.²³ n = 3.

^b Endothelin-B receptor precursor (human) membranes. n = 2.

^c Literature IC₅₀ value, *ibid*.

IC₅₀: 1.6 nM), but still shows better activity than **2b**. Further increase in length to 5 carbons (**2k**; IC₅₀ 610 nM) resulted in 6000-fold loss in activity compared to 6-*O*-*n*-propyl analog (**2c**), also seen in the cyclopenteno[1,2-*b*]pyridine series as ET_A receptor antagonist.¹⁴

Replacement of the saturated 6-O-*n*-propyl group with the unsaturated 6-O-allyl group resulted in **2e** analog (IC₅₀ = 38.11 nM) 380 times less active than the *n*-propyl analog **2c**. This suggests that a saturated alkyl chain is required at the 6-position of the 4-oxo-1,4-dihydroquinoline diacids for better ET_A antagonist activity. Also, when the second atom of the *n*-propyl chain in **2c** is replaced by isosteric O (**2f**; IC₅₀: 24.4 nM), there is 240-fold decrease in activity compared to **2c**.

Substitutions at the C1 of the 6-O-alkyl chain are not well tolerated. There is a twofold decrease in ET_A antagonist activity compared to **2b** when the methylene group of the ethyl chain is substituted, **2d** (*i*-propyl, $IC_{50} = 13.5$ nM). Similarly, substitution of the methylene group at C1 of the 6-*O*-*n*-propyl chain resulted in compound **2i** (6-*O*-sec-butyl) which is 90-fold less active, than **2c**, in inhibition of binding of the [^{125}I]ET-1 to ET_A receptor ($IC_{50} = 9.9$ nM). Likewise, branching at C2 of the alkyl chain is not tolerated, as substitution of a methyl group at C2 of the 6-*O*-*n*-propyl chain resulted in analog **2h** (6-*O*-*i*-butyl) which is 60 times less active as ET_A receptor antagonist ($IC_{50} = 6.6$ nM) than **2c**. Also, compound **2j** (6-*O*-cyclopropylmethyl) which is a cyclic analog of **2h**, shows 1000 times less ET_A receptor antagonist activity than **2c**. These results suggest that straight chain saturated alkoxy groups are required at the 6-position of the 4-oxo-1,4-dihydroquinoline diacids for good ET_A receptor antagonism. Introduction of a hydroxyl group (**2l**; IC₅₀: 7.4 nM) at C3 of the 6-*O*-*n*-propyl side chain resulted in 70-fold reduction in ET_A receptor antagonist activity compared with **2c**. Also, substitution of a carboxylic acid group (**2m**; IC₅₀: 284.46 nM) for a methyl group in **2b** causes a 43-fold decrease in binding affinity for ET_A receptor compared with **2b**. These results are consistent with the previously reported SAR of cyclopenteno[1,2-*b*]pyridine analogs as ET_A receptor antagonists.¹⁴

Compound **2a** was found to reduce lung inflammation induced by lipopolysaccharide (LPS),²⁴ cigarette smoke, and ET-1²⁵ in a hamster animal model by reducing the recruitment of broncoalveolar lavage fluid (BALF) leukocytes (neutrophils and macrophages). Therefore, reducing ET-1 activity with selective ET_A receptor antagonists could potentially decrease neutrophil derived enzymes and oxidants in the lung, thereby decreasing or inhibiting the progression of lung injury. Compound **2a** was also effective in controlling preterm delivery induced by lipopolysaccharide (LPS) in a pregnant mouse animal model.²⁶ This result, in accordance with previous literature, suggests the role of ET-1 in both normal parturition and premature birth.

In summary, a new series of 1,3,6-trisubstituted-4-oxo-1,4dihydroquinoline-2-carboxylic acid analogs were synthesized and evaluated for their ET_A antagonist activity and selectivity for ET_A receptor over ET_B receptor. Among the compounds tested, **2b**, **2c**, 2g, 2h, 2i, and 2l 4-oxo-1,4-dihydroquinoline diacids displayed significant ET_A antagonist activity with IC₅₀ values of inhibition of binding of the [¹²⁵I]ET-1 to ET_A receptor <10 nM. These compounds also showed good selectivity for ET_A antagonism over ET_B receptor. Initial SAR at C6 of the quinolone ring indicates straight, unbranched, and saturated alkyl chain up to three carbons long, with hetero atom spacer required for good ET_A antagonist activity and selectivity. Also, polar hydrophilic groups at the 6-position are unfavorable for ET_A antagonist activity, as seen with a decrease in activity when substituted by 6-O-propan-3-ol (21) and 6-O-carboxymethyl (2m) of the quinoline ring. The 6-O-n-propyl-4-oxo-1,4-dihydroquinoline diacids analog, 2c was found to be most potent (IC₅₀: 0.11 nM) with ET_B/ET_A selectivity of 8303. In vivo, compound **2a** (6-OH), a prototype of this 4-oxo-1.4-dihydroquinolone diacid series was shown to reduce LPS-induced lung inflammation and preterm labor in two different animal models.

Acknowledgments

The authors thank Maulik Patel for helping to construct a CoM-FA model. Support for this research was provided by Saint John's University.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bmcl.2010.08.074.

References and notes

- 1. Inoue, A.; Yanagisawa, M.; Kimura, S.; Kasuya, Y.; Miyauchi, T.; Goto, K., et al *Proc. Natl. Acad. Sci.* **1989**, *86*, 2863.
- Haynes, W. G.; Hand, M. F.; Johnstone, H. A.; Padfield, P. L.; Webb, D. J. J. Clin. Invest. 1994, 94, 1359.
- Nishimura, M.; Takahashi, H.; Matsusawa, M.; Ikegaki, I.; Sakamoto, M.; Nakanishi, T., et al J. Hypertens. 1991, 9, 71.
- (a) Motte, S.; McEntee, K.; Naeije, R. Pharmacol. Ther. 2006, 110, 386; (b) Grimshaw, M. J. Endocr. Related Cancer 2007, 14, 233; (c) Bagnato, A.; Rosanò, L. Int. J. Biochem. Cell Biol. 2008, 40, 1443; (d) Kohan, D. E. Curr. Hypertens. Rep. 2008, 10, 65; (e) Fagan, K.; McMurtry, I.; Rodman, D. Respir. Res. 2001, 2, 90.
- 5. Epstein, F. H. N. Engl. J. Med. 1995, 333, 356.
- 6. Masaki, T. Trends Pharmacol. Sci. 2004, 25, 219.
- Moreland, S.; Mc Mullen, D. M.; Delaney, C. L.; Lee, V. G.; Hunt, J. T. Biochem. Biophys. Res. Commun. 1992, 184, 100.
- 8. Dupuis, J.; Stewart, D. J.; Cernacek, P.; Gosselin, G. Circulation 1996, 94, 1578.
- 9. Shah, R. Eur. J. Intern. Med. 2007, 18, 272.
- 10. Boss, C.; Bolli, M.; Weller, T. Curr. Med. Chem. 2002, 9, 349.
- Elliot, J. D.; Lago, M. A.; Cousins, R. D.; Gao, A.; Lober, J. D.; Erhard, K. F.; Nambi, P.; Elshourbagy, N. A.; Kumar, C.; Lee, J. A.; Bean, J. W.; DeBrosse, C. W.; Eggleston, D. S.; Brooke, D. P.; Feuerstein, G.; Ruffulo, R. R.; Weinstock, J.; Gleason, J. G.; Poishoff, C. E.; Ohlstein, E. H. J. Med. Chem. **1994**, 37, 1553.
- Ishizuka, N.; Matsumura, K.; Sakai, K.; Fujimoto, M.; Mihara, S.; Yamamori, T. J. Med. Chem. 2002, 45, 2041.
- Niiyama, K.; Mase, T.; Takahashi, H.; Naya, A.; Katsuki, K.; Nagase, T.; Ito, S.; Hayama, T.; Hisaka, A.; Ozaki, S.; Ihara, M.; Yano, M.; Fukuroda, T.; Noguchi, K.; Nishikibe, M.; Ishikawa, K. *Bioorg. Med. Chem.* **2002**, *10*, 2461.
- Niiyama, K.; Takahashi, H.; Nagase, T.; Kojima, H.; Amano, Y.; Katsuki, K.; Yamakawa, T.; Ozaki, S.; Ihara, M.; Yano, M.; Fukuroda, T.; Nishikibe, M.; Ishikawa, K. Bioorg. Med. Chem. Lett. 2002, 12, 3041.
- Winn, M.; von Geldern, T. W.; Opgenorth, T. J.; Jae, H.-S.; Tasker, A. S.; Boyd, S. A.; Kester, J. A.; Mantei, R. A.; Bal, R.; Sorenson, B. K.; Wu-Wong, J. R.; Chiou, W. J.; Dixon, D. B.; Novosad, E. I.; Hernandez, L.; Marsh, K. C. *J. Med. Chem.* **1996**, *39*, 1039.
- Haesslein, J.-L.; Baholet, I.; Fortin, M.; Iltis, A.; Khider, J.; Periers, A. M.; Pierre, C.; Vevert, J.-P. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 1487.
- 17. Klonke, P.; Gibson, H. Chem. Ber. 1961, 94, 26.
- 18. Butenandt, A.; Hallmann, G.; Beckmann, R. Chem. Ber. 1957, 90, 1120.
- 19. Sonnet, P. E. J. Med. Chem. 1972, 15, 97.
- 20. Borch, R. F.; Bernstein, M. D.; Durst, H. D. J. Am. Chem. Soc. 1971, 93, 2897.
- Mattson, R. J.; Pham, K. M.; Leuck, D. J.; Cowen, K. A. J. Org. Chem. 1990, 55, 2552.
- 22. Abdel-Magid, A. F.; Mehrman, S. J. Org. Process Res. Dev. 2006, 10, 971.
- 23. Funk, O. F.; Kettmann, V.; Drimal, J.; Langer, T. J. Med. Chem. 2004, 47, 2750.
- Bhavsar, T. M.; Cerreta, J. M.; Liu, M.; Reznik, S.; Cantor, J. O. *Exp. Lung Res.* 2008, 34, 141.
- 25. Bhavsar, T. M.; Liu, X. J.; Patel, H.; Stephani, R.; Cantor, J. O. Int. J. COPD **2008**, 3, 1.
- Olgun, N.; Patel, H. J.; Stephani, R.; Wang, W.; Yen, H.; Reznik, S. E. Can. J. Physiol. Pharmacol. 2008, 86, 571.