



Cite this: *Org. Biomol. Chem.*, 2018, **16**, 9133

Received 27th September 2018,
Accepted 6th November 2018

DOI: 10.1039/c8ob02413h

rsc.li/obc

Iron(III) chloride-catalyzed activation of glycosyl chlorides†

Scott A. Geringer and Alexei V. Demchenko  *

Glycosyl chlorides have historically been activated using harsh conditions and/or toxic stoichiometric promoters. More recently, the Ye and the Jacobsen groups showed that glycosyl chlorides can be activated under organocatalytic conditions. However, those reactions are slow, require specialized catalysts and high temperatures, but still provide only moderate yields. Presented herein is a simple method for the activation of glycosyl chlorides using abundant and inexpensive ferric chloride in catalytic amounts. Our preliminary results indicate that both benzylated and benzoyleated glycosyl chlorides can be activated with 20 mol% of FeCl₃.

Introduced by Michael in 1879¹ and subsequently studied by many, glycosyl chlorides have been very influential building blocks that helped to establish basic principles of carbohydrate chemistry.^{2,3} Once prominent glycosyl donors, in recent years glycosyl chlorides have been outshadowed by other, more powerful glycosyl donors,^{4–11} and for a reason. Traditionally, the activation of glycosyl chlorides demanded stoichiometric and often toxic reagents, such as silver(I)^{2,12,13} or mercury(II) salts.¹⁴ This, along with a fairly high propensity to hydrolysis, hampered the application of glycosyl chlorides in recent years. Glycosyl chlorides, however, have many positive traits. They can be obtained using a variety of substrates and methods,^{15–25} many chlorides are stable, and recent studies by Ye *et al.*²⁶ and Jacobsen *et al.*²⁷ have demonstrated that these compounds can be activated without toxic promoters under organocatalytic conditions using urea- or thiourea-based catalysts. Good stereoselectivity was obtained using various additives²⁶ or with complex chiral catalytic constructs,²⁷ but these reactions are slow (24–48 h), require high temperatures and provide practical yields only with highly reactive (alkylated) chlorides. In an active pursuit of catalytic activation methods

for glycosylation,^{28,29} we observed that glycosidation of chlorides can be achieved in the presence of catalytic amounts of iron(III) chloride (FeCl₃ aka ferric chloride). This discovery is at the basis of this communication.

FeCl₃ is naturally abundant, inexpensive and relatively benign.³⁰ Ferric chloride has been employed in the introduction of protecting groups in carbohydrates.^{31,32} The application of FeCl₃ in O-glycosylation has also emerged, most prominently for the activation of glycosyl donors bearing the anomeric acetate.^{33–42} Other applications for the activation of aryl glycoside,⁴³ pivaloate,⁴⁴ bromide,⁴⁵ imidate,⁴⁶ or hemiacetal donors (as a co-catalyst)⁴⁷ have also been explored. Using this prior knowledge, we theorized that glycosyl chlorides may also offer a promising new substrate for the catalytic activation with FeCl₃. To test this hypothesis, we chose known per-benzylated glucosyl chloride donor **1**²³ to couple with the standard glycosyl acceptor **2**.⁴⁸ The glycosylation was set-up in the presence of molecular sieves (4 Å) in dichloromethane. For this preliminary study we chose excess of donor **1** (2.0 equiv.) similarly to that used by Ye *et al.*²⁶ and Jacobsen *et al.*²⁷ After a brief preliminary experimentation, we established that 20 mol% of FeCl₃ provides the most favorable balance between yields and the reaction time. Thus, the coupling of donor **1** with acceptor **2**⁴⁸ provided disaccharide **3** in 67% yield in only 2 h (Table 1, entry 1). Also, glycosidations of chloride **1** with secondary acceptors **4**, **6**, and **8**⁴⁸ were conducted under essentially the same reaction conditions. These reactions were slower (3–16 h), but the respective disaccharides **5**, **7** and **9** have successfully been obtained in 47–80% yields (entries 2–4). This preliminary set of experiments has demonstrated both the advantages and limitations of this approach. The main advantage of this approach is the availability and low cost of the catalytic activator. Also the reaction times are notably shorter than those reported for the organocatalytic reactions and even for the traditional heavy metal-based stoichiometric activators. Somewhat average yields for the formation of all products, perhaps except **9**, still on a par with traditional approaches and the results reported by Ye *et al.*²⁶ and

Department of Chemistry and Biochemistry, University of Missouri – St Louis, One University Boulevard, St Louis, Missouri 63121, USA.




E-mail: demchenkoa@umsl.edu

† Electronic supplementary information (ESI) available: Experimental details and characterization data for all new compounds. See DOI: 10.1039/c8ob02413h

Table 1 Iron(III) chloride-catalyzed glycosylations

donor (1 equiv) Glc, Gal, Man Bx = Bn, Bz acceptor (0.5 equiv)						
Entry	Donor	Acceptor	Time	Product	Yield	α/β ratio
1			2 h		67%	1.1/1
2	1		16 h		47%	1.2/1
3	1		3 h		60%	1.5/1
4	1		16 h		80%	1.0/1
5		2	0.5 h		88%	1/1.4
6	10	4	0.5 h		57%	1.6/1
7	10	6	0.5 h		80%	1.3/1
8	10	8	16 h		90%	1/2.7
9		2	2 h		80%	4.5/1
10	15	4	16 h		66%	α -Only
11	15	6	16 h		56%	α -Only
12	15	8	16 h		95%	2.6/1
13		2	16 h		98%	β -Only
14	20	4	16 h		80%	β -Only

Table 1 (Contd.)


						
Entry	Donor	Acceptor	Time	Product	Yield	α/β ratio
15	20	6	16 h		52%	β -Only
16	20	8	16 h		73%	β -Only

Jacobsen *et al.*,²⁷ are mainly attributed to a substantial formation of a side product of 1,6-anhydro-2,3,4-tri-*O*-benzyl- β -D-glucopyranose. While somewhat unexpected in this particular setting, the formation of 1,6-anhydro sugars in the presence of FeCl_3 has been reported.⁴⁹ As evident from Table 1, all four disaccharides have been produced with poor selectivity ($\alpha/\beta = 1\text{--}1.5/1$); our method, however, does not employ stereo-directing functionalities, additives,²⁶ or complex chiral catalytic constructs²⁷ at this stage.

Following the general success of glucosyl chloride donor **1** we investigated galactosyl chloride **10**²³ that provided even faster reaction times (entries 5–8), probably due to the generally higher reactivity of the galactosyl donors *versus* similarly equipped glucose counterparts, and a noticeable increase in yields. The latter could be attributed to the entire absence of the 1,6-anhydro side-product that hampered the yields with donor **1**.

Thus, primary acceptor **2** led to the formation of disaccharide **11** in a respectable yield of 88% (entry 5). For comparison, glucosyl chloride donor **1** produced the 1 $\rightarrow$ 6 linked disaccharide **3** in 67% (see entry 1). A similar enhancement in yields (up to 90%) and decrease in the reaction time have been observed for the secondary acceptors to produce the respective disaccharides **12–14** (entries 6–8). Expectedly, mannosyl donor **15**¹³ showed lower reactivity than its glucosyl and galactosyl counterparts. This was reflected by the increase in reaction times; nevertheless, we obtained respectable yields (up to 95%) for the synthesis of disaccharides **16–19** (entries 9–12). No formation of the 1,6-anhydro side product was detected in this case either. We believe this reaction follows a traditional Lewis acid-catalyzed mechanistic pathway depicted in Scheme 1. Presumably, this reaction follows the traditional unimolecular $\text{S}_{\text{N}}1$ mechanism according to which the catalyst-mediated leaving group departure results in the formation of the oxacarbenium ion. The latter exists in a flattened half-chair conformation that explains poor stereoselectivity observed.

Having demonstrated that FeCl_3 -catalyzed reactions work reasonably well with per-benzylated sugars, we wanted to investigate whether electronically deactivated benzoylated chloride **20** could be activated using our method. As expected,



Scheme 1 Proposed mechanism of the activation of glycosyl chlorides with ferric chloride.

when donor **20** was glycosidated with acceptor **2** a slower reaction time 16 h (entry 13, Table 1) was recorded in comparison to that with the benzylated glucosyl donor **1** (2 h, see entry 1). Nevertheless, the reaction still proceeded to completion and provided disaccharide **21** in an impeccable yield of 98% and no indication for the side product formation. The glycosidation of donor **20** also proceeded well with the secondary acceptors **4**, **6**, and **8** providing the corresponding disaccharides **22–24** in respectable yields of 52–80% and complete β -selectivity due to the neighboring group participation. It is noteworthy that neither Ye's nor Jacobsen's conditions were able to activate these deactivated benzoylated chlorides.

In conclusion, we have shown that a variety of glycosyl chlorides can be activated with catalytic iron(III) chloride. This method allows for a cheap and relatively benign activation of glycosyl chlorides compared to previous methods using harsher and less environmentally friendly conditions. While the yield of glycosylation reactions are still far from being ideal, a majority of results obtained herein are on a par with recently developed organocatalytic reactions reported by Ye *et al.*²⁶ and Jacobsen *et al.*²⁷ The stereoselectivity obtained in reactions with benzylated chlorides is unimpressive, which is not a surprise because we do not currently employ any directing auxiliaries, catalysts, or additives as in other similar studies. However, our study employs a very inexpensive activa-

tor, and this method can serve as a basis for refining stereoselectivity in the future. One of the possible directions for this to explore the known effect of stoichiometric FeCl_3 that is capable of producing the α -product preferentially, presumably due to post-glycosylational anomerization reaction.⁴⁰

Of particular significance is that electronically deactivated, benzoylated chlorides can also be activated using our reaction conditions, whereas other catalytic systems fail to activate those unreactive substrates. The investigation of the scope and limitations of this method, including screening other Lewis acids, are currently underway in our laboratory and will be reported in due course. Our preliminary attempt to broaden the scope of this reaction by investigating SnCl_4 , $\text{BF}_3\text{-OEt}_2$, and $\text{Fe}(\text{OTf})_3$ indicated similar reaction yields and reaction times to those reported herein.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the National Institute of General Medical Sciences (GM120673) and the National Science Foundation (CHE-1800350).

References

- 1 A. Michael, *Am. Chem. J.*, 1879, **1**, 305.
- 2 W. Koenigs and E. Knorr, *Ber. Dtsch. Chem. Ges.*, 1901, **34**, 957.
- 3 S. S. Kulkarni and J. Gervay-Hague, in *Handbook of Chemical Glycosylation*, ed. A. V. Demchenko, Wiley-VCH, Weinheim, Germany, 2008, p. 59.
- 4 K. Toshima and K. Tatsuta, *Chem. Rev.*, 1993, **93**, 1503.
- 5 R. R. Schmidt and W. Kinzy, *Adv. Carbohydr. Chem. Biochem.*, 1994, **50**, 21.
- 6 D. M. Whitfield and S. P. Douglas, *Glycoconjugate J.*, 1996, **13**, 5.
- 7 Z. Gyorgydeak and I. E. Pelyvas, in *Glycoscience: Chemistry and Chemical Biology*, ed. B. Fraser-Reid, K. Tatsuta and J. Thiem, Springer, Berlin - Heidelberg - New York, 2001, vol. 1, pp. 691.
- 8 *Handbook of chemical glycosylation: advances in stereoselectivity and therapeutic relevance*, ed. A. V. Demchenko, Wiley-VCH, Weinheim, Germany, 2008.
- 9 S. S. Nigudkar and A. V. Demchenko, *Chem. Sci.*, 2015, **6**, 2687.
- 10 R. Das and B. Mukhopadhyay, *ChemistryOpen*, 2016, **5**, 401.
- 11 *Selective Glycosylations: Synthetic Methods and Catalysts*, ed. C. S. Bennett, Wiley-VCH, Weinheim, 2017.
- 12 K. Igarashi, T. Honma and J. Irisawa, *Carbohydr. Res.*, 1970, **15**, 329.
- 13 K. Matsuoka, M. Terabatake, A. Umino, Y. Esumi, K. Hatano, D. Terunuma and H. Kuzuhara, *Biomacromolecules*, 2006, **7**, 2274.
- 14 B. Helferich and K. F. Wedemeyer, *Ann.*, 1949, **563**, 139.
- 15 V. D. Grob, T. G. Squires and J. R. Vercellotti, *Carbohydr. Res.*, 1969, **10**, 595.
- 16 K. Jansson, G. Noori and G. Magnusson, *J. Org. Chem.*, 1990, **55**, 3181.
- 17 G. J. F. Chittenden, *Carbohydr. Res.*, 1992, **242**, 297.
- 18 S. C. Hung and C. H. Wong, *Tetrahedron Lett.*, 1996, **37**, 4903.
- 19 Z. Zhang and G. Magnusson, *Carbohydr. Res.*, 1996, **925**, 41.
- 20 S. Sugiyama and J. M. Diakur, *Org. Lett.*, 2000, **2**, 2713.
- 21 F. M. Ibatulin and S. I. Selivanov, *Tetrahedron Lett.*, 2002, **43**, 9577.
- 22 L. Encinas and J. L. Chiara, *J. Comb. Chem.*, 2008, **10**, 361.
- 23 A. M. Gómez, A. Pedregosa, M. Casillas, C. Uriel and J. C. López, *Eur. J. Org. Chem.*, 2009, 3579.
- 24 T. M. Beale, P. J. Moon and M. S. Taylor, *Org. Lett.*, 2014, **16**, 3604.
- 25 M. B. Tatina, D. T. Khong and Z. M. A. Judeh, *Eur. J. Org. Chem.*, 2018, 2208.
- 26 L. Sun, X. Wu, D. C. Xiong and X. S. Ye, *Angew. Chem., Int. Ed.*, 2016, **55**, 8041.
- 27 Y. Park, K. C. Harper, N. Kuhl, E. E. Kwan, R. Y. Liu and E. N. Jacobsen, *Science*, 2017, **355**, 162.
- 28 Y. Singh, T. Wang, S. A. Geringer, K. J. Stine and A. V. Demchenko, *J. Org. Chem.*, 2018, **83**, 374.
- 29 S. S. Nigudkar, K. J. Stine and A. V. Demchenko, *J. Am. Chem. Soc.*, 2014, **136**, 921.
- 30 I. Bauer and H. J. Knolker, *Chem. Rev.*, 2015, **115**, 3170.
- 31 T. Y. Huang, M. M. Zulueta and S. C. Hung, *Org. Biomol. Chem.*, 2014, **12**, 376.
- 32 A. Gouasmat, A. Lemétais, J. Solles, Y. Bourdreux and J.-M. Beau, *Eur. J. Org. Chem.*, 2017, 3355.
- 33 M. Kiso and L. Anderson, *Carbohydr. Res.*, 1979, **72**, C12.
- 34 M. Kiso and L. Anderson, *Carbohydr. Res.*, 1979, **72**, C15.
- 35 M. Kiso, H. Nishiguchi and A. Hasegawa, *Carbohydr. Res.*, 1980, **81**, C13.
- 36 F. Dasgupta and P. J. Garegg, *Acta Chem. Scand.*, 1989, **43**, 471.
- 37 L. M. Lerner, *Carbohydr. Res.*, 1990, **207**, 138.
- 38 S. K. Chatterjee and P. Nuhn, *Chem. Commun.*, 1998, 1729.
- 39 J. Seibel, L. Hillringhaus and R. Moraru, *Carbohydr. Res.*, 2005, **340**, 507.
- 40 G. Wei, X. Lv and Y. Du, *Carbohydr. Res.*, 2008, **343**, 3096.
- 41 S. Narayanaperumal, R. César da Silva, J. L. Monteiro, A. G. Corrêa and M. W. Paixão, *J. Braz. Chem. Soc.*, 2012, **23**, 1982.
- 42 H. Marzag, G. Robert, M. Dufies, K. Bougrin, P. Auberger and R. Benhida, *Ultrason. Sonochem.*, 2015, **22**, 15.
- 43 J. B. Laursen, L. Petersen and K. J. Jensen, *Org. Lett.*, 2001, **3**, 687.

- 44 M. R. Rasmussen, M. H. Marqvorsen, S. K. Kristensen and H. H. Jensen, *J. Org. Chem.*, 2014, **79**, 11011.
- 45 G. S. Shetye, N. Singh, C. Jia, C. D. Nguyen, G. Wang and Y. Y. Luk, *Chembiochem*, 2014, **15**, 1514.
- 46 M. M. Mukherjee, N. Basu and R. Ghosh, *RSC Adv.*, 2016, **6**, 105589.
- 47 T. Mukaiyama, K. Matsubara and M. Hora, *Synthesis*, 1994, 1368.
- 48 S. C. Ranade, S. Kaeothip and A. V. Demchenko, *Org. Lett.*, 2010, **12**, 5628.
- 49 P. O. Miranda, I. Brouard, J. I. Padrón and J. Bermejo, *Tetrahedron Lett.*, 2003, **44**, 3931.