

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: A. M. R. Hall, R. Broomfield-Tagg, M. Camilleri, D. Carbery, A. Codina, D. Whittaker, S. Coombes, J. P. Lowe, U. Hintermair and U. Hintermair, *Chem. Commun.*, 2017, DOI: 10.1039/C7CC07059D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Journal Name

COMMUNICATION

Online Monitoring of a Photocatalytic Reaction by Real-time High Resolution FlowNMR Spectroscopy

Andrew M. R. Hall,^a Rachael Broomfield-Tagg,^b Matthew Camilleri,^a David R. Carbery,^{b*} Anna Codina,^c David T. E. Whittaker,^d Steven Coombes,^d John P. Lowe,^b and Ulrich Hintermair^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x
www.rsc.org/

We demonstrate how FlowNMR spectroscopy can readily be applied to investigate photochemical reactions that require sustained input of light and air to yield mechanistic insight under realistic conditions. The Eosin Y mediated photo-oxidation of N-allylbenzylamine is shown to produce imines as primary reaction products from which undesired aldehydes form after longer reaction times. Facile variation of reaction conditions during the reaction in flow allows for probe experiments that give information about the mode of action of the photocatalyst.

Photochemical reactions are an important part of the modern synthetic chemists' repertoire due to the unique chemistry occurring in the excited state.¹⁻⁵ Visible light photo(redox)catalysis is of particular current interest due to the potential for utilization of sunlight as sustainable energy source in chemical synthesis.⁶⁻¹¹ In order to gain an understanding of the mechanisms involved in these reactions it is necessary to have a means of producing high quality kinetic data under realistic conditions. Spectroscopic techniques such as Ultra-violet visible (UV-vis), Infrared (IR) and fluorescence spectroscopies have been widely used for monitoring photochemical reactions, however, these techniques are only able to provide limited structural information about reaction species and require calibration before use.^{1, 12-15} Mass Spectrometry represents a complimentary technique with higher sensitivity and resolution, but structural information may still be limited for novel compounds where fragmentation patterns are unknown, and calibration is required before quantitative results may be attained.^{16, 17} Although less sensitive, Nuclear Magnetic Resonance (NMR) spectroscopy provides a wealth of structural information over a wide detection range and does not require external calibration. However, conventional NMR

techniques are incompatible with photochemical reactions due to the difficulties in irradiating the sample once it is inside the magnet.^{13, 14}

For this reason, most NMR monitoring of photochemical reactions is performed off-line, but the delays and sample work-up procedures between reaction and analysis often entail compositional changes. *In-situ* approaches including setups where light is guided from an external source to the sample inside the spectrometer using mirrors or fibre-optic cables,^{18, 19} special sample tubes with miniature LED light sources inside the spectrometer^{20, 21} and modified NMR probes have been developed^{13, 14} but these require custom-made equipment and are the domain of specialists. In addition, these setups do not allow for mixing of the sample or changes in reaction conditions,²² and often deliver light at one point only.^{14, 19-21} The lack of control of light intensity across the sample means that recreating realistic conditions is difficult, which may result in different rates or/and mechanisms for reactions that are limited by photon density. The use of NMR as online monitoring technique has been known for some years, primarily as a detection method in HPLC-NMR coupled analysis,^{23, 24} but only recently gained interest as a method for real-time reaction monitoring. This has been spurred on by the recent commercialisation of a number of dedicated FlowNMR monitoring systems²⁵⁻²⁸ accompanied by several reports detailing general considerations for their use.^{26, 29, 30} Online FlowNMR techniques utilize an external reaction vessel, situated outside of the influence of the magnetic field, with sample continuously pumped into the spectrometer through a dedicated NMR flow tube located within a standard NMR probe before returning to the reaction vessel (Figure 1).^{25, 29} This setup permits full control over the reaction conditions during the analysis, including mixing, addition of reagents, temperature regulation as well as irradiation of the sample.²² Advanced solvent suppression techniques allow the use of non-deuterated reaction solvents without compromising data quality, reducing cost and avoiding unwanted isotope effects.³¹ Two recent papers have explored the possibility of utilizing online NMR reaction monitoring to study photochemical reactions, with one study investigating the photochemical degradation of environmental

^a Centre for Sustainable Chemical Technologies, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom.

^b Department of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom.

^c Bruker UK Ltd., Banner Lane, Coventry CV4 9GH, United Kingdom.

^d AstraZeneca, Charter Way, Macclesfield SK10 2NA, United Kingdom.

* Corresponding authors: D.Carbery@bath.ac.uk, U.Hintermair@bath.ac.uk

† Electronic Supplementary Information (ESI) including experimental details and additional figures available online: See DOI: 10.1039/x0xx00000x







Figure 1: FlowNMR setup for monitoring photochemical reactions (not to scale).

pollutants,¹³ and another using a low-field NMR spectrometer for analysis of products exiting from a photochemical flow reactor.³² Neither study reports reaction kinetics or mechanism, and to the best of our knowledge, no examples of high-resolution FlowNMR reaction monitoring for kinetic and mechanistic investigations of photochemical reactions have thus far been reported.

Examples of visible light photocatalysis are abundant in nature, forming the basis of many key biological processes.^{1, 2, 5} Flavins, based on a tricyclic isoalloxazine ring system, are an important class of natural photocatalysts.^{2, 33-35} Synthetic fluorescent dyes mimicking the core structure of flavins have been developed as staining agents for biological systems, including compounds such as Fluorescein, Eosin and Rose Bengal. In the presence of visible light and air, Flavin and Eosin Y have been found to catalyse reactions of secondary amines to give a mixture of inter- and intramolecular oxidation products (Scheme 1).³⁶ The reaction proceeds at room temperature in acetonitrile solution under white or green light irradiation in air (Scheme 1). Sampling off-line ¹H NMR analysis of the reaction showed the major products to be oxidation product **2** and intermolecular coupling product **3** (Figure 2). Small amounts of benzaldehyde were also observed throughout the reaction as an undesired by-product. Even with utmost care, data from off-line sampling was always scattered and difficult to reproduce due to slight variations between different samples and the need for work-up before the analysis (Figure 2). Monitoring the reaction under the same conditions by online ¹H FlowNMR spectroscopy however (Figure 1) yielded smooth and highly reproducible concentration profiles of all components from a single experiment (Figure 2). The ~30 second time lapse between leaving the continuously illuminated vessel and reaching the NMR probe was sufficient to avoid adverse effects on NMR data acquisition by photo-generated radicals, and the returning aliquot resumed turnover once returned to the illuminated vessel. The data obtained revealed a steady consumption of the N-allylbenzylamine starting material with corresponding



Scheme 1: Structure of Eosin Y and photocatalytic oxidation of allylic amines.




Figure 2: Off-line and on-line FlowNMR reaction profiles of N-allylbenzylamine (**1**) in MeCN (6.4 mM) at 20°C in the presence of Eosin Y catalyst (1 mol%) under green LED illumination (633 μ W) to form **2**, **3** and **4**. (On-line: 100 mL, 4 mL min⁻¹ flowrate, WET solvent suppression, 1.64 s acquisition time, 3 s relaxation delay, 12 scans. Off-line: 300 mL, 3 s acquisition time, 1 s relaxation delay, 16 scans. 20 mL samples were periodically withdrawn from the reaction mixture and concentrated under reduced pressure before dissolving in CDCl₃ for analysis). See the supporting information for details.

increase in concentration of products **2** and **3** in a 1:1 ratio at virtually identical rates. The rate of the reaction was constant until about 65% conversion, indicating pseudo-zero order kinetics in **1** under the conditions applied. Interestingly, no benzaldehyde formation was observed by FlowNMR during the initial stages of the reaction; the associated peaks did not appear until about 2 hours into the reaction, corresponding to >80% substrate consumption. This clearly showed benzaldehyde to be a secondary reaction product formed from imines **2** and **3** rather than from amine **1**, which appeared much more pronounced in the off-line reaction monitoring data due to degradation between sampling and data acquisition. Under the conditions applied, substrate conversion reached completion after ~4 hours, after which the concentration of benzaldehyde continued to increase at the expense of both **2** and **3**, although degradation of allylphenylimine **2** was slightly faster than the more stable benzylphenylimine **3**.

Increasing the concentration of **1** seven-fold resulted in a lengthening of the reaction time, again with constant rates during the first 6 hours up to ~65% conversion indicative of saturation kinetics (Figure S2). Over these longer reaction times a number of other (currently unidentified) by-products were observed to form at low concentration, as evidenced by the appearance of a variety of small peaks in the ¹H spectra (Figure S3). Sum integration of the aromatic region remained constant over the course of the reaction, however, demonstrating that the reaction mass-balance was maintained (Figure S4).

Having access to the reaction vessel during FlowNMR experiments means that the mode and power of illumination may easily be changed while ensuring even light input across the sample, as a way of investigating whether the reaction is limited by the availability of photons. When using a more powerful light source (250 mW halogen lamp) the reaction became much faster, reaching completion in just 30 minutes at 0°C (Figure 3), with fewer by-products formed than with lower power LED light sources that required longer reaction






Figure 1: ^1H FlowNMR reaction profiles of N-allylbenzylamine (**1**) in MeCN (6.4 mM) at 0°C in the presence of Eosin Y catalyst (1 mol%) under halogen light illumination (250 mW) to form **2**, **3** and **4** (200 mL, 4 mLmin^{-1} flowrate, WET solvent suppression, 4 s acquisition time, 1 s relaxation delay, 16 scans).

times. Since the Eosin Y exhibits a sharp absorption peak around 535 nm with minimal absorption at other wavelengths emitted by the halogen lamp (Figure S5), this difference in reaction rate is attributed to the greater light intensity $\sim 535\text{ nm}$ rather than the presence of higher energy UV photons. This was confirmed by performing the reaction using a UV filter, which yielded a virtually identical reaction rate (Figure S6). The formation of benzaldehyde was not significantly affected by light intensity, proceeding at roughly the same rate regardless of light source once appreciable amounts of **2** and **3** had formed in the reaction mixture.

Varying light intensity further revealed the reaction to be photon-limited under typical conditions, with higher lamp powers yielding increased initial reaction rates (Table S1 & Figure S7). While under photon-limited conditions, **2** and **3** appeared to form in parallel (Figure 2), under high illumination conditions **2** formed as soon as the light was switched on, but formation of **3** occurred only after an induction period of 2–3 minutes (Figure 3). This observation suggests the formation of **3** to proceed via reaction of **2** with **1** activated by




Figure 4: ^1H FlowNMR reaction profiles of N-allylbenzylamine (**1**) in MeCN (6.4 mM) at 20°C in the presence of Eosin Y catalyst (1 mol%) under chopped green LED illumination ($633\text{ }\mu\text{W}$) to form **2**, **3** and **4** (100 mL, 4 mLmin^{-1} flowrate, WET solvent suppression, 1.46 s acquisition time, 3 s relaxation delay, 12 scans).





Figure 5: ^1H FlowNMR reaction profiles of N-allylbenzylamine (**1**) in MeCN (6.4 mM) at 20°C in the presence of Eosin Y catalyst (1 mol%) under chopped green LED illumination ($633\text{ }\mu\text{W}$) and different atmospheres (argon to dry air) to form **2**, **3** and **4** (50 mL, 4 mLmin^{-1} flowrate, WET solvent suppression, 1.46 s acquisition time, 3 s relaxation delay, 16 scans).

the excited photocatalyst system (most likely an α -aminoradical),³⁶ leading to coupling and loss of two allylic groups. No allylic signals other than those from **1** were ever observed by NMR, but spiking a reaction mixture with allylamine showed it to rapidly disappear from the spectra as soon as the reaction was restarted by illumination (Figure S8), showing likely by-products of the formation of **3** and **4** to be undetectable under the conditions applied.

To investigate the nature of the photo-initiated radical reaction system, variations of atmosphere and illumination were undertaken during the reaction. A chopped illumination experiment performed in air (Figure 4) revealed that no further reaction took place in the absence of light after initial irradiation, showing the photo-generated reactive species to have short lifetimes and only operate under sustained input of photons. Product distribution profiles were identical to experiments performed under continuous illumination. Altering reaction atmospheres during light variation showed that some product formed very slowly in an illuminated reaction mixture under dry argon, and no reaction took place when air was introduced but the light switched off (Figure 5). Only when illuminated under aerobic conditions product formation occurred, and the reaction stalled immediately when the light was switched off again.

The formation of benzaldehyde **4** seemed to occur only under illumination in air (Figure 5). Test reactions showed the imine products **2** and **3** to be resistant to hydrolysis in the absence of Eosin Y,³⁷ and addition of water to an anhydrous photocatalytic FlowNMR



Scheme 2: Reaction network based on experimentally observed product and by-product formation profiles, and conditions.



experiment did not affect the amount of benzaldehyde formation (Figure S9). Thus, product degradation to the undesired aldehydes must be a parallel photo-initiated process rather than a simple hydrolysis reaction. This may proceed via the formation of the corresponding iminoradicals,³⁶ or due to the presence of singlet oxygen formed upon excitation of Eosin Y.³⁸ Combining all these observations leads to the proposed reaction network shown in Scheme 2.

We have shown how FlowNMR spectroscopy can be readily applied to investigate photochemical reactions under realistic conditions to provide valuable mechanistic insight. Substrate consumption profiles showed the reaction to operate under saturation kinetics (photon starvation), and product formation profiles revealed their relative order of formation. Variation of light intensities and reaction atmospheres in conjunction with chopped illumination experiments gave insight into the mode of action of the Eosin Y photocatalyst system, and showed aldehyde formation to occur from the imines via a parasitic photocatalytic pathway. This information, not easily accessible by alternative reaction monitoring techniques, will allow swift optimisation of reaction conditions and assist the design of improved photocatalytic systems in the future.

Conflicts of interest

A.C. is an employee of Bruker UK Ltd., manufacturer and supplier of NMR hard- and software solutions that have been used in this research. The other authors declare no competing financial interest.

Acknowledgements

This work was supported by a Research Grant from the Royal Society (Y0603), the EPSRC Centre for Doctoral Training in Sustainable Chemical Technologies (EP/L016354/1), the Dynamic Reaction Monitoring Facility at the University of Bath (EP/P001475/1), Bruker UK Ltd., and AstraZeneca. U.H. acknowledges the Centre for Sustainable Chemical Technologies for a Whorrod Research Fellowship. The authors would like to thank Joshua Tibbets, Dr Catherine Lyall and Dr Emma Emanuelsson from the University of Bath for support and assistance with this project.

Notes and references

‡ Total volume of FlowNMR apparatus = 3.7 mL. For a 100 mL reaction volume the sample spends 4% of the reaction time outside the reaction vessel (on average).

1. B. König, *Chemical Photocatalysis*, De Gruyter, 2013.
2. P. F. Heelis, *Chem. Soc. Rev.*, 1982, **11**, 15-39.
3. N. Hoffmann, *Chem. Rev.*, 2008, **108**, 1052-1103.
4. Y. Inoue, *Chem. Rev.*, 1992, **92**, 741-770.
5. K. Szacilowski, W. Macyk, A. Drzewiecka-Matuszek, M. Brindell and G. Stochel, *Chem. Rev.*, 2005, **105**, 2647-2694.
6. V. Balzani, A. Credi and M. Venturi, *ChemSusChem*, 2008, **1**, 26-58.

7. J.-T. Li, J.-H. Yang, J.-F. Han and T.-S. Li, *Green Chem.*, 2003, **5**, 433. DOI: 10.1039/C7CC07059D
8. M. Oelgemöller, C. Jung and J. Mattay, *Pure Appl. Chem.*, 2007, **79**.
9. C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322-5363.
10. J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102-113.
11. T. P. Yoon, M. A. Ischay and J. Du, *Nat. Chem.*, 2010, **2**, 527-532.
12. G. L. Closs and R. J. Miller, *J. Am. Chem. Soc.*, 1981, **103**, 3586-3588.
13. L. Blumkin, R. Dutta Majumdar, R. Soong, A. Adamo, J. P. D. Abbatt, R. Zhao, E. Reiner and A. J. Simpson, *Environ. Sci. Technol.*, 2016, **50**, 5506-5516.
14. G. E. Ball, in *Spectrosc. Prop. Inorg. Organomet. Compd.*, RSC, 2010, vol. 41, pp. 262-287.
15. B. Roig, E. Touraud and O. Thomas, *Spectrochim. Acta Mol. Biomol. Spectrosc.*, 2002, **58**, 2925-2930.
16. J. Luo, A. G. Oliver and J. Scott McIndoe, *Dalton Trans.*, 2013, **42**, 11312-11318.
17. R. Theron, Y. Wu, L. P. E. Yunker, A. V. Hesketh, I. Pernik, A. S. Weller and J. S. McIndoe, *ACS Catalysis*, 2016, **6**, 6911-6917.
18. T. F. Page, Jr., *Chem. Ind.*, 1969, 1462-1463.
19. A. Mills and C. O'Rourke, *J. Org. Chem.*, 2015, **80**, 10342-10345.
20. C. Feldmeier, H. Bartling, K. Magerl and R. M. Gschwind, *Angew. Chem. Int. Ed.*, 2015, **54**, 1347-1351.
21. C. Feldmeier, H. Bartling, E. Riedle and R. M. Gschwind, *J. Magn. Reson.*, 2013, **232**, 39-44.
22. D. A. Foley, A. L. Dunn and M. T. Zell, *Magn. Reson. Chem.*, 2016, **54**, 451-456.
23. K. Albert, in *On-Line LC-NMR And Related Techniques*, John Wiley & Sons, Ltd, 2003, pp. 1-22.
24. K. Albert and E. Bayer, *TrAC, Trends Anal. Chem.*, 1988, **7**, 288-293.
25. D. A. Foley, E. Bez, A. Codina, K. L. Colson, M. Fey, R. Krull, D. Piroli, M. T. Zell and B. L. Marquez, *Anal. Chem.*, 2014, **86**, 12008-12013.
26. D. A. Foley, M. T. Zell, B. L. Marquez and A. Kaerner, *Pharm. Technol.*, 2011, **11**, S19.
27. M. Khajeh, M. A. Bernstein and G. A. Morris, *Magn. Reson. Chem.*, 2010, **48**, 516-522.
28. M. V. Silva Elipse and R. R. Milburn, *Magn. Reson. Chem.*, 2016, **54**, 437-443.
29. A. M. R. Hall, J. C. Chouler, A. Codina, P. T. Gierth, J. P. Lowe and U. Hintermair, *Catal. Sci. Tech.*, 2016, **6**, 8406-8417.
30. N. Zientek, K. Meyer, S. Kern and M. Maiwald, *Chem. Ing. Tech.*, 2016, **88**, 698-709.
31. S. H. Smallcombe, S. L. Patt and P. A. Keifer, *J. Magn. Reson. A*, 1995, **117**, 295-303.
32. N. Emmanuel, C. Mendoza, M. Winter, C. R. Horn, A. Vizza, L. Dreesen, B. Heinrichs and J.-C. M. Monbaliu, *Org. Process Res. Dev.*, 2017, DOI: 10.1021/acs.oprd.7b00212.
33. D. P. Hari and B. König, *Org. Lett.*, 2011, **13**, 3852-3855.
34. Y. Pan, S. Wang, C. W. Kee, E. Dubuisson, Y. Yang, K. P. Loh and C.-H. Tan, *Green Chem.*, 2011, **13**, 3341-3344.
35. N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075-10166.
36. A. T. Murray, M. J. Dowley, F. Pradaux-Caggiano, A. Baldansuren, A. J. Fielding, F. Tuna, C. H. Hendon, A. Walsh, G. C. Lloyd-Jones, M. P. John and D. R. Carbery, *Angew. Chem. Int. Ed. Engl.*, 2015, **54**, 8997-9000.
37. A. Murray, Doctor of Philosophy, University of Bath, 2015.
38. F. Amat-Guerri, M. M. C. López-González, R. Martínez-Utrilla and R. Sastre, *J. Photochem. Photobiol. A: Chem.*, 1990, **53**, 199-210.

